

SELECTIVE NPY (Y5) ANTAGONISTS5 Background Of The InventionSub
B1

10 This application claims priority of and is a continuation-in-part of U.S. Serial No. 09/343,635, filed June 30, 1999, U.S. Serial No. 09/343,633, filed June 30, 1999, and U.S. Serial No. 09/343,993, filed June 30, 1999, the contents of all of which are hereby incorporated by reference into the subject application.

15 Throughout this application, various references are referred to within parentheses. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains. Full bibliographic citation for these references may be
20 found at the end of this application, preceding the claims.

25 The peptide neurotransmitter neuropeptide Y (NPY) is a 36 amino acid member of the pancreatic polypeptide family with widespread distribution throughout the mammalian nervous system (Dumont et al., 1992). The family includes the pancreatic polypeptide (PP), synthesized primarily by endocrine cells in the pancreas; peptide YY (PYY), synthesized primarily by endocrine cells in the gut; and
30 NPY, synthesized primarily in neurons (Michel, 1991; Dumont et al., 1992; Wahlestedt and Reis, 1993). All pancreatic polypeptide family members share a compact structure involving a "PP-fold" and a conserved C-terminal hexapeptide ending in Tyr³⁶ (or Y³⁶ in the single letter
35 code). The striking conservation of Y³⁶ has prompted the

reference to the pancreatic polypeptides' receptors as "Y-type" receptors (Wahlestedt et al., 1987), all of which are proposed to function as seven transmembrane-spanning G protein-coupled receptors (Dumont et al., 1992).

5

NPY and its relatives elicit a broad range of physiological effects through activation of at least five G protein-coupled receptor subtypes known as Y1, Y2, Y3, Y4 (or PP), and the "atypical Y1". While the Y1, Y2, Y3, and Y4 (or PP) receptors were each described previously in both radioligand binding and functional assays, the "atypical Y1" receptor is unique in that its classification is based solely on feeding behavior induced by various peptides including NPY.

10

15

The role of NPY in normal and abnormal eating behavior, and the ability to interfere with NPY-dependent pathways as a means to appetite and weight control, are areas of great interest in pharmacological and pharmaceutical research (Sahu and Kalra, 1993; Dryden et al., 1994). NPY is considered to be the most powerful stimulant of feeding behavior yet described (Clark et al., 1984; Levine and Morley, 1984; Stanley and Leibowitz, 1984). The stimulation of feeding behavior by NPY is thought to occur primarily through activation of the hypothalamic "atypical Y1" receptor. For example, direct injection of NPY into the hypothalamus of satiated rats can increase food intake up to 10-fold over a 4-hour period (Stanley et al., 1992). Similar studies using other peptides has resulted in a pharmacologic profile for the "atypical Y1" receptor according to the rank order of potencies of peptides in stimulating feeding behavior as follows: NPY₂₋₃₆ NPY ~ PYY ~ [Leu³¹,Pro³⁴]NPY > NPY₁₃₋₃₆ (Kalra et al., 1991; Stanley et al., 1992). The profile is similar to that of

20

25

30

a Y1-like receptor except for the anomalous ability of NPY₂₋₃₆ to stimulate food intake with potency equivalent or better than that of NPY. A subsequent report in J. Med. Chem. by Balasubramaniam and co-workers (1994) showed that feeding can be regulated by [D-Trp³²]NPY. While this peptide was presented as an NPY antagonist, the published data at least in part support a stimulatory effect of [D-Trp³²]NPY on feeding. In contrast to other NPY receptor subtypes, the "feeding" receptor has never been characterized for peptide binding affinity in radioligand binding assays.

This problem has been addressed by cloning rat and human cDNAs which encode a single receptor protein, referred to herein as Y5, whose pharmacologic profile links it to the "atypical Y1" receptor. The identification and characterization of a single molecular entity which explains the "atypical Y1" receptor allows the design of selective drugs which modulate feeding behavior (WO 96/16542). It is important to note, though, that any credible means of studying or modifying NPY-dependent feeding behavior must necessarily be highly selective, as NPY interacts with multiple receptor subtypes, as noted above (Dumont et al., 1992).

As used in this invention, the term "antagonist" refers to a compound which binds to, and decreases the activity of, a receptor in the presence of an agonist. In the case of a G-protein coupled receptor, activation may be measured using any appropriate second messenger system which is coupled to the receptor in a cell or tissue in which the receptor is expressed. Some specific but by no means limiting examples of well-known second messenger systems are adenylate cyclase, intracellular calcium mobilization,

ion channel activation, guanylate cyclase, and inositol phospholipid hydrolysis. Conversely, the term "agonist" refers to a compound which binds to, and increases the activity of, a receptor as compared with the activity of the receptor in the absence of any agonist.

In order to test compounds for selective binding to the human Y5 receptor the cloned cDNAs encoding both the human and rat Y2 and Y4 (or PP) receptors have been used. The human and rat Y5 receptors are described in coassigned U.S. Patent No. 5,602,024 and in PCT International Application PCT/US95/15646, published June 6, 1996, as WO 96/16542, the contents of which are hereby incorporated by reference into this application. The human and rat Y2 receptors are described in coassigned U.S. Patent No. 5,545,549 and in PCT International Application PCT/US95/01469, published August 10, 1995, as WO 95/21245, the contents of which are hereby incorporated by reference into this application. The human and rat Y4 receptors are described in coassigned U.S. Patent No. 5,516,653 and in PCT International Application PCT/US94/14436, published July 6, 1995, as WO 95/17906, the contents of which are hereby incorporated by reference into this application. The Y1 receptor has been cloned from a variety of species including human, rat and mouse (Larhammar et al., 1992; Herzog et al., 1992; Eva et al., 1990; Eva et al., 1992).

Using the NPY-Y5-selective antagonist CGP 71683A, it was demonstrated recently that food intake in free-feeding and energy-derived lean rats is mediated by the Y5 receptor (Criscione et al., 1998). CGP 71683A has high affinity for the cloned rat NPY-Y5 receptor subtype, but 1,000-fold lower affinity for the cloned rat NPY-Y1, Y2, and Y4

5


receptors. Examples of additional NPY-Y5-selective compounds are described in WO 97/20823, WO 98/35957, and WO 98/35944.

5 In different embodiments of this invention the synthesis of novel bicyclic compounds and tricyclic compounds which bind selectively to the cloned human Y5 receptor, compared to the other cloned human NPY receptors, and inhibit the activation of the cloned human Y5 receptor as measured in
10 in vitro assays is disclosed. The in vitro receptor binding and activation assays described hereinafter were performed using various cultured cell lines, each transfected with and expressing only a single Y-type receptor.

15 In addition, the compounds of the present invention may also be used to treat abnormal conditions such as feeding disorders (obesity and bulimia nervosa), sexual/reproductive disorders, depression, epileptic
20 seizure, hypertension, cerebral hemorrhage, congestive heart failure, sleep disturbances, or any condition in which antagonism of a Y5 receptor may be beneficial.

Summary Of The InventionA. Tricyclic Compounds

5 This invention provides a compound having the structure:



10 wherein each R_1 is independently H, F, Cl, Br, -CN, -OH, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, perfluoroalkyl, polyfluoroalkyl, aminoalkyl, or straight chained or branched C₁-C₇ alkyl;

15 wherein R₅ is independently H; or straight chained or branched C₁-C₇ alkyl;


wherein R₆ is independently H; or straight chained or branched C₁-C₇ alkyl;

20 wherein B is O, NH or S;

wherein X is CHR₅, O or NR₅;

25 wherein each n independently is an integer from 0 to 6 inclusive;

wherein R_8 is



5

wherein Y is C or N;

wherein R_7 is independently straight chained or branched C_1 - C_7 alkyl;

10


wherein R_9 is independently H; or straight chained or branched C_1 - C_4 alkyl;

8

wherein R_{10} is independently H; or straight chained or branched C_1 - C_4 alkyl;

wherein R_{11} is

5



;

wherein R_{12} is H, straight chained or branched C_1 - C_7 alkyl, $(\text{CH}_2)_n\text{OR}_{17}$, or $\text{O}(\text{CH}_2)_u\text{OR}_{17}$; provided that when X is O, R_{12} cannot be methyl;

10

wherein R_{13} is independently H; $-(\text{CH}_2)_u\text{OR}_5$; $-(\text{CH}_2)_t\text{CONR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{NR}_5\text{COR}_5$; $-(\text{CH}_2)_t\text{COR}_7$; $-(\text{CH}_2)_t\text{CO}_2\text{R}_5$; $-(\text{CH}_2)_u\text{NR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{CN}$; straight chained or branched C_1 - C_7 alkyl; C_1 - C_7 alkyl in which the C_2 - C_7 atoms may be optionally substituted with one or more F or Cl; C_3 - C_7 cycloalkyl- C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl; phenyl or C_1 - C_6 phenylalkyl; wherein the phenyl or C_1 - C_6 phenylalkyl may be substituted with one or more of F, Cl, -CN, -NO₂, -NR₅R₆, -SO₂R₅, $-(\text{CH}_2)_n\text{COR}_7$, $-(\text{CH}_2)_n\text{OR}_5$, $-(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$, $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{SO}_2\text{NR}_5\text{R}_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

15

20

25

or R_{12} and R_{13} together with the amide linkage to which they are attached are pyrrolidinonyl, piperidonyl, or oxazolidinonyl; provided that when X is O, R_{12} and R_{13} cannot be oxazolidinonyl;

30

wherein R_{14} is H; straight chained or branched C_1-C_4 alkyl; F; or $-(CH_2)_rOR_5$;

wherein R_{15} is H, straight chained or branched C_1-C_4 alkyl, or F;

with the proviso that when R_{14} is -OH, R_{15} cannot be F;

wherein R_{16} is NR_3R_4 , perfluoroalkyl, unsubstituted straight chained or branched C_1-C_7 alkyl, substituted straight chained or branched C_2-C_7 alkyl, wherein the C_2-C_7 alkyl may be substituted with one or more of F, Cl, -CN, $-NR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nOCF_3$, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2-C_7 alkenyl or alkynyl, or C_3-C_7 cycloalkyl or cycloalkenyl; C_3-C_7 cycloalkyl or cycloalkenyl; phenyl, heteroaryl, or C_1-C_7 phenylalkyl, wherein the phenyl, heteroaryl, or C_1-C_7 phenylalkyl may be substituted with one or more of F, Cl, Br, -CN, $-NO_2$, $-NR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, ethylenedioxy, methylenedioxy, straight chained or branched C_1-C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2-C_7 alkenyl or alkynyl, or C_3-C_7 cycloalkyl or cycloalkenyl; quinolinyl, 1-naphthyl, 2-naphthyl, or 2,1,3-benzothiadiazolyl; wherein the quinolinyl, 1-naphthyl, 2-naphthyl or 2,1,3-benzothiadiazolyl may be substituted with one or more of F, Cl, Br, -CN, $-NO_2$, $-NR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, ethylenedioxy, methylenedioxy, straight chained or branched C_1-C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

with the proviso that when X is O and R_8 is $NR_9(CH_2)_uO(CH_2)_uNR_{10}R_{11}$, R_{16} cannot be methyl;

5

wherein R_3 is independently H; $-(CH_2)_uOR_5$; $-(CH_2)_tCONR_5R_6$; $-(CH_2)_uNR_5COR_5$; $-(CH_2)_tCOR_7$; $-(CH_2)_tCO_2R_5$; $-(CH_2)_uNR_5R_6$; $-(CH_2)_uCN$; straight chained or
 10 branched C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl or cycloalkenyl; phenyl, C_1 - C_6 phenylalkyl, or C_1 - C_6 heteroarylalkyl; wherein the phenyl, C_1 - C_6 phenylalkyl or C_1 - C_6 heteroarylalkyl may be substituted with one or more of F, Cl, Br, $-CN$, $-NO_2$,
 15 $-NR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl;

20

wherein R_4 is independently H; $-(CH_2)_uOR_5$; $-(CH_2)_tCONR_5R_6$; $-(CH_2)_uNR_5COR_5$; $-(CH_2)_tCOR_7$; $-(CH_2)_tCO_2R_5$; $-(CH_2)_uNR_5R_6$; $-(CH_2)_uCN$; straight chained or
 25 branched C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl or cycloalkenyl; phenyl or C_1 - C_6 phenylalkyl; wherein the phenyl or C_1 - C_6 phenylalkyl may be substituted with one or more of F, Cl, Br, $-CN$, $-NO_2$, $-NR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$,
 30 straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl;

or R_3 and R_4 taken together with the nitrogen atom to which they are attached are 1-azetidiny1, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl, wherein the 1-azetidiny1, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl is substituted with one or more of F, -CN, $-(CH_2)_nNR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl, or phenyl or heteroaryl; wherein if $-(CH_2)_nNR_5R_6$, $-(CH_2)_nOR_5$, or $-(CH_2)_nNR_5COR_5$ are in the 2-position, then n is not 0; wherein the phenyl or heteroaryl may be substituted with one or more of F, Cl, Br, -CN, $-NO_2$, $-NR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl;

or R_3 and R_4 taken together with the nitrogen atom to which they are attached are morpholinyl, thiomorpholinyl, [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or [1,4]diazepanyl, wherein the morpholinyl, thiomorpholinyl, [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or [1,4]diazepanyl is optionally substituted with straight chained or branched C_1 - C_5 alkyl or $(CH_2)_tOR_5$; and wherein the nitrogen atom of the piperazinyl or [1,4]diazepanyl ring may be optionally substituted with $-(CH_2)_uOR_5$; $-COR_5$; straight chained or branched C_1 - C_5 alkyl; or phenyl; wherein the phenyl may be substituted with one or more of F, Cl, Br, -CN, $-NO_2$, $-NR_5R_6$, $-(CH_2)_nOR_5$, straight chained or

branched C₁-C₃ alkyl,¹² perfluoroalkyl,
polyfluoroalkyl, or aminoalkyl;

wherein R₁₇ is H, straight chained or branched C₁-C₄ alkyl,
perfluoroalkyl, or polyfluoroalkyl;

wherein R₁₉ is (CH₂)_nOR₅, NR₅R₆, phenyl, or heteroaryl,
wherein the phenyl or heteroaryl may be substituted with
one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -
(CH₂)_nNR₅COR₅, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅,
- (CH₂)_nCONR₅R₆, - (CH₂)_nCO₂R₅, - (CH₂)_nSO₂NR₅R₆, ethylenedioxy,
methylenedioxy, straight chained or branched C₁-C₇ alkyl,
perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight
chained or branched C₂-C₇ alkenyl or alkynyl, or C₃-C₇
cycloalkyl or cycloalkenyl;

wherein m is 0 or 1;

wherein each p independently is an integer from 0 to 2
inclusive;

wherein each r independently is an integer from 0 to 3
inclusive;

wherein each s independently is an integer from 1 to 6
inclusive;

wherein t is an integer from 1 to 4 inclusive;

wherein each u independently is an integer from 2 to 4
inclusive;

wherein v is 1 or 2;

13

with the proviso that when v is 2, m is 0;


wherein z is an integer from 2 to 7;

5 or a pharmaceutically acceptable salt thereof.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
22

B. Bicyclic Compounds

This invention provides a compound having the structure:



wherein Y is O, S or NH;

wherein each R_{14} independently is H, F, Cl, Br, -CN, -OH, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, -SO₂C₆H₅, -SO₂NR₅R₆, -C₆H₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, ethylenedioxy, methylenedioxy, perfluoroalkyl, polyfluoroalkyl, aminoalkyl, or straight chained or branched C₁-C₇ alkyl; or phenyl, heteroaryl, or C₁-C₇ phenylalkyl, wherein the phenyl, heteroaryl, or C₁-C₇ phenylalkyl may be substituted with one or more of F, Cl, Br, -CF₃, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, or straight chained or branched C₁-C₄ alkyl; provided that if one R_{14} is phenyl, heteroaryl or C₁-C₇ phenylalkyl, the other R_{14} is H;

wherein each R_1 independently is H, F, Cl, Br, -CN, -OH, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, -SO₂C₆H₅, -SO₂NR₅R₆, -C₆H₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, ethylenedioxy, methylenedioxy, perfluoroalkyl, polyfluoroalkyl, aminoalkyl, or straight chained or branched C₁-C₇ alkyl; or phenyl, heteroaryl, or C₁-C₇ phenylalkyl, wherein the phenyl, heteroaryl, or C₁-C₇ phenylalkyl may be substituted with one or more of F, Cl, Br, -CF₃, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, or straight chained or branched C₁-C₄ alkyl;

wherein R_2 is H, straight chained or branched C_1 - C_4 alkyl,
 $-(CH_2)_tOR_5$, phenyl optionally substituted with one or more
 of of F, Cl, Br, $-CF_3$, $-CN$, $-NO_2$, $-NR_5R_6$, $-SO_2R_5$, -
 5 $(CH_2)_nOR_5$, or straight chained or branched C_1 - C_4 alkyl;



wherein R_5 is independently H; or straight chained or
 branched C_1 - C_7 alkyl;

10 wherein R_6 is independently H; or straight chained or
 branched C_1 - C_7 alkyl;

wherein each n independently is an integer from 0 to 6
 inclusive;

15

wherein R_8 is



20

provided that R_1 or R_{14} cannot be $-OH$, when R_8 is (ii);


wherein R_9 is independently H; or straight chained or
 branched C_1 - C_4 alkyl;

25

16

wherein R_{10} is independently H; or straight chained or branched C_1 - C_4 alkyl;

wherein R_{11} is



5

wherein R_{12} is H, straight chained or branched C_1 - C_7 alkyl; or $(\text{CH}_2)_n\text{OR}_{17}$;

10

wherein R_{13} is independently $-(\text{CH}_2)_u\text{OR}_5$; $-(\text{CH}_2)_t\text{CONR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{NR}_5\text{COR}_5$; $-(\text{CH}_2)_t\text{COR}_7$; $-(\text{CH}_2)_t\text{CO}_2\text{R}_5$; $-(\text{CH}_2)_u\text{NR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{CN}$; straight chained or branched C_1 - C_7 alkyl; C_1 - C_7 alkyl in which the C_2 - C_7 atoms may be optionally substituted with one or more F or Cl; C_3 - C_7 cycloalkyl- C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl; phenyl or C_1 - C_6 phenylalkyl; wherein the phenyl or C_1 - C_6 phenylalkyl may be substituted with one or more of F, Cl, $-\text{CN}$, $-\text{NO}_2$, $-\text{NR}_5\text{R}_6$, $-\text{SO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{COR}_7$, $-(\text{CH}_2)_n\text{OR}_5$, $-(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$, $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{SO}_2\text{NR}_5\text{R}_6$, or straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

15

20

or R_{12} and R_{13} together with the amide linkage to which they are attached are pyrrolidinonyl, piperidonyl, or oxazolidinonyl;

25

wherein R_7 is independently straight chained or branched C_1 - C_7 alkyl;

30

wherein R_{16} is NR_3R_4 , perfluoroalkyl, unsubstituted straight chained or branched C_1 - C_7 alkyl, substituted straight chained or branched C_2 - C_7 alkyl, wherein the C_2 - C_7 alkyl may be substituted with one or more of F, Cl, -CN, - NR_5R_6 , - SO_2R_5 , - $(CH_2)_nCOR_7$, - $(CH_2)_nOR_5$, - $(CH_2)_nCONR_5R_6$, - $(CH_2)_nNR_5COR_5$, - $(CH_2)_nCO_2R_5$, - $(CH_2)_nOCF_3$, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl; C_3 - C_7 cycloalkyl or cycloalkenyl; or phenyl, heteroaryl, or C_1 - C_7 phenylalkyl, wherein the phenyl, heteroaryl, or C_1 - C_7 phenylalkyl may be substituted with one or more of F, Cl, Br, I, -CN, - NO_2 , - NR_5R_6 , - $(CH_2)_nNR_5COR_5$, - SO_2R_5 , - $(CH_2)_nCOR_7$, - $(CH_2)_nOR_5$, - $(CH_2)_nCONR_5R_6$, - $(CH_2)_nCO_2R_5$, - $(CH_2)_nSO_2NR_5R_6$, ethylenedioxy, methylenedioxy, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl; quinoliny, 1-naphthyl, 2-naphthyl, or 2,1,3-benzothiadiazolyl; wherein the quinoliny, 1-naphthyl, 2-naphthyl or 2,1,3-benzothiadiazolyl may be substituted with one or more of F, Cl, Br, I, -CN, - NO_2 , - NR_5R_6 , - $(CH_2)_nNR_5COR_5$, - SO_2R_5 , - $(CH_2)_nCOR_7$, - $(CH_2)_nOR_5$, - $(CH_2)_nCONR_5R_6$, - $(CH_2)_nCO_2R_5$, - $(CH_2)_nSO_2NR_5R_6$, ethylenedioxy, methylenedioxy, or straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

wherein R_3 is independently H; - $(CH_2)_uOR_5$; - $(CH_2)_tCONR_5R_6$; - $(CH_2)_uNR_5COR_5$; - $(CH_2)_tCOR_7$; - $(CH_2)_tCO_2R_5$; - $(CH_2)_uNR_5R_6$; - $(CH_2)_uCN$; straight chained or branched C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl or cycloalkenyl; or phenyl, C_1 - C_6 phenylalkyl, or C_1 - C_6 heteroarylalkyl;

wherein the phenyl, C₁-C₆ phenylalkyl, or C₁-C₆ heteroarylalkyl may be substituted with one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, -(CH₂)_nCO₂R₅, -(CH₂)_nSO₂NR₅R₆,
 5 straight chained or branched C₁-C₇ alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C₂-C₇ alkenyl or alkynyl, or C₃-C₇ cycloalkyl or cycloalkenyl;

wherein R₄ is independently H; -(CH₂)_uOR₅; -(CH₂)_tCONR₅R₆; -(CH₂)_uNR₅COR₅; -(CH₂)_tCOR₇; -(CH₂)_tCO₂R₅; -(CH₂)_uNR₅R₆; -(CH₂)_uCN; straight chained or branched C₁-C₇ alkyl; straight chained or branched C₂-C₇ alkenyl or alkynyl; or C₃-C₇ cycloalkyl or cycloalkenyl;
 15 or phenyl or C₁-C₆ phenylalkyl; wherein the phenyl or C₁-C₆ phenylalkyl may be substituted with one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, -(CH₂)_nCO₂R₅, -(CH₂)_nSO₂NR₅R₆, straight chained or branched C₁-C₇ alkyl,
 20 perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C₂-C₇ alkenyl or alkynyl, or C₃-C₇ cycloalkyl or cycloalkenyl;

or R₃ and R₄ taken together with the nitrogen atom to which they are attached are 1-azetidiny, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl, wherein the 1-azetidiny, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl is substituted with one or more of F, -CN, -(CH₂)_nNR₅R₆, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, -(CH₂)_nCO₂R₅, straight chained or branched C₁-C₇ alkyl,
 30 perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C₂-C₇ alkenyl or alkynyl, or C₃-C₇ cycloalkyl or cycloalkenyl, or phenyl or heteroaryl; wherein if -(CH₂)_nNR₅R₆, -(CH₂)_nOR₅, or -(CH₂)_nNR₅COR₅ are

19

in the 2-position, then n is not 0; wherein the phenyl or heteroaryl may be substituted with one or more of F, Cl, Br, I, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, -(CH₂)_nCO₂R₅, -(CH₂)_nSO₂NR₅R₆, straight chained or branched C₁-C₇ alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C₂-C₇ alkenyl or alkynyl, or C₃-C₇ cycloalkyl or cycloalkenyl;

or R₃ and R₄ taken together with the nitrogen atom to which they are attached are morpholinyl, thiomorpholinyl, [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or [1,4]diazepanyl, wherein the morpholinyl, thiomorpholinyl, [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or [1,4]diazepanyl is optionally substituted with straight chained or branched C₁-C₅ alkyl or (CH₂)_tOR₅; and wherein the nitrogen atom of the piperazinyl or [1,4]diazepanyl ring may be optionally substituted with -(CH₂)_uOR₅; -COR₅; straight chained or branched C₁-C₅ alkyl; or phenyl; wherein the phenyl may be substituted with one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -(CH₂)_nOR₅, straight chained or branched C₁-C₃ alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

wherein R₁₇ is straight chained or branched C₁-C₄ alkyl, perfluoroalkyl, or polyfluoroalkyl;

wherein each p independently is an integer from 0 to 2 inclusive;

wherein each r independently is an integer from 0 to 3 inclusive;

wherein t is an integer from 1 to 4 inclusive;


wherein each u independently is an integer from 2 to 4 inclusive;

5 or a pharmaceutically acceptable salt thereof.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

C. Tricyclic Compounds

The invention provides a compound having the structure:



5 wherein each R_1 is independently H, F, Cl, Br, -CN, -OH, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, perfluoroalkyl, polyfluoroalkyl, aminoalkyl, or straight chained or branched C₁-C₇ alkyl;

10 wherein R₅ is independently H; or straight chained or branched C₁-C₇ alkyl;

wherein R₆ is independently H; or straight chained or branched C₁-C₇ alkyl;


15

wherein B is O, NH or S;

wherein X is S, SO or SO₂;

20 wherein each n independently is an integer from 0 to 6 inclusive;

wherein R_8 is



5

wherein Y is C or N;

wherein R_7 is independently straight chained or branched
 10 C_1 - C_7 alkyl;


wherein R_9 is independently H; or straight chained or
 branched C_1 - C_4 alkyl;

23

wherein R_{10} is independently H; or straight chained or branched C_1 - C_4 alkyl;

wherein R_{11} is

5



wherein R_{12} is H, straight chained or branched C_1 - C_7 alkyl, $(\text{CH}_2)_n\text{OR}_{17}$, or $\text{O}(\text{CH}_2)_u\text{OR}_{17}$;

10

wherein R_{13} is independently H; $-(\text{CH}_2)_u\text{OR}_5$; $-(\text{CH}_2)_t\text{CONR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{NR}_5\text{COR}_5$; $-(\text{CH}_2)_t\text{COR}_7$; $-(\text{CH}_2)_t\text{CO}_2\text{R}_5$; $-(\text{CH}_2)_u\text{NR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{CN}$; straight chained or branched C_1 - C_7 alkyl; C_1 - C_7 alkyl in which the C_2 - C_7 atoms may be optionally substituted with one or more F or Cl; C_3 - C_7 cycloalkyl- C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl; phenyl or C_1 - C_6 phenylalkyl; wherein the phenyl or C_1 - C_6 phenylalkyl may be substituted with one or more of F, Cl, $-\text{CN}$, $-\text{NO}_2$, $-\text{NR}_5\text{R}_6$, $-\text{SO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{COR}_7$, $-(\text{CH}_2)_n\text{OR}_5$, $-(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$, $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{SO}_2\text{NR}_5\text{R}_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

15

20

25

or R_{12} and R_{13} together with the amide linkage to which they are attached are pyrrolidinonyl, piperidonyl, or oxazolidinonyl;

wherein R_{14} is H; straight chained or branched C_1 - C_4 alkyl; F; or $-(\text{CH}_2)_r\text{OR}_5$;

30

wherein R_{15} is H, straight chained or branched C_1 - C_4 alkyl, or F;

with the proviso that when R_{14} is -OH, R_{15} cannot be F;

5

wherein R_3 is independently H; $-(CH_2)_uOR_5$; $-(CH_2)_tCONR_5R_6$; $-(CH_2)_uNR_5COR_5$; $-(CH_2)_tCOR_7$; $-(CH_2)_tCO_2R_5$; $-(CH_2)_uNR_5R_6$; $-(CH_2)_uCN$; straight chained or branched C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl or cycloalkenyl; phenyl, C_1 - C_6 phenylalkyl or C_1 - C_6 heteroarylalkyl; wherein the phenyl, C_1 - C_6 phenylalkyl, or C_1 - C_6 heteroarylalkyl may be substituted with one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -SO₂R₅, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl;

wherein R_4 is independently H; $-(CH_2)_uOR_5$; $-(CH_2)_tCONR_5R_6$; $-(CH_2)_uNR_5COR_5$; $-(CH_2)_tCOR_7$; $-(CH_2)_tCO_2R_5$; $-(CH_2)_uNR_5R_6$; $-(CH_2)_uCN$; straight chained or branched C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl or cycloalkenyl; phenyl or C_1 - C_6 phenylalkyl; wherein the phenyl or C_1 - C_6 phenylalkyl may be substituted with one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -SO₂R₅, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl;

or R_3 and R_4 taken together with the nitrogen atom to which they are attached are 1-azetidiny1, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl, wherein the 1-azetidiny1, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl is substituted with one or more of F, -CN, $-(CH_2)_nNR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl, or phenyl or heteroaryl; wherein if $-(CH_2)_nNR_5R_6$, $-(CH_2)_nOR_5$, or $-(CH_2)_nNR_5COR_5$ are in the 2-position, then n is not 0; wherein the phenyl or heteroaryl may be substituted with one or more of F, Cl, Br, -CN, $-NO_2$, $-NR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl;

or R_3 and R_4 taken together with the nitrogen atom to which they are attached are morpholinyl, thiomorpholinyl, [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or [1,4]diazepanyl, wherein the morpholinyl, thiomorpholinyl, [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or [1,4]diazepanyl is optionally substituted with straight chained or branched C_1 - C_5 alkyl or $(CH_2)_tOR_5$; and wherein the nitrogen atom of the piperazinyl or [1,4]diazepanyl ring may be optionally substituted with $-(CH_2)_uOR_5$; $-COR_5$; $-CO_2R_5$; straight chained or branched C_1 - C_5 alkyl; or phenyl; wherein the phenyl may be substituted with one or more of F, Cl, Br, -CN, $-NO_2$, $-NR_5R_6$, $-(CH_2)_nOR_5$, straight chained or branched C_1 - C_3 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

wherein R_{17} is straight chained or branched C_1-C_4 alkyl, perfluoroalkyl, or polyfluoroalkyl;

5 wherein m is 0 or 1;

wherein each p independently is an integer from 0 to 2 inclusive;

10 wherein each r independently is an integer from 0 to 3 inclusive;

wherein each s independently is an integer from 1 to 6 inclusive;

15 wherein t is an integer from 1 to 4 inclusive;

wherein each u independently is an integer from 2 to 4 inclusive;

20 wherein v is 1 or 2;

with the proviso that when v is 2, m is 0;

25 wherein z is an integer from 2 to 7;

or a pharmaceutically acceptable salt thereof.

30 The invention also provides a pharmaceutical composition comprising a therapeutically effective amount of the compound of the invention and a pharmaceutically acceptable carrier. This invention further provides a pharmaceutical composition made by combining a therapeutically effective amount of the compound of this


invention and a pharmaceutically acceptable carrier. This invention further provides a process for making a pharmaceutical composition comprising combining a therapeutically effective amount of the compound of the invention and a pharmaceutically acceptable carrier.

5

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
221

Detailed Description Of The InventionA. Tricyclic Compounds

This invention provides a compound having the structure:



wherein each R_1 is independently H, F, Cl, Br, -CN, -OH, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, perfluoroalkyl, polyfluoroalkyl, aminoalkyl, or straight chained or branched C₁-C₇ alkyl;

wherein R_5 is independently H; or straight chained or branched C₁-C₇ alkyl;

wherein R_6 is independently H; or straight chained or branched C₁-C₇ alkyl;


wherein B is O, NH or S;

wherein X is CHR₅, O or NR₅;

wherein each n independently is an integer from 0 to 6 inclusive;

wherein R_8 is

5



10


wherein Y is C or N;

wherein R_7 is independently straight chained or branched C_1 - C_7 alkyl;

wherein R_9 is independently H; or straight chained or branched C_1 - C_4 alkyl;

5 wherein R_{10} is independently H; or straight chained or branched C_1 - C_4 alkyl;

wherein R_{11} is



;

10 wherein R_{12} is H, straight chained or branched C_1 - C_7 alkyl, $(\text{CH}_2)_n\text{OR}_{17}$, or $\text{O}(\text{CH}_2)_u\text{OR}_{17}$; provided that when X is O, R_{12} cannot be methyl;

15 wherein R_{13} is independently H; $-(\text{CH}_2)_u\text{OR}_5$; $-(\text{CH}_2)_t\text{CONR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{NR}_5\text{COR}_5$; $-(\text{CH}_2)_t\text{COR}_7$; $-(\text{CH}_2)_t\text{CO}_2\text{R}_5$; $-(\text{CH}_2)_u\text{NR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{CN}$; straight chained or branched C_1 - C_7 alkyl; C_1 - C_7 alkyl in which the C_2 - C_7 atoms may be optionally substituted with one or more F or Cl; C_3 - C_7 cycloalkyl- C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl; phenyl or C_1 - C_6 phenylalkyl; wherein the phenyl or C_1 - C_6 phenylalkyl may be substituted with one or more of F, Cl, $-\text{CN}$, $-\text{NO}_2$, $-\text{NR}_5\text{R}_6$, $-\text{SO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{COR}_7$, $-(\text{CH}_2)_n\text{OR}_5$, $-(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$, $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{SO}_2\text{NR}_5\text{R}_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

or R_{12} and R_{13} together with the amide linkage to which they are attached are pyrrolidinonyl, piperidonyl, or oxazolidinonyl; provided that when X is O, R_{12} and R_{13} cannot be oxazolidinonyl;

5

wherein R_{14} is H; straight chained or branched C_1 - C_4 alkyl; F; or $-(CH_2)_rOR_5$;

wherein R_{15} is H, straight chained or branched C_1 - C_4 alkyl, or F;

10

with the proviso that when R_{14} is -OH, R_{15} cannot be F;

wherein R_{16} is NR_3R_4 , perfluoroalkyl, unsubstituted straight chained or branched C_1 - C_7 alkyl, substituted straight chained or branched C_2 - C_7 alkyl, wherein the C_2 - C_7 alkyl may be substituted with one or more of F, Cl, -CN, $-NR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nOCF_3$,

15

perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl; C_3 - C_7 cycloalkyl or cycloalkenyl; phenyl, heteroaryl, or C_1 - C_7 phenylalkyl, wherein the phenyl, heteroaryl, or C_1 - C_7 phenylalkyl may be substituted with one or more of F, Cl, Br, -CN, $-NO_2$, $-NR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, ethylenedioxy, methylenedioxy, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl; quinolinyl, 1-naphthyl, 2-naphthyl, or 2,1,3-benzothiadiazolyl; wherein the quinolinyl, 1-naphthyl, 2-naphthyl or 2,1,3-benzothiadiazolyl may be substituted with one or more of

25

30

32

F, Cl, Br, -CN, -NO₂, -NR₅R₆, -
 (CH₂)_nNR₅COR₅, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -
 (CH₂)_nCONR₅R₆, -(CH₂)_nCO₂R₅, -(CH₂)_nSO₂NR₅R₆, ethylenedioxy,
 methylenedioxy, straight chained or branched C₁-C₇ alkyl,
 5 perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

with the proviso that when X is O and R₈ is
 NR₉(CH₂)_uO(CH₂)_uNR₁₀R₁₁, R₁₆ cannot be methyl;

10

wherein R₃ is independently H; -(CH₂)_uOR₅; -
 (CH₂)_tCONR₅R₆; -(CH₂)_uNR₅COR₅; -(CH₂)_tCOR₇; -
 (CH₂)_tCO₂R₅; -(CH₂)_uNR₅R₆; -(CH₂)_uCN; straight chained or
 15 branched C₁-C₇ alkyl; straight chained or branched C₂-C₇
 alkenyl or alkynyl; or C₃-C₇ cycloalkyl or cycloalkenyl;
 phenyl, C₁-C₆ phenylalkyl, or C₁-C₆ heteroarylalkyl; wherein
 the phenyl, C₁-C₆ phenylalkyl or C₁-C₆ heteroarylalkyl may
 be substituted with one or more of F, Cl, Br, -CN, -NO₂,
 20 -NR₅R₆, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -
 (CH₂)_nNR₅COR₅, -(CH₂)_nCO₂R₅, -(CH₂)_nSO₂NR₅R₆, straight chained
 or branched C₁-C₇ alkyl, perfluoroalkyl, polyfluoroalkyl,
 or aminoalkyl, straight chained or branched C₂-C₇ alkenyl
 or alkynyl, or C₃-C₇ cycloalkyl or cycloalkenyl;

25

wherein R₄ is independently H; -(CH₂)_uOR₅; -
 (CH₂)_tCONR₅R₆; -(CH₂)_uNR₅COR₅; -(CH₂)_tCOR₇; -
 (CH₂)_tCO₂R₅; -(CH₂)_uNR₅R₆; -(CH₂)_uCN; straight chained or
 30 branched C₁-C₇ alkyl; straight chained or branched C₂-C₇
 alkenyl or alkynyl; or C₃-C₇ cycloalkyl or cycloalkenyl;
 phenyl or C₁-C₆ phenylalkyl; wherein the phenyl or C₁-C₆
 phenylalkyl may be substituted with one or more of F, Cl,
 Br, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -

$(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$, $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{SO}_2\text{NR}_5\text{R}_6$,
 straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl,
 polyfluoroalkyl, or aminoalkyl, straight chained or
 branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or
 cycloalkenyl;

or R_3 and R_4 taken together with the nitrogen atom to which
 they are attached are 1-azetidiny, 1-pyrrolidinyl, 1-
 piperidinyl, or 1H-azepanyl, wherein the 1-azetidiny, 1-
 pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl is substituted
 with one or more of F, -CN, $-(\text{CH}_2)_n\text{NR}_5\text{R}_6$, $-\text{SO}_2\text{R}_5$,
 $-(\text{CH}_2)_n\text{COR}_7$, $-(\text{CH}_2)_n\text{OR}_5$, $-(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$,
 $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$, straight chained or branched C_1 - C_7 alkyl,
 perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight
 chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7
 cycloalkyl or cycloalkenyl, or phenyl or heteroaryl;
 wherein if $-(\text{CH}_2)_n\text{NR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{OR}_5$, or $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$ are
 in the 2-position, then n is not 0; wherein the phenyl or
 heteroaryl may be substituted with one or more of F, Cl,
 Br, -CN, $-\text{NO}_2$, $-\text{NR}_5\text{R}_6$, $-\text{SO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{COR}_7$,
 $-(\text{CH}_2)_n\text{OR}_5$, $-(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$, $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$,
 $-(\text{CH}_2)_n\text{SO}_2\text{NR}_5\text{R}_6$, straight chained or branched C_1 - C_7 alkyl,
 perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight
 chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7
 cycloalkyl or cycloalkenyl;

or R_3 and R_4 taken together with the nitrogen atom to which
 they are attached are morpholinyl, thiomorpholinyl,
 [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or
 [1,4]diazepanyl, wherein the morpholinyl, thiomorpholinyl,
 [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or
 [1,4]diazepanyl is optionally substituted with straight
 chained or branched C_1 - C_5 alkyl or $(\text{CH}_2)_t\text{OR}_5$; and wherein
 the nitrogen atom of the piperazinyl or [1,4]diazepanyl

34

ring may be optionally substituted with -
 $(\text{CH}_2)_u\text{OR}_5$; $-\text{COR}_5$; straight chained or branched $\text{C}_1\text{-C}_5$ alkyl;
 or phenyl; wherein the phenyl may be substituted with one
 or more of F, Cl, Br, $-\text{CN}$, $-\text{NO}_2$, $-\text{NR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{OR}_5$, straight
 5 chained or branched $\text{C}_1\text{-C}_3$ alkyl, perfluoroalkyl,
 polyfluoroalkyl, or aminoalkyl;

wherein R_{17} is H, straight chained or branched $\text{C}_1\text{-C}_4$ alkyl,
 perfluoroalkyl, or polyfluoroalkyl;

10

wherein R_{19} is $(\text{CH}_2)_n\text{OR}_5$, NR_5R_6 , phenyl, or heteroaryl,
 wherein the phenyl or heteroaryl may be substituted with
 one or more of F, Cl, Br, $-\text{CN}$, $-\text{NO}_2$, $-\text{NR}_5\text{R}_6$, -
 $(\text{CH}_2)_n\text{NR}_5\text{COR}_5$, $-\text{SO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{COR}_7$, $-(\text{CH}_2)_n\text{OR}_5$,
 15 $-(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{SO}_2\text{NR}_5\text{R}_6$, ethylenedioxy,
 methylenedioxy, straight chained or branched $\text{C}_1\text{-C}_7$ alkyl,
 perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight
 chained or branched $\text{C}_2\text{-C}_7$ alkenyl or alkynyl, or $\text{C}_3\text{-C}_7$
 cycloalkyl or cycloalkenyl;

20

wherein m is 0 or 1;

wherein each p independently is an integer from 0 to 2
 inclusive;

25

wherein each r independently is an integer from 0 to 3
 inclusive;

30

wherein each s independently is an integer from 1 to 6
 inclusive;

wherein t is an integer from 1 to 4 inclusive;

35

wherein each u independently is an integer from 2 to 4 inclusive;

wherein v is 1 or 2;

5

with the proviso that when v is 2, m is 0;

wherein z is an integer from 2 to 7;

10


or a pharmaceutically acceptable salt thereof.

In one embodiment, the compound comprises the (+) enantiomer. In another embodiment, the compound comprises the (-) enantiomer.

15

36


In one embodiment, the compound has the structure:



wherein X is CR_5R_6 , O or NR_5 .


5

In another embodiment, the compound has the structure:




37




In still another embodiment, the compound has the structure:




5 In a further embodiment, the compound has the structure:



In still further embodiments, the compound has the structure selected from the group consisting of:




, and



39





In another embodiment, the compound has the structure:






wherein s is 2, 3 or 4.


5

In still other embodiments, the compound has the structure selected from the group consisting of:









, and










In a further embodiment, the compound has the structure:







5 In still further embodiments, the compound has the structure selected from the group consisting of:









In still other embodiments, the compound has the structure:




or



In still further embodiments, the compound has the structure selected from the group consisting of:



and



5

In the present invention as relates to tricyclic compounds, the term "heteroaryl" is used to include five and six membered unsaturated rings that may contain one or more heteroatoms such as oxygen, sulfur, and nitrogen. Examples of heteroaryl groups include, but are not limited to, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl. In addition the term "heteroaryl" is used to include fused bicyclic ring systems that may contain one or more heteroatoms such as oxygen, sulfur and nitrogen. Examples of such heteroaryl groups include, but are not limited to, indolizinyl, indolyl, isoindolyl, benzo[b]furanyl, benzo[b]thiophenyl, indazolyl, benzimidazolyl, benzthiazolyl, purinyl, imidazo[2,1-b]thiazolyl, quinolinyl, isoquinolinyl, quinolizinyl, and 2,1,3-

10


15

20

benzothiazolyl. Furthermore, any of the heteroaryl groups recited above may be substituted with thienyl, isoxazolyl, or pyridyl.

B. Bicyclic Compounds

This invention provides a compound having the structure:



5 wherein Y is O, S or NH;

wherein each R_{14} independently is H, F, Cl, Br, -CN, -OH, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, -SO₂C₆H₅, -SO₂NR₅R₆, -C₆H₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, ethylenedioxy, methylenedioxy, perfluoroalkyl, polyfluoroalkyl, aminoalkyl, or straight chained or branched C₁-C₇ alkyl; or phenyl, heteroaryl, or C₁-C₇ phenylalkyl, wherein the phenyl, heteroaryl, or C₁-C₇ phenylalkyl may be substituted with one or more of F, Cl, Br, -CF₃, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, or straight chained or branched C₁-C₄ alkyl; provided that if one R_{14} is phenyl, heteroaryl or C₁-C₇ phenylalkyl, the other R_{14} is H;

wherein each R_1 independently is H, F, Cl, Br, -CN, -OH, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, -SO₂C₆H₅, -SO₂NR₅R₆, -C₆H₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, ethylenedioxy, methylenedioxy, perfluoroalkyl, polyfluoroalkyl, aminoalkyl, or straight chained or branched C₁-C₇ alkyl; or phenyl, heteroaryl, or C₁-C₇ phenylalkyl, wherein the phenyl, heteroaryl, or C₁-C₇ phenylalkyl may be substituted with one or more of F, Cl, Br, -CF₃, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, or straight chained or branched C₁-C₄ alkyl;

wherein R_2 is H, straight chained or branched C_1 - C_4 alkyl,
 $-(CH_2)_tOR_5$, phenyl optionally substituted with one or more
 of F, Cl, Br, $-CF_3$, $-CN$, $-NO_2$, $-NR_5R_6$, $-SO_2R_5$,
 5 $-(CH_2)_nOR_5$, or straight chained or branched C_1 - C_4 alkyl;


wherein R_5 is independently H; or straight chained or
 branched C_1 - C_7 alkyl;

10 wherein R_6 is independently H; or straight chained or
 branched C_1 - C_7 alkyl;

wherein each n independently is an integer from 0 to 6
 inclusive;

15

wherein R_8 is



20

provided that R_1 or R_{14} cannot be $-OH$, when R_8 is (ii);


wherein R_9 is independently H; or straight chained or
 branched C_1 - C_4 alkyl;

25

49

wherein R_{10} is independently H; or straight chained or branched C_1 - C_4 alkyl;

wherein R_{11} is



5

wherein R_{12} is H, straight chained or branched C_1 - C_7 alkyl; or $(\text{CH}_2)_n\text{OR}_{17}$;

10

wherein R_{13} is independently $-(\text{CH}_2)_u\text{OR}_5$; $-(\text{CH}_2)_t\text{CONR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{NR}_5\text{COR}_5$; $-(\text{CH}_2)_t\text{COR}_7$; $-(\text{CH}_2)_t\text{CO}_2\text{R}_5$; $-(\text{CH}_2)_u\text{NR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{CN}$; straight chained or branched C_1 - C_7 alkyl; C_1 - C_7 alkyl in which the C_2 - C_7 atoms may be optionally substituted with one or more F or Cl; C_3 - C_7 cycloalkyl- C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl; phenyl or C_1 - C_6 phenylalkyl; wherein the phenyl or C_1 - C_6 phenylalkyl may be substituted with one or more of F, Cl, $-\text{CN}$, $-\text{NO}_2$, $-\text{NR}_5\text{R}_6$, $-\text{SO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{COR}_7$, $-(\text{CH}_2)_n\text{OR}_5$, $-(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$, $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{SO}_2\text{NR}_5\text{R}_6$, or straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

15

20

or R_{12} and R_{13} together with the amide linkage to which they are attached are pyrrolidinonyl, piperidonyl, or oxazolidinonyl;

25

wherein R_7 is independently straight chained or branched C_1 - C_7 alkyl;

30

wherein R_{16} is NR_3R_4 , perfluoroalkyl, unsubstituted straight chained or branched C_1 - C_7 alkyl, substituted straight chained or branched C_2 - C_7 alkyl, wherein the C_2 - C_7 alkyl may be substituted with one or more of F, Cl, -CN, - NR_5R_6 , - SO_2R_5 , - $(CH_2)_nCOR_7$, - $(CH_2)_nOR_5$, - $(CH_2)_nCONR_5R_6$, - $(CH_2)_nNR_5COR_5$, - $(CH_2)_nCO_2R_5$, - $(CH_2)_nOCF_3$, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl; C_3 - C_7 cycloalkyl or cycloalkenyl; or phenyl, heteroaryl, or C_1 - C_7 phenylalkyl, wherein the phenyl, heteroaryl, or C_1 - C_7 phenylalkyl may be substituted with one or more of F, Cl, Br, I, -CN, - NO_2 , - NR_5R_6 , - $(CH_2)_nNR_5COR_5$, - SO_2R_5 , - $(CH_2)_nCOR_7$, - $(CH_2)_nOR_5$, - $(CH_2)_nCONR_5R_6$, - $(CH_2)_nCO_2R_5$, - $(CH_2)_nSO_2NR_5R_6$, ethylenedioxy, methylenedioxy, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl; quinolinyl, 1-naphthyl, 2-naphthyl, or 2,1,3-benzothiadiazolyl; wherein the quinolinyl, 1-naphthyl, 2-naphthyl or 2,1,3-benzothiadiazolyl may be substituted with one or more of F, Cl, Br, I, -CN, - NO_2 , - NR_5R_6 , - $(CH_2)_nNR_5COR_5$, - SO_2R_5 , - $(CH_2)_nCOR_7$, - $(CH_2)_nOR_5$, - $(CH_2)_nCONR_5R_6$, - $(CH_2)_nCO_2R_5$, - $(CH_2)_nSO_2NR_5R_6$, ethylenedioxy, methylenedioxy, or straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

wherein R_3 is independently H; - $(CH_2)_uOR_5$; - $(CH_2)_tCONR_5R_6$; - $(CH_2)_uNR_5COR_5$; - $(CH_2)_tCOR_7$; - $(CH_2)_tCO_2R_5$; - $(CH_2)_uNR_5R_6$; - $(CH_2)_uCN$; straight chained or branched C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl or cycloalkenyl; or phenyl, C_1 - C_6 phenylalkyl, or C_1 - C_6 heteroarylalkyl; wherein the phenyl, C_1 - C_6 phenylalkyl, or C_1 - C_6

heteroarylalkyl may be substituted with one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, -(CH₂)_nCO₂R₅, -(CH₂)_nSO₂NR₅R₆, straight chained or branched C₁-C₇ alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C₂-C₇ alkenyl or alkynyl, or C₃-C₇ cycloalkyl or cycloalkenyl;

wherein R₄ is independently H; -(CH₂)_uOR₅; -(CH₂)_tCONR₅R₆; -(CH₂)_uNR₅COR₅; -(CH₂)_tCOR₇; -(CH₂)_tCO₂R₅; -(CH₂)_uNR₅R₆; -(CH₂)_uCN; straight chained or branched C₁-C₇ alkyl; straight chained or branched C₂-C₇ alkenyl or alkynyl; or C₃-C₇ cycloalkyl or cycloalkenyl; or phenyl or C₁-C₆ phenylalkyl; wherein the phenyl or C₁-C₆ phenylalkyl may be substituted with one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, -(CH₂)_nCO₂R₅, -(CH₂)_nSO₂NR₅R₆, straight chained or branched C₁-C₇ alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C₂-C₇ alkenyl or alkynyl, or C₃-C₇ cycloalkyl or cycloalkenyl;

or R₃ and R₄ taken together with the nitrogen atom to which they are attached are 1-azetidiny, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl, wherein the 1-azetidiny, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl is substituted with one or more of F, -CN, -(CH₂)_nNR₅R₆, -SO₂R₅, -(CH₂)_nCOR₇, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, -(CH₂)_nCO₂R₅, straight chained or branched C₁-C₇ alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C₂-C₇ alkenyl or alkynyl, or C₃-C₇ cycloalkyl or cycloalkenyl, or phenyl or heteroaryl; wherein if -(CH₂)_nNR₅R₆, -(CH₂)_nOR₅, or -(CH₂)_nNR₅COR₅ are in the 2-position, then n is not 0; wherein the phenyl or

52

heteroaryl may be substituted with one or more
 of F, Cl, Br, I, -CN, -NO₂, -NR₅R₆, -SO₂R₅, -
 (CH₂)_nCOR₇, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, -
 (CH₂)_nCO₂R₅, -(CH₂)_nSO₂NR₅R₆, straight chained or branched C₁-
 5 C₇ alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl,
 straight chained or branched C₂-C₇ alkenyl or alkynyl, or
 C₃-C₇ cycloalkyl or cycloalkenyl;

or R₃ and R₄ taken together with the nitrogen atom to which
 10 they are attached are morpholinyl, thiomorpholinyl,
 [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or
 [1,4]diazepanyl, wherein the morpholinyl, thiomorpholinyl,
 [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or
 15 [1,4]diazepanyl is optionally substituted with straight
 chained or branched C₁-C₅ alkyl or (CH₂)_tOR₅; and wherein
 the nitrogen atom of the piperazinyl or [1,4]diazepanyl
 ring may be optionally substituted with -(CH₂)_uOR₅; -COR₅;
 straight chained or branched C₁-C₅ alkyl; or phenyl;
 wherein the phenyl may be substituted with one or more of
 20 F, Cl, Br, -CN, -NO₂, -NR₅R₆, -(CH₂)_nOR₅, straight chained or
 branched C₁-C₃ alkyl, perfluoroalkyl, polyfluoroalkyl, or
 aminoalkyl;

wherein R₁₇ is straight chained or branched C₁-C₄ alkyl,
 25 perfluoroalkyl, or polyfluoroalkyl;

wherein each p independently is an integer from 0 to 2
 inclusive;

30 wherein each r independently is an integer from 0 to 3
 inclusive;

wherein t is an integer from 1 to 4 inclusive;

53

wherein each u independently is an integer from 2 to 4 inclusive;

or a pharmaceutically acceptable salt thereof.


5

In one embodiment, the compound comprises the (+) enantiomer. In another embodiment, the compound comprises the (-) enantiomer.

10
20
30
40
50
60
70
80
90
100
110
120
130
140
150
160
170
180
190
200
210
220
230
240
250
260
270
280
290
300
310
320
330
340
350
360
370
380
390
400
410
420
430
440
450
460
470
480
490
500
510
520
530
540
550
560
570
580
590
600
610
620
630
640
650
660
670
680
690
700
710
720
730
740
750
760
770
780
790
800
810
820
830
840
850
860
870
880
890
900
910
920
930
940
950
960
970
980
990
1000


54

In one embodiment, the compound has the structure:




5

In further embodiments, the compound has the structure selected from the group consisting of:




and




10

55


In another embodiment, the compound has the structure:



5 In further embodiment, the compound has the structure selected from the group consisting of:




and



In the present invention as relates to bicyclic compounds, the term "heteroaryl" is used to include five and six membered unsaturated rings that may contain one sulfur or nitrogen atom or one or more oxygen, sulfur, or nitrogen atoms. Examples of heteroaryl groups include, but are not limited to, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl. In addition the term "heteroaryl" is used to include fused bicyclic ring systems that may contain one or more heteroatoms such as oxygen, sulfur and nitrogen. Examples of such heteroaryl groups include, but are not limited to, indoliziny, indolyl, isoindolyl, benzo[b]furanyl, benzo[b]thiophenyl, indazolyl, benzimidazolyl, benzthiazolyl, purinyl, imidazo[2,1-b]thiazolyl, quinolinyl, isoquinolinyl, quinoliziny, and 2,1,3-benzothiazolyl.

C. Tricyclic Compounds

The invention provides a compound having the structure:



5 wherein each R_1 is independently H, F, Cl, Br, -CN, -OH, -NO₂, -NR₅R₆, -SO₂R₅, -(CH₂)_nOR₅, -(CH₂)_nCONR₅R₆, -(CH₂)_nNR₅COR₅, perfluoroalkyl, polyfluoroalkyl, aminoalkyl, or straight chained or branched C₁-C₇ alkyl;

10 wherein R₅ is independently H; or straight chained or branched C₁-C₇ alkyl;


wherein R₆ is independently H; or straight chained or branched C₁-C₇ alkyl;

15 wherein B is O, NH or S;

wherein X is S, SO or SO₂;

20 wherein each n independently is an integer from 0 to 6 inclusive;

wherein R_8 is



5

wherein Y is C or N;

wherein R_7 is independently straight chained or branched
 10 C_1 - C_7 alkyl;


wherein R_9 is independently H; or straight chained or
 branched C_1 - C_4 alkyl;

59

wherein R_{10} is independently H; or straight chained or branched C_1 - C_4 alkyl;

wherein R_{11} is

5



wherein R_{12} is H, straight chained or branched C_1 - C_7 alkyl, $(\text{CH}_2)_n\text{OR}_{17}$, or $\text{O}(\text{CH}_2)_u\text{OR}_{17}$;

10

wherein R_{13} is independently H; $-(\text{CH}_2)_u\text{OR}_5$; $-(\text{CH}_2)_t\text{CONR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{NR}_5\text{COR}_5$; $-(\text{CH}_2)_t\text{COR}_7$; $-(\text{CH}_2)_t\text{CO}_2\text{R}_5$; $-(\text{CH}_2)_u\text{NR}_5\text{R}_6$; $-(\text{CH}_2)_u\text{CN}$; straight chained or branched C_1 - C_7 alkyl; C_1 - C_7 alkyl in which the C_2 - C_7 atoms may be optionally substituted with one or more F or Cl; C_3 - C_7 cycloalkyl- C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl; phenyl or C_1 - C_6 phenylalkyl; wherein the phenyl or C_1 - C_6 phenylalkyl may be substituted with one or more of F, Cl, $-\text{CN}$, $-\text{NO}_2$, $-\text{NR}_5\text{R}_6$, $-\text{SO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{COR}_7$, $-(\text{CH}_2)_n\text{OR}_5$, $-(\text{CH}_2)_n\text{CONR}_5\text{R}_6$, $-(\text{CH}_2)_n\text{NR}_5\text{COR}_5$, $-(\text{CH}_2)_n\text{CO}_2\text{R}_5$, $-(\text{CH}_2)_n\text{SO}_2\text{NR}_5\text{R}_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

20

or R_{12} and R_{13} together with the amide linkage to which they are attached are pyrrolidinonyl, piperidonyl, or oxazolidinonyl;

25

wherein R_{14} is H; straight chained or branched C_1 - C_4 alkyl; F; or $-(\text{CH}_2)_r\text{OR}_5$;

30

wherein R_{15} is H, straight chained or branched C_1 - C_4 alkyl, or F;

with the proviso that when R_{14} is -OH, R_{15} cannot be F;

5

wherein R_3 is independently H; $-(CH_2)_uOR_5$; $-(CH_2)_tCONR_5R_6$; $-(CH_2)_uNR_5COR_5$; $-(CH_2)_tCOR_7$; $-(CH_2)_tCO_2R_5$; $-(CH_2)_uNR_5R_6$; $-(CH_2)_uCN$; straight chained or branched C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl or cycloalkenyl; phenyl, C_1 - C_6 phenylalkyl or C_1 - C_6 heteroarylalkyl; wherein the phenyl, C_1 - C_6 phenylalkyl, or C_1 - C_6 heteroarylalkyl may be substituted with one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -SO₂R₅, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl;

20

wherein R_4 is independently H; $-(CH_2)_uOR_5$; $-(CH_2)_tCONR_5R_6$; $-(CH_2)_uNR_5COR_5$; $-(CH_2)_tCOR_7$; $-(CH_2)_tCO_2R_5$; $-(CH_2)_uNR_5R_6$; $-(CH_2)_uCN$; straight chained or branched C_1 - C_7 alkyl; straight chained or branched C_2 - C_7 alkenyl or alkynyl; or C_3 - C_7 cycloalkyl or cycloalkenyl; phenyl or C_1 - C_6 phenylalkyl; wherein the phenyl or C_1 - C_6 phenylalkyl may be substituted with one or more of F, Cl, Br, -CN, -NO₂, -NR₅R₆, -SO₂R₅, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl;

30

61

or R_3 and R_4 taken together with the nitrogen atom to which they are attached are 1-azetidiny1, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl, wherein the 1-azetidiny1, 1-pyrrolidinyl, 1-piperidinyl, or 1H-azepanyl is substituted with one or more of F, -CN, $-(CH_2)_nNR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl, or phenyl or heteroaryl; wherein if $-(CH_2)_nNR_5R_6$, $-(CH_2)_nOR_5$, or $-(CH_2)_nNR_5COR_5$ are in the 2-position, then n is not 0; wherein the phenyl or heteroaryl may be substituted with one or more of F, Cl, Br, -CN, $-NO_2$, $-NR_5R_6$, $-SO_2R_5$, $-(CH_2)_nCOR_7$, $-(CH_2)_nOR_5$, $-(CH_2)_nCONR_5R_6$, $-(CH_2)_nNR_5COR_5$, $-(CH_2)_nCO_2R_5$, $-(CH_2)_nSO_2NR_5R_6$, straight chained or branched C_1 - C_7 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl, straight chained or branched C_2 - C_7 alkenyl or alkynyl, or C_3 - C_7 cycloalkyl or cycloalkenyl;

20

or R_3 and R_4 taken together with the nitrogen atom to which they are attached are morpholinyl, thiomorpholinyl, [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or [1,4]diazepanyl, wherein the morpholinyl, thiomorpholinyl, [1,4]oxazepanyl, [1,4]thiazepanyl, piperazinyl, or [1,4]diazepanyl is optionally substituted with straight chained or branched C_1 - C_5 alkyl or $(CH_2)_tOR_5$; and wherein the nitrogen atom of the piperazinyl or [1,4]diazepanyl ring may be optionally substituted with $-(CH_2)_uOR_5$, $-COR_5$, $-CO_2R_5$; straight chained or branched C_1 - C_5 alkyl; or phenyl; wherein the phenyl may be substituted with one or more of F, Cl, Br, -CN, $-NO_2$, $-NR_5R_6$, $-(CH_2)_nOR_5$, straight chained or branched C_1 - C_3 alkyl, perfluoroalkyl, polyfluoroalkyl, or aminoalkyl;

30

wherein R_1 is straight chained or branched C_1 - C_4 alkyl, perfluoroalkyl, or polyfluoroalkyl;

5 wherein m is 0 or 1;

wherein each p independently is an integer from 0 to 2 inclusive;

10 wherein each r independently is an integer from 0 to 3 inclusive;

wherein each s independently is an integer from 1 to 6 inclusive;

15 wherein t is an integer from 1 to 4 inclusive;

wherein each u independently is an integer from 2 to 4 inclusive;

20 wherein v is 1 or 2;

with the proviso that when v is 2, m is 0;

25 wherein z is an integer from 2 to 7;


or a pharmaceutically acceptable salt thereof.

30 In one embodiment, the compound comprises the (+) enantiomer. In another embodiment, the compound comprises the (-) enantiomer.

63


In one embodiment, the

compound has the structure:





In another embodiment, the compound has the structure:

5




64

In still other embodiments, the compound has the structure:




or





5

In a further embodiment, the compound has the structure:




10

In still further embodiments, the compound has the structure:




5

In another embodiment, the compound has the structure:




In still another embodiment, ⁶⁶ the compound has the structure:




5

In a further embodiment, the compound has the structure:



67

In a still further embodiment, the compound has the structure:



5

10

15

20

25

In the present invention as relates to tricyclic compounds, the term "heteroaryl" is used to include five and six membered unsaturated rings that may contain one or more heteroatoms such as oxygen, sulfur, and nitrogen. Examples of heteroaryl groups include, but are not limited to, furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, and triazinyl. In addition the term "heteroaryl" is used to include fused bicyclic ring systems that may contain one or more heteroatoms such as oxygen, sulfur and nitrogen. Examples of such heteroaryl groups include, but are not limited to, indolizinyl, indolyl, isoindolyl, benzo[b]furanyl, benzo[b]thiophenyl, indazolyl, benzimidazolyl, benzthiazolyl, purinyl, imidazo[2,1-b]thiazolyl, quinolinyl, isoquinolinyl, quinolizinyl, and 2,1,3-benzothiazolyl. Furthermore, any of the heteroaryl groups recited above may be substituted with thienyl, isoxazolyl, or pyridyl.

Included in this invention are pharmaceutically acceptable salts and complexes of all of the compounds described herein. The salts include but are not limited to the acids and bases listed herein. The salts include, but are not limited to the following inorganic acids: hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid and boric acid. The salts include, but are not limited to the following organic acids: acetic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, maleic acid, citric acid, methanesulfonic acid, benzoic acid, glycolic acid, lactic acid and mandelic acid. The salts include, but are not limited to the inorganic base, ammonia. The salts include, but are not limited to the following organic bases: methylamine, ethylamine, propylamine, dimethylamine, diethylamine, trimethylamine, triethylamine, ethylenediamine, hydroxyethylamine, morpholine, piperazine and guanidine. This invention further provides for the hydrates and polymorphs of all of the compounds described herein.

The present invention includes within its scope prodrugs of the compounds of the invention. In general, such prodrugs will be functional derivatives of the compounds of the invention which are readily convertible in vivo into the required compound. Thus, in the present invention, the term "administering" shall encompass the treatment of the various conditions described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound in vivo after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in Design of Prodrugs, ed. H. Bundgaard, Elsevier, 1985.

5 The present invention further includes metabolites of the compounds of the present invention. Metabolites include active species produced upon introduction of compounds of this invention into the biological milieu.

10 This invention further provides a pharmaceutical composition comprising a therapeutically effective amount of a compound of the invention and a pharmaceutically acceptable carrier. In one embodiment, the amount of the compound is an amount from about 0.01 mg to about 800 mg. In another embodiment, the amount of the compound is an amount from about 0.01 mg to about 500 mg. In another embodiment, the amount of the compound is an amount from about 0.01 mg to about 250 mg. In another embodiment, the amount of the compound is an amount from about 0.1 mg to about 60 mg. In another embodiment, the amount of the compound is an amount from about 1 mg to about 20 mg. In a further embodiment, the carrier is a liquid and the composition is a solution. In another embodiment, the carrier is a solid and the composition is a tablet. In a further embodiment, the carrier is a gel and the composition is a suppository.

25 This invention provides a pharmaceutical composition made by combining a therapeutically effective amount of a compound of this invention and a pharmaceutically acceptable carrier.

30 This invention provides a process for making a pharmaceutical composition comprising combining a therapeutically effective amount of a compound of this invention and a pharmaceutically acceptable carrier.

This invention provides a use of a compound of this invention for the preparation of a pharmaceutical composition for treating an abnormality, wherein the abnormality is alleviated by decreasing the activity of a human Y5 receptor. In different embodiments, the abnormality is an eating disorder, obesity, bulimia nervosa, a sexual disorder, a reproductive disorder, depression, an epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure, or a sleep disturbance.

In the subject invention a "therapeutically effective amount" is any amount of a compound which, when administered to a subject suffering from a disease against which the compounds are effective, causes reduction, remission, or regression of the disease.

In the practice of this invention the "pharmaceutically acceptable carrier" is any physiological carrier known to those of ordinary skill in the art useful in formulating pharmaceutical compositions.

In one preferred embodiment the pharmaceutical carrier may be a liquid and the pharmaceutical composition would be in the form of a solution. In another equally preferred embodiment, the pharmaceutically acceptable carrier is a solid and the composition is in the form of a powder or tablet. In a further embodiment, the pharmaceutical carrier is a gel and the composition is in the form of a suppository or cream. In a further embodiment the compound may be formulated as a part of a pharmaceutically acceptable transdermal patch.

A solid carrier can include one or more substances which may also act as flavoring agents, lubricants,

solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents; it can also be an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

Liquid carriers are used in preparing solutions, suspensions, emulsions, syrups, elixirs and pressurized compositions. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fats. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (partially containing additives as above, e.g. cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g. glycols) and their derivatives, and oils (e.g. fractionated coconut oil and arachis oil). For parenteral administration, the carrier can also be an oily ester such as ethyl oleate and

isopropyl myristate. Sterile liquid carriers are useful in sterile liquid form compositions for parenteral administration. The liquid carrier for pressurized compositions can be halogenated hydrocarbon or other pharmaceutically acceptable propellant.

Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by for example, intramuscular, intrathecal, epidural, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. The compounds may be prepared as a sterile solid composition which may be dissolved or suspended at the time of administration using sterile water, saline, or other appropriate sterile injectable medium. Carriers are intended to include necessary and inert binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings.

The compound can be administered orally in the form of a sterile solution or suspension containing other solutes or suspending agents (for example, enough saline or glucose to make the solution isotonic), bile salts, acacia, gelatin, sorbitan monoleate, polysorbate 80 (oleate esters of sorbitol and its anhydrides copolymerized with ethylene oxide) and the like.

The compound can also be administered orally either in liquid or solid composition form. Compositions suitable for oral administration include solid forms, such as pills, capsules, granules, tablets, and powders, and liquid forms, such as solutions, syrups, elixirs, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions, and suspensions.

Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound in use, the strength of the preparation, the mode of administration, and the advancement of the disease condition. Additional factors depending on the particular subject being treated will result in a need to adjust dosages, including subject age, weight, gender, diet, and time of administration.

10 One skilled in the art will readily appreciate that appropriate biological assays will be used to determine the therapeutic potential of the claimed compounds for treating the above noted disorders.

15 This invention further provides compositions which need not be pharmaceutical as that term is understood in the art. Such compositions comprise a compound in accordance with the subject invention in an amount effective to agonize and/or antagonize a Y5 receptor and a suitable carrier.

20 Still further, the invention provides a method of agonizing and/or antagonizing a Y5 receptor which comprises contacting the receptor, e.g. in vitro or in vivo, with an amount of a compound of this invention effective to agonize and/or antagonize the receptor.

25 This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow

30 thereafter.

Experimental Details and ResultsI. Synthetic Methods for ExamplesA. Tricyclic Compounds

5

General Procedures Relating to Examples:

10

For the formation of 2-aminothiazoles from 2-haloketones and thioureas, see, for example, Kearney, P.C., et al., 1998; Di Fabio, R. and Pentassuglia, G., 1998; De Kimpe, N., et al., 1996; Plazzi, P.V., et al., 1995; and Novikova, A. P., 1991.

15

For the formation of thiazoles from 2-haloketones and thioamides, see, for example, Critcher, D. J. and Pattenden, G., 1996; and Friedman, B. S., et al., 1937.

20

For the formation of 2-aminoimidazoles from 2-haloketones and guanidines, see, for example, Little, T. L. and Webber, 1994; and Chabaka, L.M., et al., 1994.

25

For the formation of imidazoles from 2-haloketones and amidines, see, for example, Demchenko, A. M., et al., 1997; and Nagao, Y., et al., 1996.

30

For the synthesis of 2-aminooxazoles from 2-haloketones and ureas, see, for example, Pathak, V.N., et al., 1993; Crangk, G. and Foulis, M.J., 1971; and Marchetti, E., et al., 1968.

For the formation of oxazoles from 2-haloketones and amides, see, for example, Hammar, W.J. and Rustad, M.A., 1981; and Zhao, Z., et al., 1991.

All reactions were performed under an inert atmosphere (Argon) and the reagents, neat or in appropriate solvents, were transferred to the reaction vessel via syringe and cannula techniques. Anhydrous solvents were purchased from Aldrich Chemical Company and used as received. Examples 1-90 described in this patent application were named using ACD/Name program (version 2.51, Advanced Chemistry Development Inc., Toronto, Ontario, M5H2L3, Canada).

¹H and ¹³C spectra were recorded at 300 and 75 MHz (QE Plus) with CDCl₃ as solvent (unless otherwise noted) and tetramethylsilane as internal standard. s = singlet; d = doublet; t = triplet; q = quartet; p = pentet; sextet; septet; b = broad; m = multiplet. Elemental analyses were performed by Robertson Microlit Laboratories, Inc. Low-resolution electrospray MS spectra were measured (ESMS, MS) and MH⁺ is reported. Thin-layer chromatography (TLC) was carried out on glass plates precoated with silica gel 60 F₂₅₄ (0.25 mm, EM Separations Tech.). Preparative thin-layer chromatography was carried out on glass sheets precoated with silica gel GF (2 mm, Analtech). Flash column chromatography was performed on Merck silica gel 60 (230 - 400 mesh). Melting points were determined in open capillary tubes on a Med-Temp apparatus and are uncorrected.

General Procedure for the Synthesis of Bromoketones:

To a cooled solution of the ketone (1 equivalent) in acetic acid was slowly added bromine (1 equivalent). The reaction mixture was stirred at room temperature for 3 hours. The solvent was evaporated and the residue

76

dissolved in dichloromethane, washed with saturated sodium bicarbonate and water. The organic phase was dried over sodium sulfate. Evaporation of the combined decolorized organic phase afforded a light yellow oil in > 80% yield in most cases.

Similarly, the following were prepared:

6-Bromo-3-fluoro-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one was obtained as a light yellow oil in 97% yield from 8-fluoro 1-benzosuberone and bromine: ^1H NMR δ 7.30 (dd, 1H, $JJ = 2.7, 9.0$ Hz), 7.20-7.09 (m, 2H), 4.85 (dd, 1H, $JJ = 4.2, 7.5$ Hz), 3.06-2.97 (m, 1H), 2.91-2.81 (m, 1H), 2.43-2.24 (m, 2H), 2.05-1.97 (m, 2H).

6-Bromo-2-methoxy-6,7,8,9-tetrahydro-5H-benzo[a]-cyclohepten-5-one was obtained as a light brown oil in 97% yield from 2-methoxy-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one and bromine: ^1H NMR δ 7.68 (d, 1H, $J = 8.6$ Hz), 6.80 (dd, 1H, $J = 2.5, 8.6$ Hz), 6.70 (d, 1H, $J = 2.5$ Hz), 4.87 (dd, 1H, $J = 4.3, 7.9$ Hz), 3.85 (s, 3H), 3.10-2.99 (m, 1H), 2.94-2.82 (m, 1H), 2.40-2.30 (m, 2H), 2.0 (m, 2H).

2-Methoxy-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one: To a homogeneous mixture of phosphorus pentoxide and 85% phosphorus acid at 100 °C, was added 5-(3-methoxyphenyl)pentanoic acid in small portions over 20 minutes. The vigorously stirred mixture was heated at 100 °C for 6 hours. Then it was poured on ice-water, and extracted with ether (3 times). The organic phase was washed with water and dried over anhydrous sodium sulfate. Evaporation of the combined organic phase afforded a light yellow oil in more than 95% yield; ^1H NMR δ 7.77 (d,

77

1H, $J = 8.6$ Hz), 6.80 (dd, 1H, $J = 2.5, 8.6$ Hz), 6.69 (d, 1H, $J = 2.5$ Hz), 3.84 (s, 3H), 2.90 (t, 2H, $J = 6.2$ Hz), 2.71 (t, 2H, $J = 5.9$ Hz), 1.91-1.76 (m, 4H).

5 5-(3-Methoxyphenyl)pentanoic acid. A mixture of ethyl 5-(3-methoxyphenyl)pentanoate (1 equivalent) and sodium hydroxide (3 equivalents) in THF/H₂O was heated at reflux overnight. THF was evaporated, the aqueous phase acidified with 6N HCl to pH < 5, and then extracted with a mixture
10 of chloroform : isopropyl alcohol (3 : 1, 5 times). The organic phase was dried over anhydrous sodium sulfate. A light yellow oil was obtained in more than 85% yield after removal of the solvent: ¹H NMR δ 7.20 (m, 1H), 6.79-6.73 (m, 3H), 3.80 (s, 3H), 2.62 (m, 2H), 2.38 (m, 2H), 1.68
15 (m, 4H).

Ethyl 5-(3-methoxyphenyl)pentanoate. A reaction mixture comprising ethyl (*E*) 5-(3-methoxyphenyl) 4-pentenoate and 5% palladium on carbon in methanol was stirred under 300
20 psi H₂ overnight. A light yellow oil was obtained after filtration and evaporation: ¹H NMR δ 7.20 (m, 1H), 6.78-6.72 (m, 3H), 4.12 (q, 2H, $J = 7.1$ Hz), 3.80 (s, 3H), 2.60 (m, 2H), 2.32 (m, 2H), 1.66 (m, 4H), 1.25 (t, 3H, $J = 7.1$ Hz).

25 Ethyl (*E*) 5-(3-methoxyphenyl) 4-pentenoate. To a refluxed solution of 3-iodoanisole (5.0 g, 21.37 mmol, 1 equivalent), tetrakis(triphenylphosphine)-palladium(0) ([C₆H₅)₃P]₄Pd, 740 mg, 0.64 mmol, 0.03 equivalent), and
30 triethyl amine (6.0 ml, 42.74 mmol, 2.0 equivalents) in CH₃CN/THF, was added 1.0 g of ethyl 4-pentenoate every two hours (total 3.3 g, 25.64 mmol, 1.2 equivalents). After the addition was complete, the reaction mixture was stirred at reflux overnight. Solvent was removed and the

78

dark brown residue dissolved in 5% HCl and extracted with methylene chloride (3 times). The combined organic extracts were washed with saturated sodium bicarbonate. A dark brown oil was obtained after the organic solvent was removed. The crude product was purified by flash column (silica gel, chloroform : hexane = 1 : 5) to afford 3.6 g of a light yellow oil in 72% yield: ^1H NMR δ 7.21 (t, 1H, J = 7.9 Hz), 6.93 (m, 1H), 6.87 (m, 1H), 6.76 (dd, 1H, J = 2.6, 8.1 Hz), 6.40 (d, 1H, J = 18 Hz), 6.25-6.16 (m, 1H), 4.15 (q, 2H, J = 7.2 Hz), 3.81 (s, 3H), 2.50 (m, 4H), 1.26 (t, 3H, J = 7.2 Hz).

4-Bromo-9-methoxy-3,4-dihydro-1-benzoxepin-5(2H)-one was obtained as a light yellow oil in 100% yield from 9-methoxy-3,4-dihydro-1-benzoxepin-5(2H)-one and bromine: ^1H NMR δ 7.32 (t, 1H, J = 4.8 Hz), 7.06 (d, 2H, J = 5.1 Hz), 4.99 (dd, 1H, J = 7.0, 8.4 Hz), 4.60-4.53 (m, 1H), 4.20-4.11 (m, 1H), 3.90 (s, 3H), 3.02-2.91 (m, 1H), 2.56-2.45 (m, 1H).

9-Methoxy-3,4-dihydro-1-benzoxepin-5(2H)-one. To PPA in toluene at 100 °C, was added 4-(2-methoxyphenoxy)butanoic acid and the mixture stirred overnight. The mixture was poured onto ice-water, and extracted with ether (3 times). The combined organic phase was washed with water and dried over anhydrous sodium sulfate. Evaporation of the organic phase afforded a light yellow oil in 10% yield: ^1H NMR δ 7.33 (dd, 1H, J = 4.4, 5.1 Hz), 7.03 (d, 2H, J = 4.3 Hz), 4.31 (t, 2H, J = 6.6 Hz), 3.90 (s, 3H), 2.90 (t, 2H, J = 6.9 Hz), 2.22 (p, 2H, J = 6.8 Hz).

4-(2-Methoxyphenoxy)butanoic acid. To the mixture of guaiacol (2-methoxyphenol, 1.0 equivalent) and γ -butyrolactone, cooled in an ice-water bath, was added

potassium ethoxide in small portions. The reaction mixture was heated at 110 °C overnight, and then at 155 °C for an additional 10 hours. The mixture was then poured into ice water, basified to pH 9.5 with saturated sodium bicarbonate, and extracted with chloroform. The aqueous phase was separated, acidified to pH 2.5 with 2N HCl, and extracted with a mixture of chloroform : isopropyl alcohol (3 : 1). The organic phase was dried over anhydrous sodium sulfate and evaporated to afford 4-(2-methoxyphenoxy)butanoic acid as a light brown oil in 55% yield. The crude product was used for next reaction without further purification: ¹H NMR δ 6.90 (m, 4H), 4.08 (t, 2H, *J* = 6.2 Hz), 3.85 (s, 3H), 2.62 (t, 2H, *J* = 7.2 Hz), 2.16 (p, 2H, *J* = 6.6 Hz).

3-Bromo-2,3-dihydro-4*H*-chromen-4-one was obtained as a yellow oil in 100% yield from 4-chromanone and bromine. ¹H NMR δ 7.93 (dd, 1H, *J* = 1.5, 7.8 Hz), 7.53 (td, 1H, *J* = 1.8, 7.8 Hz), 7.11-7.04 (m, 2H), 4.64 (m, 3H).

General Procedure for the Synthesis of Thioureas:

A protected diamine such as N-Boc-1,4-diaminobutane or N-Boc-1,5-diaminopentane (1 equivalent) was dissolved in tetrahydrofuran and stirred at room temperature. Benzoyl thioisocyanate (1 equivalent) was added dropwise to the reaction mixture. The resulting mixture was stirred at room temperature for 24 hours, and the solvent was removed under reduced pressure to afford a yellow oil.

The yellow oil (1 equivalent) was then dissolved in methanol, an aqueous potassium carbonate (3 equivalents) solution added, and the mixture stirred for 48 hours. Water was added to the reaction mixture which was then

extracted with 2x75 ml ethyl acetate. The combined extracts were washed with water, dried with anhydrous sodium sulfate, filtered and the solvent removed under reduced pressure to give the desired thiourea.

5

tert-Butyl 5-[(aminocarbothioyl)amino]pentylcarbamate was obtained as a light yellow wax from tert-butyl 5-[[[(benzoylamino)carbothioyl]amino]-pentylcarbamate. ¹H NMR (CD₃OD) δ 3.44 (m, 1H), 3.10 (m, 1H), 3.01 (t, 2H, J = 6.7 Hz), 1.60-1.31 (m, 6H), 1.41 (s, 9H); 262 (ESMS, MH⁺).

10

tert-Butyl 5-[[[(benzoylamino)carbothioyl]amino]-pentylcarbamate was obtained a light yellow solid in 79% yield from N-BOC-1,5-diaminopentate and benzoyl isothiocyanate; m.p. 90-93 °C; ¹H NMR δ (17001-131-1)

15

tert-Butyl 4-[(aminocarbothioyl)amino]butylcarbamate was obtained as a light yellow wax from tert-butyl 4-[[[(benzoylamino)carbothioyl]amino]-butylcarbamate. ¹H NMR (CD₃OD) δ 3.48 (m, 1H), 3.10 (m, 1H), 3.05 (t, 2H, J = 6.5 Hz), 1.60 (m, 4H), 1.42 (s, 9H); 248 (ESMS, MH⁺).

20

tert-Butyl 4-[[[(benzoylamino)carbothioyl]amino]-butylcarbamate was obtained as a light brown oil in 93% yield from N-BOC-1,4-diaminobutane and benzoyl isothiocyanate.

25

trans-tert-Butyl-{4-[(aminocarbothioyl)amino]-cyclohexyl}methylcarbamate was obtained as a light yellow wax from trans-tert-butyl (4-[[[(benzoylamino)carbothioyl]amino]cyclohexyl)-methylcarbamate. ¹H NMR (CD₃OD) δ 3.92 (m, 1H), 2.86 (m, 2H), 2.00 (m, 2H), 1.76 (m, 2H), 1.41 (s, 9H), 1.37 (m, 1H), 1.06 (m, 4H); 288 (ESMS, MH⁺).

30

trans-*tert*-Butyl (4-{[(benzoylamino)carbothioyl]-amino}cyclohexyl)-methylcarbamate was obtained as a yellow solid in 97% yield from *tert*-butyl 4-aminocyclohexylmethylcarbamate and benzoyl isothiocyanate.

trans-*tert*-Butyl 4-aminocyclohexylmethylcarbamate was obtained in more than 95 % yield by hydrogenation of benzyl 4-{[(*tert*-butoxycarbonyl)amino]methyl} cyclocarbamate.

Benzyl-4-[[[*tert*-butoxycarbonyl]amino]methyl]cyclohexylcarbamate: To a stirred suspension of 4-[[(*tert*-butoxycarbonyl)amino]methyl]cyclohexanecarboxylic acid (Maybridge Chemical Co., Ltd.) (45g) and diphenylphosphoryl azide (44 ml) in toluene (600 ml) was added triethylamine (32 ml) over a period of 20 min whilst maintaining the internal temperature at -10-0 C. The mixture was slowly warmed and then stirred at 70 C for 4 h. After cooling to 40 C, benzyl alcohol (36 ml) was added and the reaction mixture heated at reflux for 20 h. The cold reaction mixture was washed with water and brine and dried over anhydrous magnesium sulfate. Removal of the solvent and recrystallization of the organic residue from ethyl acetate and diethyl ether gave the title compound, benzyl-4-[[[*tert*-butoxycarbonyl]amino]methyl]cyclohexylcarbamate as a white solid, m.p. 129-131 C.

trans-*tert*-Butyl {4-[(aminocarbothioyl)amino]cyclohexyl}carbamate was obtained as a yellow solid from *trans*-*tert*-Butyl 4-(benzoylamino)carbothioylamino}cyclohexyl)-carbamate: ¹H NMR (CD₃OD) δ 3.94 (m, 1H),

82

3.30 (m, 1H), 2.00 (m, 2H), 1.90 (m, 2H), 1.41 (s, 9H), 1.26 (m, 4H); 274 (ESMS, MH⁺).

trans-*tert*-Butyl-4-{[(benzoylamino)carbothioyl]-

5 amino)cyclohexyl)carbamate was obtained as a white solid in 66% yield from *tert*-butyl 4-aminocyclohexylcarbamate and benzoyl isothiocyanate.

10 *trans*-*tert*-Butyl 4-aminocyclohexylcarbamate was obtained as a light yellow wax in more than 95% yield by hydrogenation of benzyl 4-[(*tert*-butoxycarbonyl)-amino]cyclohexylcarbamate.

15 *trans*-Benzyl 4-{[(aminocarbothioyl)amino]methyl}cyclohexylcarbamate was obtained as a yellow solid in 71% yield from *trans*-Benzyl-4-{[(Benzoylamino) carbothioyl]-amino}methyl)-cyclohexylcarbamate; 322 (ESMS, MH⁺).

20 *trans*-Benzyl-4-({[(benzoylamino)carbothioyl]-amino}methyl)-cyclohexylcarbamate was obtained as a yellow solid from benzyl 4-(aminomethyl)cyclohexylcarbamate and benzoyl isothiocyanate.

25 *trans*-Benzyl 4-(aminomethyl)cyclohexylcarbamate was obtained as a white solid in more than 95% yield by stirring benzyl-4-{[(*tert*-butoxycarbonyl)amino]-methyl}cyclocarbamate in 2N HCl (made from 1 : 1 of EtOAc and 4N HCl in dioxane).

30 **General Procedure for the Synthesis of Tricyclic Thiazoles:**

A mixture of a bromoketone such as 6-bromo-3-fluoro-6,7,8,9-tetrahydro-5*H*-benzo[a]cyclohepten-5-one (1.2

83

equivalent), thiourea (1 equivalent), and diisopropylethylamine (2 equivalents) in anhydrous ethanol was stirred and heated at reflux overnight. The solvent was evaporated and the brown residue dissolved in dichloromethane and washed with saturated aqueous sodium bicarbonate. The mixture was extracted with dichloromethane three times. The combined extracts were dried over anhydrous sodium sulfate and the solvent removed to afford a crude product which was purified by flash column chromatography (silica gel).

General Procedure for the Deprotection of BOC-Protected Amines:

The Boc protected amine was dissolved in Et₂O. The same volume of 4N HCl in dioxane was added to make a 2N solution of HCl. The reaction mixture was stirred at room temperature overnight, and the solvent removed under reduced pressure to afford the desired product as its HCl salt.

tert-Butyl-5-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-d][1,3]thiazol-2-yl)amino]-pentylcarbamate was obtained as a light yellow oil in 80% yield from 6-bromo-3-fluoro-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one and *tert*-butyl 5-[(aminocarbothioyl)amino]-pentylcarbamate: ¹H NMR δ 7.70 (dd, 1H, *J* = 2.7, 10.8 Hz), 7.05 (dd, 1H, *J* = 6.0, 8.4 Hz), 6.83 (td, 1H, *J* = 2.7, 8.4 Hz), 5.12 (m, 1H), 4.53 (m, 1H), 3.26 (q, 2H, *J* = 5.7 Hz), 3.14 (m, 2H), 2.89 (t, 2H, *J* = 6.9 Hz), 2.79 (m, 2H), 2.09 (m, 2H), 1.68 (m, 4H), 1.50 (m, 2H), 1.44 (s, 9H); 420 (ESMS, MH⁺).

N-(5-Aminopentyl)-*N*-(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-yl)amine

dihydrogen chloride was obtained as a yellow solid in more than 95% yield from deprotection of *tert*-butyl 5-[(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-yl)amino]pentylcarbamate by using 2N HCl (made from 1 : 1 of ethyl acetate and 4N HCl in dioxane);

¹H NMR (CD₃OD) δ 7.38-7.31 (m, 2H), 7.10 (td, 1H, *J* = 2.4, 8.4 Hz), 3.71 (m, 2H), 3.49 (t, 2H, *J* = 6.9 Hz), 2.95 (m, 2H), 2.78 (m, 2H), 2.17 (m, 2H), 1.85-1.68 (m, 4H), 1.54 (m, 2H); 320 (ESMS, MH⁺).

tert-Butyl 4-[(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-yl)amino]-

butylcarbamate was obtained as a yellow oil in 31% yield from

6-bromo-3-fluoro-6,7,8,9-tetrahydro-5*H*-benzo[*a*]cyclohepten-5-one and *tert*-butyl 4-[(aminocarbothioyl)amino]-butylcarbamate: ¹H NMR δ 7.70 (dd, 1H, *J* = 3.0, 10.8 Hz), 7.05 (dd, 1H, *J* = 6.0, 8.4 Hz), 6.82 (td, 1H, *J* = 3.0, 8.3 Hz), 5.14 (m, 1H), 4.59 (m, 1H), 3.30 (m, 2H), 3.18 (m, 2H), 2.86 (t, 2H, *J* = 6.9 Hz), 2.79 (m, 2H), 2.10 (m, 2H), 1.74-1.53 (m, 4H), 1.44 (s, 9H).

N-(4-Aminobutyl)-*N*-(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-yl)amine hydrogen chloride was obtained as a yellow solid in more than 95% yield by deprotection of *tert*-butyl 4-[(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-

yl)amino]butylcarbamate with 2N HCl (made from 1 : 1 of 4N HCl in dioxane and ethyl acetate): ¹H NMR (CD₃OD) δ 7.39-7.31 (m, 2H), 7.10 (td, 1H, *J* = 3.0, 8.4 Hz), 3.54 (m, 2H), 2.97 (m, 2H), 2.79 (m, 4H), 2.18 (m, 2H), 1.83 (m, 4H). (no MS).

trans-*tert*-Butyl-{4-[(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta-[1,2-*d*][1,3]thiazol-2-yl)amino]cyclohexyl}methylcarbamate was obtained as a yellow oil in 80% yield from 6-bromo-3-fluoro-6,7,8,9-tetrahydro-5*H*-benzo[*a*]cyclohepten-5-one and *trans*-*tert*-butyl {4-[(aminocarbothioyl)-amino]cyclohexyl}methylcarbamate: ^1H NMR δ 7.69 (dd, 1H, J = 2.8, 10.8 Hz), 7.05 (dd, 1H, J = 5.9, 8.4 Hz), 6.82 (td, 1H, J = 2.7, 8.3 Hz), 4.87 (m, 1H), 4.61 (m, 1H), 3.28 (m, 1H), 3.00 (m, 1H), 2.85 (t, 2H, J = 6.9 Hz), 2.79 (m, 2H), 2.22 (m, 2H), 2.08 (m, 2H), 1.85 (m, 2H), 1.74 (m, 1H), 1.44 (s, 9H), 1.45-1.02 (m, 4H); 446 (ESMS, MH^+).

N-(4-Aminomethyl)cyclohexyl]-*N*-(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-yl)amine hydrogen chloride was obtained as a yellow solid in more than 95% yield by deprotection of *tert*-butyl {4-[(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-yl)amino]cyclohexyl}methylcarbamate: ^1H NMR (CD_3OD) δ 7.34 (td, 2H, J = 2.7, 9.6 Hz), 7.11 (m, 1H), 3.61 (m, 1H), 2.82-2.76 (m, 6H), 2.18 (m, 4H), 1.96 (m, 2H), 1.69 (m, 1H), 1.50 (m, 2H), 1.24 (m, 2H); 346 (ESMS, MH^+).

trans-*tert*-Butyl-4-[(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta-[1,2-*d*][1,3]thiazol-2-yl)amino]cyclohexylcarbamate was obtained as a yellow solid in 59% yield from 6-bromo-3-fluoro-6,7,8,9-tetrahydro-5*H*-benzo[*a*]cyclohepten-5-one and *trans*-*tert*-butyl {4-[(aminocarbothioyl)-amino]cyclohexyl}-carbamate: ^1H NMR δ 7.69 (dd, 1H, J = 2.7, 10.8 Hz), 7.04 (dd, 1H, J = 6.0, 8.1 Hz), 6.82 (td, 1H, J = 2.7, 8.2 Hz), 4.95 (m, 1H), 4.45 (m, 1H), 3.39 (m, 2H), 2.84 (t, 2H, J = 6.9 Hz),

86

2.77 (m, 2H), 2.10 (m, 2H), 2.07 (m, 4H), 1.44 (s, 9H), 1.28 (m, 4H); 432 (ESMS, MH⁺).

N-(4-Aminocyclohexyl)-*N*-(9-fluoro-5,6-dihydro-4*H*-

5 benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-yl)amine

hydrogen chloride was obtained as a yellow solid in more than 95% yield by deprotection of *tert*-butyl 4-[(9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[1,2-

10 *d*][1,3]thiazol-2-yl)amino]-cyclohexylcarbamate with 2N HCl (made from 1 : 1 of ethyl acetate and 4N HCl in dioxane):

¹H NMR (CD₃OD) δ 7.35 (m, 2H), 7.12 (m, 1H), 3.68 (m, 1H), 3.17 (m, 1H), 2.76 (m, 4H), 2.25-2.13 (m, 6H), 1.59 (m, 4H); 332 (ESMS, MH⁺).

15 *trans*-*tert*-Butyl-{4-[(8-Methoxy-5,6-dihydro-4*H*-

benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-yl)amino]-

cyclohexyl}methylcarbamate was obtained as a light yellow oil in 46% yield from 6-bromo-2-methoxy-6,7,8,9-tetrahydro-5*H*-benzo[*a*]cyclohepten-5-one and *trans*-*tert*-

20 butyl-{4-[(aminocarbothioyl)amino]-

cyclohexyl}methylcarbamate; 458 (ESMS, MH⁺).

N-[4-(Aminomethyl)cyclohexyl]-8-methoxy-5,6-dihydro-4*H*-

25 benzo[6,7]cyclohepta[1,2-*d*][1,3]thiazol-2-amine hydrogen chloride was obtained as a yellow solid in more than 95% yield by deprotection of *trans*-*tert*-butyl-{4-[(8-methoxy-5,6-dihydro-4*H*-benzo[6,7]-cyclohepta[1,2-*d*][1,3]thiazol-2-

yl)amino]cyclohexyl}-methylcarbamate) with 2N HCl (made from 1 : 1 of ethyl acetate and 4N HCl in dioxane): ¹H NMR

30 (CD₃OD) δ 7.48 (m, 1H), 6.91 (m, 2H), 3.81 (s, 3H), 3.54 (m, 1H), 2.83-2.70 (m, 6H), 2.10 (m, 4H), 1.95 (m, 2H), 1.70 (m, 1H), 1.50 (m, 2H), 1.24 (m, 2H); 358 (ESMS, MH⁺).

Benzyl-{4-[(8-Methoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-d][1,3]thiazol-2-yl)amino]methyl}-cyclohexylcarbamate was obtained as a light yellow oil in 60% yield from 6-bromo-2-methoxy-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one and trans-benzyl-4-[[aminocarbothioyl)amino]methyl}-cyclohexylcarbamate: ¹H NMR δ 7.88 (d, 1H, J = 8.6 Hz), 7.35 (m, 5H), 6.79 (dd, 1H, J = 2.4, 8.8 Hz), 6.68 (d, 1H, J = 2.6 Hz), 5.30 (m, 1H), 5.08 (s, 2H), 4.62 (m, 1H), 3.81 (s, 3H), 3.45 (m, 1H), 3.08 (m, 2H), 2.83 (t, 2H, J = 7.0 Hz), 2.77 (m, 2H), 2.08 (m, 4H), 1.85 (m, 2H), 1.55 (m, 1H), 1.10 (m, 4H); 492 (ESMS, MH⁺).

N-[4-(Aminocyclohexyl)methyl]-8-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-d][1,3]thiazol-2-amine trifluoroacetic acid salt was obtained as a brown syrup in more than 95% yield by refluxing benzyl {4-[(8-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-d][1,3]thiazol-2-yl)amino]methyl}cyclohexylcarbamate in TFA/H₂O (95 : 5) for 1 hour: ¹H NMR (CD₃OD) δ 7.47 (d, 1H, J = 8.4 Hz), 6.86 (m, 2H), 3.80 (s, 3H), 3.28 (m, 2H), 3.07 (m, 1H), 2.79-2.73 (m, 4H), 2.17-2.05 (m, 4H), 1.97 (m, 2H), 1.75 (m, 1H), 1.42 (m, 2H), 1.23 (m, 2H); 358 (ESMS, MH⁺).

tert-Butyl {4-[(7-Methoxy-4,5-dihydro[1]benzoxepino-[5,4-d][1,3]thiazol-2-yl)amino]cyclohexyl}methyl-carbamate was obtained as a light yellow oil in 69 % yield from 4-bromo-9-methoxy-3,4-dihydro-1-benzoxepin-5(2H)-one and trans-tert-butyl {4-[[aminocarbothioyl)amino]cyclohexyl}methylcarbamate: ¹H NMR δ 7.85 (dd, 1H, J = 1.5, 8.2 Hz), 7.02 (t, 1H, J = 8.1 Hz), 6.81 (dd, 1H, J = 1.4, 8.0 Hz), 4.83 (m, 1H), 4.60 (m, 1H), 4.40 (t, 2H, J = 5.1 Hz), 3.88 (s, 3H), 3.29 (m,

88

1H), 3.18 (t, 2H, $J = 5.1$ Hz), 3.00 (m, 2H), 2.24 (m, 2H), 1.85 (m, 2H), 1.60 (m, 1H), 1.45 (s, 9H); 460 (ESMS, MH^+).

5 N-[4-(Aminomethyl)cyclohexyl]-7-methoxy-4,5-dihydro-[1]-benzoxepino[5,4-d][1,3]thiazol-2-amine hydrogen chloride was obtained as a yellow solid in more than 95% yield by stirring tert-butyl {4-[(7-methoxy-4,5-dihydro[1]benzoxepino[5,4-d][1,3]thiazol-2-yl)amino]cyclohexyl}methylcarbamate in 2N HCl (made from 1 : 1 of ethyl acetate and 4N HCl in dioxane) for 3hr: 1H NMR (CD₃OD) δ 7.25 (m, 1H), 7.20-7.06 (m, 2H), 4.32 (m, 2H), 3.85 (s, 3H), 3.56 (m, 1H), 3.19 (m, 2H), 2.86 (m, 2H), 2.21 (m, 2H), 1.94 (m, 2H), 1.73 (m, 1H), 1.55 (m, 2H), 15 1.27 (m, 2H); 360 (ESMS, MH^+).

tert-Butyl [4-(4H-Chromeno[4,3-d][1,3]thiazol-2-yiamino)cyclohexyl]-methylcarbamate was obtained as a light yellow oil from 3-Bromo-2,3-dihydro-4H-chromen-4-one and trans-tert-butyl {4-[(aminocarbothioyl)amino]cyclohexyl}methylcarbamate: 1H NMR δ 7.58 (dd, 1H, $J = 1.5, 7.5$ Hz), 7.12 (m, 1H), 6.96 (m, 1H), 6.87 (dd, 1H, $J = 0.6, 8.1$ Hz), 5.07 (m, 1H), 4.60 (m, 1H), 3.29 (m, 1H), 3.00 (m, 2H), 2.07 (m, 2H), 1.85 (m, 2H), 1.50 (m, 1H), 1.45 (s, 9H), 1.25 (m, 2H), 1.05 (m, 2H); 416 (ESMS, MH^+).

30 N-[4-(Aminomethyl)cyclohexyl]-4H-chromeno[4,3-d][1,3]thiazol-2-amine hydrogen chloride was obtained as a yellow solid in more than 95% yield by stirring tert-butyl [4-(4H-chromeno[4,3-d][1,3]thiazol-2-yiamino)cyclohexyl]methylcarbamate in 2N HCl (made from 1 : 1 of ethyl acetate and 4N HCl in dioxane) for 3hr: 1H NMR (CD₃OD) δ 7.65 (dd, 1H, $J = 1.3, 7.8$ Hz), 7.29 (td, 1H, $J =$

1.5, 7.8 Hz), 7.06 (td, 1H, $J = 1.0, 7.7$ Hz), 6.97 (dd, 1H, $J = 0.9, 8.4$ Hz), 5.23 (s, 2H), 3.73 (m, 1H), 2.84 (m, 2H), 2.21 (m, 2H), 1.95 (m, 2H), 1.70 (m, 1H), 1.50 (m, 2H), 1.25 (m, 2H); 316 (ESMS, MH^+).

5

trans-N2-(4-(2-Methoxyethoxycarbonyl)aminomethyl)-cyclohexyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]-cyclohepta-[d][1,3]-thiazol-2-amine:

10 a. *t*-Butyl-*trans*-N-(4-(Benzoylaminocarbothioylamino)-cyclohexyl)methyl-carbamate: A solution of *t*-butyl *trans*-N-(4-aminocyclohexyl)methylcarbamate (4.683g, 20.51mmol) and benzoylisothiocyanate (3.77g, 23.1mmol) in THF (180 mL) was stirred at room temperature for 20 hours. The
15 solvent was removed in vacuo to yield a golden viscous oil. The oil was triturated with hexane (250 mL) and the resulting solid was collected by filtration and washed with hexane to yield 91% (7.32g) of the product as an off-white solid: 1H NMR ($CDCl_3$) δ 1.20 (6H, m), 1.44 (9H, s),
20 1.83 (2H, m), 2.25 (2H, m), 3.01 (2H, t, $J = 6.4$ Hz), 4.20 (1H, m), 4.62 (1H, m), 7.51 (2H, t, $J = 7.9$ Hz), 7.60 (1H, t, $J = 7.4$ Hz), 7.81 (2H, d, $J = 7.2$ Hz), 8.93 (1H, s); 392 (ESMS, MH^+).

25 b. *t*-Butyl *trans*-N-(4-(aminocarbothioylamino)-cyclohexyl)methylcarbamate: To a solution of *t*-butyl *trans*-N-(4-(benzoylaminocarbothioylamino)cyclohexyl)-methylcarbamate (7.30g, 18.6mmol) in MeOH (110 mL) was
30 added a solution of potassium carbonate (5.15 g, 37.3 mmol) in water (50 mL). The solution was stirred at room temperature for 18 hours. The solvent was removed in vacuo and the residue was extracted several times with EtOH. The combined EtOH extracts were evaporated to dryness to yield 6.15g of an off-white solid. The solid

90

was dissolved in acetone (200 mL) and stirred for 20 minutes. The solution was filtered by suction to remove any insoluble salts and the filtrate evaporated to yield 98% (5.23 g) of the desired product as a slightly yellow solid: ^1H NMR (CDCl_3) δ 1.10 (6H, m), 1.43 (9H, s), 1.80 (2H, m), 2.10 (2H, m), 2.95 (2H, t, $J = 6.4$ Hz), 4.75 (1H, broad), 6.03 (2H, s), 6.64 (1H, broad); 288 (ESMS, MH^+).

c. 6-Bromo-3-fluoro-6,7,8,9-tetrahydro-5H-benzo[a]-cyclohepten-5-one: To a solution of 8-fluoro-1-benzosuberone (20.00 g, 112.2 mmol) in glacial acetic acid (125 ml) was added bromine (5.8 mL, 112.6 mmol) dropwise. After addition was complete, the solution was poured into water (400 ml) and extracted with EtOAc (500 ml). The EtOAc layer was separated and washed with water (500 ml). The organic layer was dried over MgSO_4 , filtered and evaporated to yield 100% (28.9g) of the desired product as a viscous oil: ^1H NMR (CDCl_3) δ 2.00 (2H, m), 2.33 (2H, m), 2.84 (1H, m), 3.00 (1H, m), 4.83 (1H, m), 7.13 (2H, m), 7.29 (1H, dd).

d. *t*-Butyl *trans*-N-(4-(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-ylamino)-cyclohexyl)methylcarbamate: A solution of *t*-butyl *trans*-N-(4-(aminocarbothioylamino)cyclohexyl)-methylcarbamate (22.4g, 78.0 mmol), 6-bromo-3-fluoro-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one (20.05g, 78.0 mmol) and *N,N*-diisopropylethylamine (22.5 ml) in absolute ethanol (260 ml) was heated at reflux for 3 hours. The solvent was removed *in vacuo*, and the resulting viscous golden oil was taken up in EtOAc (400 ml) and washed with water (600 ml). The EtOAc layer was separated and evaporated to yield 39.0 g of a viscous golden oil. The oil was chromatographed over silica gel eluting with EtOAc to yield 34.5g of the

91

desired product (99% yield): ^1H NMR (CDCl_3) δ 1.18 (6H, h, $J = 12.2$ Hz), 1.44 (9H, s), 1.81 (2H, m), 2.22 (2H, m), 2.79 (2H, m), 2.86 (2H, t, $J = 7.1$ Hz), 3.00 (2H, t, $J = 6.3$ Hz), 3.25 (1H, broad), 4.62 (1H, broad), 4.86 (1H, broad), 6.83 (1H, dt, $J = 2.9$ Hz), 7.04 (1H, m), 7.70 (1H, dd, $J = 2.8$ Hz).

e. *trans*-N2-(4-Aminomethyl)cyclohexyl-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine hydrochloride: A solution of t-butyl *trans*-N-(4-(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-[d][1,3]thiazol-2-ylamino)cyclohexyl)-methylcarbamate (34.5g, 77.0 mmol) in EtOAc (200 ml) and 4N HCl in dioxane (100 ml) was stirred at room temperature for 2 hours. The resulting solid was collected by filtration, washed with Et_2O and dried under vacuum to yield 25.9g of the desired product as a slightly yellow solid. An additional 1.43g of the product was obtained by allowing the filtrate to stand in a closed flask for several days. The overall yield was 85%: ESMS $m/e = 346$ (MH^+); ^1H NMR (CD_3OD) δ 1.25 (2H, m), 1.51 (2H, m), 1.70 (1H, broad), 1.95 (2H, m), 2.18 (4H, m), 2.80 (6H, m), 3.63 (1H, broad), 7.11 (1H, dt, $J = 2.5\text{Hz}, 8.4$ Hz), 7.36 (2H, m).

f. *trans*-N2-(4-(2-Methoxyethoxycarbonyl)aminomethyl)-cyclohexyl-9-fluoro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[d][1,3]-thiazol-2-amine: A solution of *trans*-N2-(4-aminomethyl)cyclohexyl-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine hydrochloride (0.30 g, 0.72 mmol) and triethylamine (1.0 ml) in anhydrous THF (3 ml) was treated dropwise with chloroformic acid 2-methoxyethyl ester (0.114 g, 0.82 mmol). The solution was stirred at room temperature for 2 hours. The solvent was removed *in vacuo* and the residue

92

chromatographed over silica gel eluting with 1:1 EtOAc/Hexane to yield the desired product (80 mg, 25% yield) as a highly viscous colorless oil: 448 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 1.11 (4H, h, J = 9.8 Hz), 1.42 (1H, broad), 1.80 (2H, m), 2.04 (2H, m), 2.20 (2H, m), 2.77 (2H, m), 2.84 (2H, t, J = 6.9 Hz), 3.04 (2H, t, J = 6.4 Hz), 3.24 (1H, broad), 3.38 (3H, s), 3.58 (2H, m), 4.22 (2H, m), 4.90 (1H, broad), 5.00 (1H, d), 6.81 (1H, dt, J = 2.8 Hz), 7.03 (1H, m), 7.68 (1H, dd, J = 2.7 Hz, 10.9 Hz).

Similarly prepared were:

N2-(5-Aminopent-1-yl)-9-fluoro-5,6-dihydro-4H-

benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine hydrochloride:

Yellow solid; m.p. 220 °C (dec.); 320 (ESMS, MH⁺); ¹H NMR (CD₃OD) δ 7.34 (m, 2H), 7.10 (td, 1H, J = 2.4, 8.4 Hz), 3.49 (t, 2H, J = 7.0 Hz), 2.95 (t, 2H, J = 7.7 Hz), 2.78 (m, 4H), 2.17 (m, 2H), 1.82-1.67 (m, 4H), 1.54 (m, 2H).

General Procedure for the Derivatization of Amines with Carboxylic Acid and Sulfonic Acid Derivatives:

An amine such as N-[4-(aminomethyl)cyclohexyl]-4H-chromeno[4,3-d][1,3]thiazol-2-amine (0.305 mmol) was dissolved in 2ml CH₂Cl₂ with 1-3 equivalents of diisopropylethylamine. The appropriate sulfonyl chloride, acid chloride, isocyanate or carbamyl chloride (1-3 equivalents) was then added dropwise. The reaction solution was stirred at room temperature for 1-3 days, quenched with water and washed with 10% NaHCO₃, extracted with EtOAc, and then dried over Na₂SO₄. The crude product was purified using column chromatography or preparative TLC.

General Procedure for the Formation of Formamides:

The following sequence for the preparation of formamides is typical:

5

N2-[4-(aminomethyl)phenyl]-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine-tert-butyl N-(4-aminobenzyl)carbamate: To a stirred solution of 4-amino-benzylamine (7.5 g, 61.39 mmol) in anhydrous THF (100 ml) under N₂ was added, in portions, di-*t*-butyl dicarbonate (13.5 g, 61.9 mmol). The solution was stirred at room temperature under N₂ for 90 minutes. The solvent was removed in vacuo to yield 100% (13.7 g) of the Boc-protected amine as a white solid: m.p. 70-72°C. ¹H-NMR (CDCl₃) δ 1.43 (9H, s), 3.62 (2H, broad), 4.18 (2H, d), 4.77 (1H, broad), 6.63 (2H, d), 7.06 (2H, d).

15

tert-butyl N-(4-{[(benzoylamino)carbothioyl]amino}-benzyl)carbamate: To a stirred solution of N2-[4-(aminomethyl)phenyl]-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine-tert-butyl N-(4-aminobenzyl)carbamate (6.00 g, 27.0 mmol) in anhydrous THF (50 ml) was added benzoylisothiocyanate (4.43 g, 27.0 mmol). The solution was stirred for 2 hours and the solvent removed in vacuo to yield 100% (10.4 g) of the protected thiourea as a yellow solid: m.p. 161-163°C. ¹H-NMR (CDCl₃) δ 1.45 (9H, s), 4.33 (2H, d), 4.85 (1H, broad), 7.34 (2H, d), 7.60 (5H, m), 7.90 (2H, d), 9.13 (1H, s), 12.58 (1H, s); 386 (ESMS, MH⁺).

25

30

tert-butyl N-{4-[(aminocarbothioyl)amino]benzyl}-carbamate: To a stirred solution of tert-butyl N-(4-{[(benzoylamino)carbothioyl]amino}benzyl)carbamate 10.4 g,

94

27.0 mmol) in MeOH (150 ml) (not completely dissolved) was added a solution of K_2CO_3 (8.5 g, 61.5 mmol) in circa. 15 ml water. The solution was stirred at room temperature for 24 hours and filtered to remove a white precipitate. The filtrate was evaporated *in vacuo* to yield 100% (7.6 g) of an off-white solid.

tert-butyl N-{4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]-cyclo-hepta[d][1,3]thiazol-2-yl)amino]benzyl}-carbamate:

A solution containing tert-butyl N-{4-[(aminocarbothioyl)amino]benzyl}carbamate (7.60 g, 27.0 mmol), 6-bromo-3-fluoro-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one (6.95 g, 27.0 mmol), and diisopropylethylamine (5.0 ml) in EtOH (60 ml) was heated at reflux for 2.5 hours. The solvent was removed *in vacuo* and the residue dissolved in EtOAc (125 ml). The resulting solution was filtered to remove the insoluble (*i*-pr)₂NEt•HBr. The filtrate was transferred to a 250 ml round bottom flask and 4N HCl in dioxane (35 ml) added with stirring. The solution was stirred for 90 minutes and filtered to collect the hydrochloride salt. The solid was washed with EtOAc and then with Et₂O. Upon drying, 86% (11.1 g) of the Boc-protected amine hydrochloride was obtained as a slightly yellow solid: ¹H-NMR (d₄-MeOH) δ 1.43 (9H, s), 2.18 (2H, m), 2.80 (4H, m), 4.25 (2H, s), 7.12 (1H, dt), 7.33 (2H, m), 7.43 (4H, s).

2-(4-aminomethyl)anilino-1-aza-9-fluoro-4,5-dihydro-3-thia-benzo[e]azulene: To a stirred solution of tert-butyl N-{4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[d][1,3]-thiazol-2-yl)amino]benzyl}-carbamate (11.1 g, 23.3 mmol) in MeOH (150 ml) (partially dissolved) was added 4N HCl in dioxane (40 ml). The solution was

95

brought to a brief gentle reflux and the solvent was then removed in vacuo. The resulting solid was re-suspended in Et₂O, filtered and dried to yield 90% (9.25 g) of the product di-hydrochloride as a white solid: m.p. 236-239°C. ¹H-NMR (CDCl₃) δ 1.75 (2H, broad), 2.06 (2H, m), 2.77 (2H, m), 2.83 (2H, t), 3.79 (2H, s), 6.82 (1H, t), 7.04 (1H, m), 7.22 (4H, m), 7.79 (1H, dd), 8.8 (1H, broad); 340 (ESMS, MH⁺).

10 N1-{4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-
[d][1,3]thiazol-2-yl)amino]benzyl}-2-methoxyacetamide: To
a stirred solution of 2-(4-aminomethyl)anilino-1-aza-9-
fluoro-4,5-dihydro-3-thia-benzo[e]azulene (1.50 g, 3.64
mmol) in anhydrous pyridine (10 ml) and
15 diisopropylethylamine (5.0 ml) was added methoxyacetyl
chloride (0.40 ml, 4.38 mmol) dropwise. The solution was
stirred overnight, poured into water (200 ml), and
extracted with EtOAc (200 ml). The EtOAc extract was
evaporated to yield 1.8 g of a viscous golden oil. The
20 oil was chromatographed on silica-gel eluting with EtOAc
to yield 80% (1.2 g) of the desired product as a foam.
The product was converted to its HCl salt in chloroform
using 1M HCl in Et₂O: m.p. 175-178°C; ¹H-NMR (CDCl₃) δ 2.15
(2H, m), 2.86 (4H, m), 3.42 (3H, s), 3.96 (2H, s), 4.50
25 (2H, d, J = 6.1 Hz), 7.00 (1H, t), 7.18-7.40 (7H, m), 7.61
(1H, dd, J = 2.5 Hz, 9.5 Hz); 412 (ESMS, MH⁺).

N-{4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-
[d][1,3]thiazol-2-yl)amino]benzyl}-N-isopropyl-formamide:

30

N2-{4-[(isopropylamino)methyl]phenyl}-9-fluoro-5,6-
dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine: A
solution of 2-(4-aminomethyl)anilino-1-aza-9-fluoro-4,5-
dihydro-3-thia-benzo[e]azulene (free-base) (0.54 g, 1.59

mmol), anhydrous THF (8 ml), anhydrous DMF (4 ml) and 2-iodopropane (1.4 g, 8.24 mmol) was heated to reflux for 1 hour. The solution was then stirred at room temperature overnight. TLC (95% EtOAc/5% 2M NH₃ in MeOH) indicated a major product (R_f ~ 0.4) and some starting material. The solution was poured into water (200 ml) and extracted with EtOAc. The EtOAc layer was evaporated to yield 99% (0.6 g) of the product as a viscous golden oil. ¹H NMR spectrum showed the presence of an isopropyl group (δ 1.08, CDCl₃) as a doublet. The amine was not purified before the next step.

1-aza-9-fluoro-4,5-dihydro-2-(4-(n-(2-propyl)-formamido)-methyl)anilino-3-thia-benzo[e]azulene: To a stirred solution of N2-{4-[(isopropylamino)methyl]phenyl}-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine (0.60 g, 1.57 mmol) in anhydrous THF (10 ml) was added 1H-benzotriazole-1-carboxaldehyde (0.245 g, 1.67 mmol). The solution was stirred for 30 minutes at room temperature, poured into 2N NaOH (60 ml) and extracted with EtOAc. The EtOAc layer was again washed with 2N NaOH (60 ml) and evaporated to yield 0.55 g of a viscous golden oil. The oil was chromatographed on silica-gel eluting with 40% EtOAc/ 60% hexane to yield 23% (0.15 g) of the desired product as a viscous colorless oil. The oil was converted to its HCl salt in chloroform with 1N HCl in Et₂O. Upon evaporation of solvent and trituration with Et₂O, 104 mg of the desired HCl salt was isolated: m.p. 180-183°C. ¹H-NMR (CDCl₃) δ 1.10 (1.5H, d, J = 6.8 Hz), 1.20 (4.5H, d, 6.3 Hz), 2.10 (2H, m), 2.78 (2H, m), 2.83 (2H, m), 3.72 (1H, h), 4.30 (0.5H, s), 4.51 (1.5H, s), 6.82 (1H, m), 7.18 (3H, m), 7.34 (2H, m), 7.76 (1H, d), 8.23 (0.25H, s), 8.39 (1.75H, broad s); 410 (ESMS, MH⁺).

Example 1

trans-N²-(4-Dimethylaminosulfonylaminoethyl)-cyclohexyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]-cyclohepta[d][1,3]thiazol-2-amine

a. *trans*-1-Aminocarbothioylamino-4-(dimethylaminosulfonyl-aminomethyl)-cyclohexane: 4-(Dimethylaminosulfonylamino-methyl)cyclohexylamine (500 mg, 2.12 mmol) was suspended in dry THF (8 mL), cooled in an ice water bath and treated dropwise with benzoylisothiocyanate (290 μ L, 2.16 mmol). The yellow solution was stirred at room temperature overnight and then concentrated. The residual yellow solid was dissolved in chloroform and flash chromatographed over silica gel (49 g) eluting with EtOAc/hexane (2:3) to give a yellow solid (490 mg, 58%). This solid was suspended in MeOH (4 mL), cooled in a water bath and treated with potassium carbonate (170 mg, 1.23 mmol) in water (3 mL). The mixture was stirred at room temperature for 1 hour before more MeOH (2 mL) and water (1.5 mL) were added. After 3 hours, the organic solvent was evaporated and the aqueous layer extracted with dichloromethane and EtOAc. The combined extracts were dried (Na₂SO₄), filtered and concentrated to give a white foam (378 mg).

b. *trans*-N²-(4-Dimethylaminosulfonylaminoethyl)-cyclohexyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]-cyclohepta[d][1,3]-thiazol-2-amine: 6-Bromo-3-fluoro-6,7,8,9-tetrahydro-5*H*-benzo[a]cyclohepten-5-one (69 mg, 0.27 mmol) and *trans*-1-aminocarbothioylamino-4-(dimethylaminosulfonylaminoethyl)-cyclohexane (119 mg, 0.40 mmol) were dissolved in EtOH (4 mL) and heated at reflux overnight. The solvent was evaporated and the

98

residue partitioned between EtOAc and potassium carbonate solution. The organic phase was dried (Na_2SO_4), filtered and concentrated to give a light orange oil (179 mg) which was flash chromatographed over silica gel (14 g) eluting with EtOAc/hexane (2:3) to afford a pale yellow foam (105 mg, 86%): m.p. 60-65 °C; 453 (ESMS, MH^+); ^1H NMR (CDCl_3) δ 1.15 (4H, m), 1.50 (1H, broad), 1.90 (2H, m), 2.07 (2H, m), 2.25 (2H, m), 2.80 (2H, m), 2.82 (6H, s), 2.86 (2H, t, $J = 6.9$ Hz), 2.94 (2H, t, $J = 6.6$ Hz), 3.33 (1H, broad), 4.18 (1H, m), 4.81 (1H, d, $J = 7.6$ Hz), 6.83 (1H, m), 7.06 (1H, dd, $J = 6.0, 8.3$ Hz), 7.69 (1H, dd, $J = 2.8, 10.8$ Hz).

Example 2

1-Aza-9-fluoro-4,5-dihydro-2-{5-(dimethyl-aminosulfonyl-amino)pentyl}amino-3-thia-benzo[e]azulene: 86% yield; 427 (ESMS, MH^+); ^1H NMR (CDCl_3) δ 7.67 (dd, 1H, $J = 10.8, 3.0$ Hz), 7.05 (dd, 1H, $J = 6.0, 8.2$), 6.82 (apparent dt, 1H, $J = 2.8, 8.3$ Hz), 5.74 (broad, 1H), 5.24 (t, 1H, $J = 5.7$ Hz), 3.16 (q, 2H, $J = 4.2$ Hz), 2.97 (q, 2H, $J = 6.3$ Hz), 2.84 (t, 2H, $J = 6.9$ Hz), 2.77 (s, 6H), 2.75 (masked m, 2H), 2.08 (m, 2H), 1.58 (p, 2H, $J = 6.75$ Hz), 1.43 (p, 2H, $J = 6.9$ Hz), 1.26 (p, 2H, $J = 7.2$ Hz).

Example 3

1-Aza-9-fluoro-2-(5-(2-fluorophenyl)sulfonylamino)-pentylamino-4,5-dihydro-3-thia-benzo[e]azulene was obtained as a yellow oil in 36% yield; ^1H NMR δ 7.89 (td, 1H, $J = 1.6, 7.5$ Hz), 7.62 (dd, 1H, $J = 2.7, 10.8$ Hz), 7.55 (m, 1H), 7.29-7.17 (m, 2H), 7.06 (dd, 1H, $J = 6.0, 8.1$ Hz), 6.81 (td, 1H, $J = 3.0, 8.2$ Hz), 5.44 (b, 2H), 3.16 (m, 2H), 2.95 (m, 2H), 2.84 (t, 2H, $J = 6.9$ Hz), 2.08

99

(m, 2H), 1.58-1.25 (m, 6H); 478 (ESMS, MH⁺); ; m.p. (HCl salt) 55-57 °C.

Example 4

5

1-Aza-9-fluoro-4,5-dihydro-2-(5-(1-naphthyl)sulfonyl-amino)-pentylamino-3-thia-benzo[e]azulene: was obtained as a yellow oil in 29% yield; ¹H NMR δ 8.69 (d, 1H, J = 8.7 Hz), 8.25 (dd, 1H, J = 0.9, 7.2 Hz), 8.05 (d, 1H, J = 8.4 Hz), 7.93 (d, 1H, J = 8.1 Hz), 7.68-7.49 (m, 4H), 7.04 (dd, 1H, J = 5.7, 8.1 Hz), 6.80 (td, 1H, J = 2.9, 8.1 Hz), 5.52 (b, 1H), 5.41 (b, 1H), 3.01 (m, 2H), 2.89-2.74 (m, 6H), 2.07 (m, 2H), 1.39-1.12 (m, 6H); 510 (ESMS, MH⁺).

15

Example 5

1-Aza-9-fluoro-4,5-dihydro-2-(4-(methane-sulfonylamino)-butyl)amino-3-thia-benzo[e]azulene was obtained as a yellow oil in 59% yield: ¹H NMR δ 7.63 (dd, 1H, J = 2.7, 10.8 Hz), 7.07 (dd, 1H, J = 6.0, 8.4 Hz), 6.84 (td, 1H, J = 2.9, 8.4 Hz), 5.61 (b, 1H), 5.34 (b, 1H), 3.22 (m, 2H), 3.09 (m, 2H), 2.94 (s, 3H), 2.86-2.76 (m, 4H), 2.09 (m, 2H), 1.60 (m, 6H); 384 (ESMS, MH⁺).

25

Example 6

1-Aza-9-fluoro-4,5-dihydro-2-(4-(dimethyl-aminosulfonyl-amino)butyl)amino-3-thia-benzo[e]-azulene was obtained as a yellow oil in 75% yield: ¹H NMR δ 7.65 (dd, 1H, J = 2.4, 10.8 Hz), 7.06 (t, 1H, J = 7.1 Hz), 6.83 (td, 1H, J = 2.7, 8.1 Hz), 5.57 (b, 1H), 5.05 (b, 1H), 3.21 (m, 2H), 3.02 (q, 2H, J = 5.7 Hz), 2.87-2.76 (m, 10H), 2.08 (m, 2H), 1.59 (m, 4H); 413 (ESMS, MH⁺).

30

Example 7

1-Aza-9-fluoro-2-(4-(2-fluorophenyl)sulfonylamino)-
butylamino-4,5-dihydro-3-thia-benzo[e]azulene was obtained
5 as a light yellow oil in 81% yield: ^1H NMR δ 7.88 (td, 1H, J = 1.5, 9.0 Hz), 7.57 (m, 2H), 7.28-7.17 (m, 2H), 7.05 (dd, 1H, J = 6.0, 8.1 Hz), 6.81 (td, 1H, J = 2.7, 8.1 Hz), 5.64 (b, 2H), 3.14 (t, 2H, J = 6.3 Hz), 2.95 (t, 2H, J = 6.3 Hz), 2.83 (t, 2H, J = 6.9 Hz), 2.76 (m, 2H), 2.06 (m, 2H), 1.52 (m, 4H); 464 (ESMS, MH^+).

Example 8

1-Aza-9-fluoro-4,5-dihydro-2-(4-(1-naphthyl)sulfonyl-
15 amino)butylamino-3-thia-benzo[e]azulene was obtained as a yellow oil in 78% yield: ^1H NMR δ 8.68 (d, 1H, J = 8.4 Hz), 8.24 (d, 1H, J = 7.2 Hz), 8.05 (d, 1H, J = 8.4 Hz), 7.93 (d, 1H, J = 7.8 Hz), 7.66-7.50 (m, 4H), 7.05 (dd, 1H, J = 6.0, 8.4 Hz), 6.80 (td, 1H, J = 2.7, 8.1 Hz), 5.62 (b, 1H), 5.39 (b, 1H), 3.02 (m, 2H), 2.87-2.74 (m, 6H), 2.07 (m, 2H), 1.41 (m, 4H); 496 (ESMS, MH^+).

Example 9

1-Aza-9-fluoro-4,5-dihydro-2-(4-((2(S)-methoxy-methyl)-
pyrrolidine-1-yl)sulfonyl)phenylamino-3-thia-
benzo[e]azulene was obtained as a light yellow solid in
72% yield: m.p. 186-188 °C ^1H NMR δ 7.94 (b, 1H), 7.76 (m, 3H), 7.57 (d, 2H, J = 8.7 Hz), 7.11 (dd, 1H, J = 6.0, 8.4 Hz), 3.75-3.64 (m, 2H), 3.45-3.32 (m, 2H), 3.38 (s, 3H), 3.11 (m, 1H), 2.94 (t, 2H, J = 6.9 Hz), 2.83 (m, 2H), 2.15 (m, 2H), 1.84 (m, 2H), 1.60 (m, 2H); 488 (ESMS, MH^+).
The hydrogen chloride salt was obtained as a yellow solid:
m.p. 151-155 °C.

Example 10

5

1-Aza-9-fluoro-4,5-dihydro-2-(5-(methylsulfonyl-amino)-pentyl)amino-3-thia-benzo[e]azulene (hydrogen chloride salt) was obtained as a yellow oil in 53% yield: ^1H NMR (CD_3OD) δ 7.35 (dd, 1H, $J = 2.5, 9.9$ Hz), 7.28 (dd, 1H, $J = 6.0, 8.4$ Hz), 7.05 (td, 1H, $J = 2.5, 8.4$ Hz), 3.42 (t, 2H, $J = 7.0$ Hz), 3.06 (t, 2H, $J = 6.6$ Hz), 2.90 (s, 3H), 2.78 (t, 4H, $J = 7.1$ Hz), 2.14 (m, 2H), 1.74 (p, 2H, $J = 7.2$ Hz), 1.63-1.47 (m, 4H); 398 (ESMS, MH^+).

10

15

Example 11

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-(methylsulfonylamino-methyl)cyclohexyl)amino-3-thia-benzo[e]azulene: Anal. Calcd for $\text{C}_{20}\text{H}_{26}\text{FN}_3\text{O}_2\text{S}_2 + 0.25\text{H}_2\text{O}$: C, 56.12; H, 6.24; N, 9.82. Found: C, 56.15; H, 6.40; N, 9.52.

20

Example 12

25

1-Aza-9-fluoro-4,5-dihydro-2-(5-(2,4-difluorophenyl)sulfonylamino)pentylamino-3-thia-benzo[e]-azulene (hydrogen chloride salt) was obtained as a yellow syrup in 83% yield: ^1H NMR (CD_3OD) δ 7.91 (m, 1H), 7.64 (dd, 1H, $J = 2.8, 10.8$ Hz), 7.08-6.92 (m, 3H), 6.82 (td, 1H, $J = 2.8, 8.3$ Hz), 5.23 (b, 2H), 3.21 (m, 2H), 2.97 (m, 2H), 2.86 (t, 2H, $J = 6.9$ Hz), 2.78 (m, 2H), 2.07 (m, 2H), 1.63-1.34 (m, 6H); 496 (ESMS, MH^+).

30

Example 13

1-Aza-9-fluoro-4,5-dihydro-2-(5-isopropyl-sulfonylamino)-
pentylamino-3-thia-benzo[e]azulene (hydrogen chloride
5 salt) was obtained as a yellow syrup in 12% yield: ^1H NMR
(free base) δ 7.69 (dd, 1H, $J = 3.0, 10.8$ Hz), 7.05 (dd,
1H, $J = 5.9, 8.1$ Hz), 6.82 (td, 1H, $J = 2.8, 8.1$ Hz), 5.34
(b, 1H), 3.92 (t, 1H, $J = 6.0$ Hz), 3.29-3.11 (m, 3H), 2.85
(t, 2H, $J = 6.9$ Hz), 2.79 (m, 2H), 2.10 (m, 2H), 1.80 (s,
10 3H), 1.73 (s, 3H), 1.64 (m, 4H), 1.44 (m, 2H).

Example 14

1-Aza-9-fluoro-4,5-dihydro-2-(5-(diethyl-aminosulfonyl-
15 amino)pentyl)amino-3-thia-benzo[e]-azulene (hydrogen
chloride salt) was obtained as a yellow syrup in 12%
yield: ^1H NMR (free base) δ 7.67 (dd, 1H, $J = 2.7, 10.8$
Hz), 7.06 (dd, 1H, $J = 6.0, 8.3$ Hz), 6.83 (td, 1H, $J =$
2.7, 8.3 Hz), 5.39 (b, 1H), 4.57 (t, 1H, $J = 6.1$ Hz), 3.24
20 (m, 6H), 2.95 (q, 2H, $J = 6.7$ Hz), 2.85 (t, 2H, $J = 6.9$
Hz), 2.79 (m, 2H), 2.08 (m, 2H), 1.60 (m, 2H), 1.50 (m,
2H), 1.39 (m, 2H), 1.08 (t, 6H, $J = 7.1$ Hz); 455 (ESMS,
MH $^+$).

25

Example 15

1-Aza-9-fluoro-4,5-dihydro-2-(5-(2-methoxy-5-methyl-
phenyl)sulfonylamino)pentylamino-3-thia-benzo[e]-azulene
(hydrogen chloride salt) was obtained as a yellow syrup in
30 80% yield: ^1H NMR (free base) δ 7.70 (m, 1H), 7.66 (dd, 1H,
 $J = 2.9, 10.8$ Hz), 7.31 (dd, 1H, $J = 1.8, 8.4$ Hz), 7.05
(dd, 1H, $J = 6.0, 8.1$), 6.91 (d, 1H, $J = 8.4$ Hz), 6.82
(td, 1H, $J = 2.8, 8.3$ Hz), 5.26 (m, 1H), 5.10 (t, 1H, $J =$
6.1 Hz), 3.93 (s, 3H), 3.18 (q, 2H, $J = 5.7$ Hz), 2.85 (m,

103

4H), 2.78 (m, 2H), 2.33 (s, 3H), 2.09 (m, 2H), 1.58 (m, 2H), 1.48 (m, 2H), 1.34 (m, 2H); 504 (ESMS, MH⁺).

Example 16

5

1-Aza-2-(5-benzylsulfonylamino)pentylamino-9-fluoro-4,5-dihydro-3-thia-benzo[e]azulene was obtained as a yellow oil in 68% yield: ¹H NMR δ 7.66 (dd, 1H, J = 2.7, 10.8 Hz), 7.38 (m 5H), 7.06 (dd, 1H, J = 6.0, 8.1 Hz), 6.84 (td, 1H, J = 2.9, 8.3 Hz), 5.21 (m, 1H), 4.62 (m, 1H), 4.24 (s, 2H), 3.21 (q, 2H, J = 6.2 Hz), 2.93 (q, 2H, J = 6.4 Hz), 2.85 (t, 2H, J = 6.9 Hz), 2.78 (m, 2H), 2.07 (m, 2H), 1.60 (m, 2H), 1.45 (m, 2H), 1.35 (m, 2H); 474 (ESMS, MH⁺).

10

15

Example 17

1-Aza-2-(5-(3,4-difluorophenyl)sulfonylamino)-pentylamino-9-fluoro-4,5-dihydro-3-thia-benzo[e]-azulene was obtained as a yellow oil in 78% yield: ¹H NMR δ 7.73-7.61 (m, 2H), 7.56 (dd, 1H, J = 2.9, 10.8 Hz), 7.34-7.25 (m, 1H), 7.06 (dd, 1H, J = 2.9, 8.2 Hz), 6.82 (td, 1H, J = 2.8, 8.2 Hz), 5.42 (b, 1H), 5.34 (b, 1H), 3.19 (m, 2H), 2.92 (m, 2H), 2.86 (t, 2H, J = 7.0 Hz), 2.78 (m, 2H), 2.07 (m, 2H), 1.58 (m, 2H), 1.43 (m, 2H), 1.36 (m, 2H); 496 (ESMS, MH⁺). The hydrogen chloride salt was obtained as a tan solid: m.p. 68-70 °C.

20

25

Example 18

30

1-Aza-9-fluoro-4,5-dihydro-2-(5-(4-methoxyphenyl)sulfonylamino)pentylamino-3-thia-benzo[e]azulene was obtained as a yellow oil in 73% yield: ¹H NMR δ 7.79 (d, 2H, J = 9.0 Hz), 7.62 (dd, 1H, J = 2.9, 10.8 Hz), 7.06 (dd, 1H, J = 6.0, 8.4 Hz), 6.97 (d, 2H, J = 8.7 Hz), 6.82

104

(td, 1H, $J = 2.8, 8.3$ Hz), 5.22 (m, 1H), 4.92 (m, 1H), 3.85 (s, 3H), 3.19 (q, 2H, $J = 5.7$ Hz), 2.91 (m, 2H), 2.86 (t, 2H, $J = 6.9$ Hz), 2.78 (m, 2H), 2.07 (m, 2H), 1.57 (p, 2H, $J = 7.2$ Hz), 1.45 (m, 2H), 1.36 (m, 2H); 490 (ESMS, MH⁺).

Example 19

1-Aza-9-fluoro-4,5-dihydro-2-(5-(2-thienyl)-sulfonylamino)-pentylamino-3-thia-benzo[e]azulene was obtained as a light yellow solid in 70% yield: m.p. 152-154 °C; ¹H NMR δ 7.64-7.56 (m, 3H), 7.08 (m, 2H), 6.82 (td, 1H, $J = 2.8, 8.2$ Hz), 5.25 (b, 1H), 5.16 (b, 1H), 3.20 (m, 2H), 3.01 (m, 2H), 2.86 (t, 2H, $J = 6.9$ Hz), 2.77 (m, 2H), 2.07 (m, 2H), 1.58 (p, 2H, $J = 7.2$ Hz), 1.47 (m, 2H), 1.36 (m, 2H); 466 (ESMS, MH⁺).

Example 20

1-Aza-9-fluoro-2-(5-(2-trifluoroethyl)sulfonylamino)-pentylamino-4,5-dihydro-3-thia-benzo[e]azulene was obtained as a yellow oil in 25% yield: ¹H NMR δ 7.65 (dd, 1H, $J = 2.9, 10.8$ Hz), 7.07 (dd, 1H, $J = 5.9, 8.4$ Hz), 6.84 (td, 1H, $J = 2.9, 8.3$ Hz), 5.30 (b, 2H), 3.77 (q, 2H, $J = 9.1$ Hz), 3.25 (t, 2H, $J = 6.6$ Hz), 3.14 (t, 2H, $J = 6.6$ Hz), 2.85 (t, 2H, $J = 6.9$ Hz), 2.77 (m, 2H), 2.08 (m, 2H), 1.72-1.52 (m, 4H), 1.42 (m, 2H); 466 (ESMS, MH⁺).

Example 21

1-Aza-9-fluoro-2-(5-ethylsulfonylamino)pentylamino-4,5-dihydro-3-thia-benzo[e]azulene was obtained as a yellow oil in 72% yield: ¹H NMR δ 7.65 (dd, 1H, $J = 2.7, 10.8$ Hz), 7.06 (dd, 1H, $J = 6.0, 8.2$ Hz), 6.83 (td, 1H, $J = 3.0, 8.2$

105

Hz), 5.52 (m, 1H), 5.07 (m, 1H), 3.22 (m, 2H), 3.05 (m, 4H), 2.85 (t, 2H, $J = 7.0$ Hz), 2.77 (m, 2H), 2.05 (m, 2H), 1.60 (m, 2H), 1.51 (m, 2H), 1.38 (m, 5H); 412 (ESMS, MH^+).

5

Example 22

1-Aza-2-(4-diethylaminosulfonylamino)butylamino-9-fluoro-4,5-dihydro-3-thia-benzo[e]azulene was obtained as a yellow oil in 36% yield: 1H NMR δ 7.68 (dd, 1H, $J = 2.7$, 10.8 Hz), 7.06 (dd, 1H, $J = 6.0$, 8.3 Hz), 6.84 (td, 1H, $J = 2.4$, 8.2 Hz), 5.18 (m, 1H), 4.36 (t, 1H, $J = 6.3$ Hz), 3.26 (m, 6H), 3.01 (q, 2H, $J = 6.5$ Hz), 2.86 (t, 2H, $J = 6.9$ Hz), 2.08 (m, 2H), 1.67 (m, 4H), 1.18 (t, 6H, $J = 7.1$ Hz).

15

Example 23

1-Aza-9-fluoro-4,5-dihydro-2-(5-(1-methylimidazol-4-yl)sulfonylamino)pentylamino-3-thia-benzo[e]azulene was obtained as a yellow solid in 74% yield: m.p. 50-53 °C; 1H NMR δ 7.66 (dd, 1H, $J = 2.7$, 10.8 Hz), 7.53 (s, 1H), 7.46 (s, 1H), 7.05 (dd, 1H, $J = 6.0$, 8.2 Hz), 6.82 (td, 1H, $J = 2.7$, 8.2 Hz), 5.96 (m, 1H), 5.47 (m, 1H), 3.71 (s, 3H), 3.18 (m, 2H), 2.96 (m, 2H), 2.85 (t, 2H, $J = 6.9$ Hz), 2.06 (m, 2H), 1.56 (m, 2H), 1.48 (m, 2H), 1.37 (m, 2H).

25

Example 24

1-Aza-9-fluoro-4,5-dihydro-2-(5-(3,5-dimethyl-isoxazol-4-yl)sulfonylamino)pentylamino-3-thia-benzo[e]azulene was obtained as a yellow oil in 80% yield: 1H NMR δ 7.63 (dd, 1H, $J = 2.7$, 10.8 Hz), 7.06 (dd, 6.0, 8.3 Hz), 6.83 (dd, 1H, $J = 2.8$, 8.2 Hz), 5.24 (b, 2H), 3.48 (m, 2H), 2.94 (m, 2H), 2.85 (t, 2H, $J = 7.0$ Hz), 2.77 (m, 2H), 2.63 (s, 3H),

30

2.40 (s, 3H), 2.08 (m, 2H), ¹⁰⁶ 1.60 (p, 2H, $J = 7.5$ Hz),
1.49 (m, 2H), 1.38 (m, 2H).

Example 25

5

1-Aza-9-fluoro-4,5-dihydro-2-(5-aminosulfonylamino)-
pentylamino-3-thia-benzo[e]azulene was obtained as a
yellow oil in 39% yield: ¹H NMR δ 7.58 (dd, 1H, $J = 2.7$,
10.8 Hz), 7.06 (dd, 1H, $J = 6.0$, 8.3 Hz), 6.82 (td, 1H, J
10 = 2.7, 8.2 Hz), 5.30 (b, 2H), 5.18 (b, 2H), 3.19 (t, 2H, J
= 6.4 Hz), 3.09 (m, 2H), 2.82 (t, 2H, $J = 5.4$ Hz), 2.76
(m, 2H), 2.07 (m, 2H), 1.85-1.47 (m, 4H), 1.43 (m, 2H).

Example 26

15

trans-1-aza-9-fluoro-2-(4-(2-fluorophenyl)sulfonyl-amino-
methyl)cyclohexylamino-4,5-dihydro-3-thia-benzo[e]-
azulene: ¹H NMR δ 7.89 (apparent td, 1H, $J = 1.4$, 7.5 Hz),
7.62 (dd, 1H, $J = 2.7$, 10.8 Hz), 7.59 (m, 1H), 7.25-7.19
20 (m, 2H), 7.04 (dd, 1H, $J = 6.0$, 8.1 Hz), 6.81 (td, 1H, J
= 3.0, 8.2 Hz), 5.08 (broad m, 1H), 4.95 (m, 1H), 3.02 (m,
1H), 2.90-2.75 (m, 6H), 2.25-0.80 (m, 10H).

Example 27

25

trans-1-Aza-9-fluoro-4,5-dihydro-2-{4-(4-methoxy-phenyl)-
sulfonylaminomethyl}cyclohexylamino-3-thia-
benzo[e]azulene: 6.2% yield; ¹H NMR (CDCl₃) δ 7.79 (d, 2H,
 $J=8.7$ Hz), 7.62 (dd, 1H, $J=10.8$, 2.7 Hz), 7.05 (dd, 1H,
 $J=6.0$, 8.3 Hz), 6.98 (d, 2H, $J=8.7$ Hz), 6.82 (apparent dt,
30 1H, $J=2.8$, 8.2 Hz), 4.95 (d, 1H, $J=8$ Hz), 4.66 (t, 1H,
 $J=5.1$ Hz), 3.85 (s, 3H), 3.24 (b, 1H), 2.90-0.80 (m, 17H).

Example 28

trans-N2-(4-(2,6-Difluorophenylsulfonyl)aminomethyl)-
cyclohexyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]-cyclohepta-
5 [d][1,3]-thiazol-2-amine: A solution of *trans*-N2-(4-
aminomethyl)cyclohexyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]-
cyclohepta[d][1,3]thiazol-2-amine hydrochloride (200 mg,
0.52 mmol) and *N,N*-diisopropylethylamine (0.22 ml, 1.29
mmol) in dry THF (8 ml) was treated with 2,6-
10 difluorobenzenesulfonyl chloride (117 mg, 0.55 mmol) in
THF (2 ml). The mixture was stirred at room temperature
overnight. The solvent was evaporated. The residue was
dissolved in dichloromethane. The solution was then washed
with 5% HCl solution, dried (MgSO₄), filtered and
15 concentrated. The product was purified by preparative TLC
(EtOAc/hexane 1:1) to give a tan solid (53% yield): ¹H NMR
(CDCl₃) δ 1.10 (4H, m), 1.42 (1H, m), 1.80 (2H, m), 2.10
(2H, m), 2.20 (2H, m), 2.80 (2H, m), 2.84 (2H, t, *J* = 6.9
Hz), 2.92 (2H, m), 3.23 (1H, m), 5.05 (1H, broad), 5.55
20 (1H, broad), 6.80 (1H, m), 7.05 (3H, m), 7.50 (1H, m),
7.65 (1H, m)

Similarly prepared were:

25

Example 29

trans-1-Aza-2-{4-benzylsulfonylaminomethyl}-
cyclohexylamino-9-fluoro-4,5-dihydro-3-thia-benzo[e]-
30 azulene: 23.6% yield; ¹H NMR (CDCl₃) δ 7.62 (dd, 1H,
J=10.8, 2.7 Hz), 7.50-7.20 (m, 5H), 7.05 (dd, 1H, *J*=6.0,
8.3 Hz), 6.81 (apparent dt, 1H, *J*=2.8, 8.2 Hz), 4.95 (m,
1H), 4.30 (m, 1H), 4.24 (m, 2H), 4.10 (m, 2H), 3.25
(broad, 1H), 2.90-0.80 (m, 15H).

Example 30

trans-N2-(4-(2-Thienylsulfonyl)aminomethyl)-cyclohexyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]-

5 cyclohepta[d][1,3]thiazol-2-amine: ¹H NMR (CDCl₃) δ 1.15 (4H, m), 1.45 (1H, m), 1.80 (2H, m), 2.07 (2H, m), 2.20 (2H, m), 2.82 (6H, m), 3.25 (1H, m), 5.00 (2H, m), 6.82 (1H, m), 7.05 (1H, m), 7.10 (1H, t, *J* = 4.4 Hz), 7.62 (3H, m).

10

Example 31

trans-N2-(4-Ethylsulfonylaminomethyl)cyclohexyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-

15 amine: ¹H NMR (CDCl₃) δ 1.15 (4H, m), 1.38 (3H, t, *J* = 7.4 Hz), 1.45 (1H, m), 1.90 (2H, m), 2.10 (2H, m), 2.25 (2H, m), 2.80 (2H, m), 2.86 (2H, t, *J* = 6.9 Hz), 2.97 (2H, t, *J* = 6.6 Hz), 3.05 (2H, q, *J* = 7.4 Hz), 3.30 (1H, broad), 4.50 (1H, m), 4.97 (1H, m), 6.83 (1H, m), 7.07 (1H, m),
20 7.66 (1H, m).

Example 32

trans-1-Aza-9-fluoro-4,5-dihydro-2-{4-(1-methylimidazolyl-4-yl)sulfonylaminomethyl}cyclohexyl-amino-3-thia-

25 benzo[e]azulene in 33% yield: ¹H NMR (CDCl₃) δ 7.67 (dd, 1H, *J*=10.8, 3.0 Hz), 7.52 (s, 1H), 7.48 (s, 1H), 7.05 (dd, 1H, *J*=6.0, 8.2 Hz), 6.82 (apparent dt, 1H, *J*=2.8, 8.3 Hz), 5.43 (broad, 1H), 5.01 (d, 1H, *J*=8.0 Hz), 3.76 (s, 3H),
30 3.22 (m, 1H), 2.90-2.70 (m, 5H), 2.30-0.80 (m, 9H).

Example 33

trans-1-Aza-9-fluoro-4,5-dihydro-2-{4-(3,5-dimethylisoxazol-4-yl)sulfonylaminomethyl}cyclohexyl-amino-3-thia-benzo[e]azulene in 87% yield: ^1H NMR (CDCl_3) δ 7.55 (dd, 1H, $J=10.8$, 3.0 Hz), 7.05 (dd, 1H, $J=6.0$, 8.2 Hz), 6.81 (apparent dt, 1H, $J=2.8$, 1.3 Hz), 5.95 (broad, 1H), 5.30 (broad, 1H), 3.20-2.70 (m, 5H), 2.61 (s, 3H), 2.38 (s, 3H), 2.20-0.80 (m, 9H).

Example 34

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-methylsulfonylamino)-cyclohexylamino-3-thia-benzo[e]azulene was obtained as a white solid in 41% yield: m.p. 179-181 °C; ^1H NMR δ 7.67 (dd, 1H, $J = 2.8$, 10.8 Hz), 7.07 (dd, 1H, $J = 6.0$, 8.2 Hz), 6.84 (td, 1H, $J = 2.8$, 8.2 Hz), 4.96 (b, 1H), 4.52 (b, 1H), 3.32 (b, 2H), 3.00 (s, 3H), 2.86 (t, 2H, $J = 6.9$ Hz), 2.79 (m, 2H), 2.26 (m, 2H), 2.17 (m, 4H), 1.35 (m, 4H); 410 (ESMS, MH^+).

Example 35

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-diethylamino-sulfonylamino)cyclohexylamino-3-thia-benzo[e]azulene was obtained as a tan solid in 13% yield: m.p. 172-174 °C; ^1H NMR δ 7.68 (dd, 1H, $J = 2.8$, 10.8 Hz), 7.06 (dd, 1H, $J = 5.8$, 8.3 Hz), 6.84 (td, 1H, $J = 2.7$, 8.2 Hz), 4.82 (m, 1H), 3.96 (d, 1H, $J = 7.5$ Hz), 3.38 (m, 1H), 3.27 (q, 4H, $J = 7.1$ Hz), 3.17 (m, 1H), 2.86 (t, 2H, $J = 6.9$ Hz), 2.80 (m, 2H), 2.27-2.03 (m, 6H), 1.32 (m, 2H), 1.20 (t, 6H, $J = 7.1$ Hz); 467 (ESMS, MH^+).

Example 36

110

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-(4-methoxy-phenyl)sulfonylamino)cyclohexylamino-3-thia-benzo[e]-azulene was obtained as a yellow syrup in 50% yield: ^1H NMR δ 7.82 (d, 2H, J = 8.9 Hz), 7.65 (dd, 1H, J = 2.8, 10.8 Hz), 7.05 (dd, 1H, J = 6.0, 8.2 Hz), 6.98 (d, 2H, J = 8.9 Hz), 6.82 (td, 1H, J = 2.9, 8.3 Hz), 4.83 (d, 1H, J = 7.7 Hz), 4.55 (d, 1H, J = 7.5 Hz), 3.88 (s, 3H), 3.27 (m, 1H), 3.10 (m, 1H), 2.84 (t, 2H, J = 6.9 Hz), 2.78 (m, 2H), 2.16-2.02 (m, 4H), 1.89 (m, 2H), 1.24 (m, 4H); 502 (ESMS, MH^+).

Example 37

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-(2-thienyl)sulfonyl-amino)cyclohexylamino-3-thia-benzo[e]azulene was obtained as a tan solid in 36% yield: m.p. 230-231 $^{\circ}\text{C}$; ^1H NMR δ 7.67-7.58 (m, 3H), 7.10-7.03 (m, 2H), 6.81 (td, 1H, J = 2.8, 8.3 Hz), 4.87 (m, 1H), 4.79 (m, 1H), 3.26 (m, 2H), 2.84 (t, 2H, J = 6.9 Hz), 2.77 (m, 2H), 2.18-2.02 (m, 4H), 1.93 (m, 2H), 1.28 (m, 4H); 478 (ESMS, MH^+).

Example 38

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-(2,2,2-trifluoroethyl)sulfonylamino)cyclohexylamino-3-thia-benzo[e]azulene was obtained as a yellow solid in 32% yield: m.p. 89-91 $^{\circ}\text{C}$; ^1H NMR δ 7.68 (dd, 1H, J = 2.7, 10.8 Hz), 7.07 (dd, 1H, J = 6.0, 8.2 Hz), 6.84 (td, 1H, J = 2.8, 8.3 Hz); 4.78 (m, 1H), 4.72 (m, 1H), 3.82 (q, 2H, J = 8.9 Hz), 3.38 (m, 2H), 2.89-2.81 (m, 4H), 2.28 (m, 2H), 2.10 (m, 4H), 1.47-1.30 (m, 4H); 478 (ESMS, MH^+).

Example 39

1-Aza-9-fluoro-4,5-dihydro-2-(4-(2,2,2-trifluoroethyl)-sulfonylamino)butylamino-3-thia-benzo[e]azulene (TFA salt) was obtained as a white solid in 35% yield: m.p. 140-141 °C; 452 (ESMS, MH⁺).

10

Example 40

trans-1-Aza-9-fluoro-2-{4-(3,4-difluorophenyl)-sulfonylaminomethyl}cyclohexylamino-4,5-dihydro-3-thia-benzo[e]azulene: 522 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 7.74-7.63 (m, 2H), 7.60 (dd, 1H, J=2.7, 10.8 Hz), 7.32 (apparent dq, 1H, J=7.6, 1.0), 7.06 (dd, 1H, J=6.0, 8.3 Hz), 6.82 (apparent dt, 1H, J=2.8, 8.2 Hz), 5.05-4.95 (m, 2H), 3.25 (m, 1H), 2.95-2.60 (m, 6H), 2.05 (m, 2H), 2.01 (ABm, 4H), 1.62 (broad, 1H), 1.41 (m, 1H), 1.30-0.80 (m, 4H).

20

Example 41

trans-1-Aza-9-fluoro-2-{4-trifluoromethylsulfonylaminomethyl}cyclohexylamino-4,5-dihydro-3-thiabenz[e]azulene in 17% yield: 446 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 7.69 (dd, 1H, J=2.7, 10.8 Hz), 7.05 (dd, 1H, J=6.0, 8.3 Hz), 6.82 (apparent dt, 1H, J=2.8, 8.2 Hz), 4.87 (d, 1H, J=8.0 Hz), 4.61 (m, 1H), 3.29 (m, 1H), 3.00 (t, 2H, J=9.2 Hz), 2.86 (t, 2H, J=6.9 Hz), 2.83-2.77 (m, 2H), 2.06 (m, 2H), 2.01 (ABm, 4H), 1.80-1.00 (m, 6H).

30

Example 42

trans-1-Aza-9-fluoro-2-{4-(2-fluoro)phenylsulfonyl-amino}-
cyclohexylmethylamino-4,5-dihydro-3-thia-benzo[e]azulene
in 100% yield: 504 (ESMS, MH^+); 1H NMR ($CDCl_3$) δ 7.90
5 (apparent dt, 1H, $J=1.1$ average, 7.7 Hz), 7.63 (dd, 1H,
 $J=2.8$, 10.7 Hz), 7.57 (apparent dq, 1H, $J=1.1$, 3.5 Hz),
7.56-7.17 (m, 2H), 7.04 (dd, 1H, $J=6.0$, 8.3 Hz), 6.82
(apparent dt, 1H, $J=2.8$, 8.3 Hz), 5.46 (broad, 1H), 4.94
(d, 1H, $J=5.7$ Hz), 3.15 (m, 1H), 3.05 (m, 2H), 2.83 (t,
10 2H, $J=6.9$ Hz), 2.05-0.75 (m, 12H).

Example 43

trans-N2-(4-Methylsulfonylamino)cyclohexylmethyl-9-fluoro-
15 5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-
amine: A mixture of trans-N2-(4-amino)cyclohexylmethyl-9-
fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-
2-amine dihydrochloride (418 mg, 1.0 mmol) and
triethylamine (1.25 mL) in THF (6 ml) was cooled in an ice
20 water bath and treated with methanesulfonyl chloride
dropwise (0.10 ml). The mixture was stirred at room
temperature for 30 minutes and then poured into water (60
ml). The product was extracted into chloroform and
purified by flash chromatography over silica gel
25 (EtOAc/hexane 1:1) to afford a yellow solid (227 mg, 54%
yield): CIMS m/e = 424 (MH^+); 1H NMR (300 MHz, $CDCl_3$) δ
1.0-1.3 (4H, m), 1.4-1.6 (1H, m), 1.84 (2H, d, J = 12.3
Hz), 2.0-2.2 (4H, m), 2.77 (2H, t, J = 5.4 Hz), 2.84 (2H,
t, J = 6.9 Hz), 2.97 (3H, s), 3.07 (2H, t, J = 6.1 Hz),
30 3.1-3.30 (1H, m), 4.66 (1H, d, J = 7.6 Hz), 5.52 (1H, t, J
= 4.5 Hz), 6.83 (1H, dt, J = 2.8, 8.2 Hz), 7.06 (1H, dd, J
= 5.9, 8.2 Hz), 7.65 (1H, dd, J = 2.7, 10.7 Hz).

Example 44

5 *trans*-N2-(4-Aminosulfonylamino)cyclohexylmethyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine

a. *trans*-1-Aminomethyl-4-(benzyloxycarbonylamino)-cyclohexane: To a solution of *trans*-1-(*t*-butyloxycarbonylaminomethyl)-4-(benzyloxycarbonyl-amino)-cyclohexane (20 g, 55.2 mmol) in EtOAc (790 mL) was added 4N HCl in dioxane (200 mL). The mixture was stirred overnight and then concentrated to give a white solid: ¹H NMR (300 MHz, CD₃OD) δ 1.0-1.3 (4H, m), 1.50-1.67 (1H, m), 1.83 (2H, d, *J* = 12.3 Hz), 1.96 (2H, d, *J* = 10.3 Hz), 2.77 (2H, d, *J* = 7.0 Hz), 3.20-3.40 (1H, m), 5.03 (1H, s), 7.20-7.40 (5H, m).

b. *trans*-1-Aminocarbothioylaminomethyl-4-(benzyloxy-carbonylamino)cyclohexane: *trans*-1-Aminomethyl-4-(benzyloxycarbonylamino)cyclohexane (55.2 mmol) was dissolved in a mixture of chloroform (350 ml) and triethylamine (16 mL), cooled in an ice water bath and treated with benzoylisothiocyanate (10 ml, 74.4 mmol). The mixture was stirred for 6 hours before it was allowed to slowly warm to room temperature overnight. The solvent was evaporated and the residue triturated with MeOH (800 ml) and potassium carbonate (70 g) solution in water (800 ml). The mixture was heated at reflux for 10 min. and then additional aqueous potassium carbonate solution (34 g/140 ml) was added. The mixture was cooled to 8 °C. The resulting precipitate was collected by filtration and washed with water and ether (200 mL) to give the desired product as a white solid (12.02 g, 68%): m.p. 155-158 °C; ¹H NMR (300 MHz, CD₃OD) δ 0.9-1.3 (4H, m), 1.4-1.55 (1H,

115

solid: ^1H NMR of the free base (300 MHz, CDCl_3) δ 1.1-1.6 (4H, m), 1.3-1.7 (4H, m), 1.80-2.00 (4H, m), 2.00-2.20 (2H, m), 2.30 (1H, m), 2.79 (2H, t, $J = 5.5$ Hz), 2.86 (2H, 5, $J = 6.9$ Hz), 3.12 (2H, t, $J = 6.2$ Hz), 5.07 (1H, t), 6.82 (1H, dt, $J = 2.8$ Hz), 7.05 (1H, dd, $J = 5.9, 8.3$ Hz), 7.70 (1H, dd, $J = 2.8, 10.8$ Hz).

e. *trans*-N2-(4-Aminosulfonylamino)cyclohexylmethyl-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine: A mixture of *trans*-N2-(4-amino)cyclohexylmethyl-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine dihydrochloride (524 mg, 1.25 mmol), sulfamide (371 mg) and triethylamine (1 mL) in THF (7 ml), methanol (5 mL) and water (7 ml) was heated at reflux for 18 hours. Most of the solvent was evaporated and then water (50 ml) was added. The product was extracted into chloroform and then purified by flash chromatography over silica gel (EtOAc/hexane 5:4) to give the desired product as a yellow solid (238 mg, 56%): m.p. 180-182 °C; CIMS $m/e = 425$ (MH^+); ^1H NMR (300 MHz, CDCl_3) δ 1.00-1.30 (4H, m), 1.40-1.60 (1H, m), 1.81 (2H, d, $J = 12.0$ Hz), 2.00-2.20 (4H, m), 2.75 (2H, t, $J = 5.3$ Hz), 2.82 (2H, t, $J = 6.9$ Hz), 3.02 (2H, d, $J = 6.4$ Hz), 3.20 (1H, m), 4.85 (1H, d, $J = 6.7$ Hz), 5.20 (2H, s), 5.85 (1H, s), 6.83 (1H, dt, $J = 2.8, 8.2$ Hz), 7.05 (1H, dd, $J = 5.9, 8.3$ Hz), 7.58 (1H, dd, $J = 2.8, 10.7$ Hz).

Example 45

9-Fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine: 6-Bromo-3-fluoro-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one (1.0 g, 3.89 mmole) was mixed with thiourea (355 mg, 4.66 mmole) in 20 ml of ethanol and heated at reflux temperature for 16 hours. The reaction

114

m), 1.79 (2H, d, $J = 9.5$ Hz), 1.92 (2H, d, $J = 10.0$ Hz), 2.95 (1H, d, $J = 6.0$ Hz), 3.20-3.40 (2H, m), 5.03 (2H, s), 7.20-7.40 (5H, m).

5 c. *trans*-N2-(4-Benzyloxycarbonylamino)cyclohexyl-methyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]-cyclohepta[d][1,3]thiazol-2-amine: A mixture of *trans*-1-aminocarbothioylaminomethyl-4-(benzyloxy-carbonylamino)cyclohexane (4.46 g, 13.9 mmol), 2-bromo-8-fluoro-1-benzosuberone (4.08 g, 15.9 mmol), and *N,N*-diisopropylethylamine (4.6 mL) in ethanol (95 ml) was heated at reflux for 6 hours. The solvent was evaporated and the residue washed with water (200 ml). Ether (10 ml) was then added and the suspension filtered. The solid was washed with ether (10 ml) to give a white solid (5.55 g, 83%): CIMS $m/e = 480$ (MH^+); 1H NMR (300 MHz, $CDCl_3$) δ 1.0-1.2 (4H, m), 1.40-1.55 (1H, m), 1.89 (2H, d, $J = 7.7$ Hz), 2.00-2.16 (4H, m), 2.77 (2H, t, $J = 5.4$ Hz), 2.86 (2H, t, $J = 6.9$ Hz), 3.13 (2H, t, $J = 6.4$ Hz), 3.4-3.6 (1H, m), 4.30 (1H, d, $J = 6$ Hz), 5.05 (1H, m), 5.09 (2H, s), 6.8 (1H, dt, $J = 2.8, 8.4$ Hz), 7.06 (1H, dd, $J = 5.7, 8.4$ Hz), 7.35 (5H, m), 7.69 (1H, dd, $J = 2.8, 10.8$ Hz).

25 d. *trans*-N2-(4-Amino)cyclohexylmethyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine: *trans*-N2-(4-Benzyloxycarbonylamino)cyclohexyl-methyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine (5.55 g) was heated at reflux in a mixture of trifluoroacetic acid (50 ml) and water (5 ml) for 30 min. Most of the solvent was evaporated and the residue poured into water (150 ml). Potassium carbonate was added until the pH reached 10. The product was extracted into chloroform and converted to the hydrochloride salt which was isolated as an off-white

116

mixture was cooled and filtered to give a solid (m.p. 270 °C). The filtrate was concentrated and then partitioned between ethyl acetate and saturated sodium carbonate solution. The organic extract was concentrated and chromatographed (silica, ethyl acetate/hexanes) to give the desired product as a yellow solid (m.p. 142-144 °C, 34%).

N1-(9-Fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-[d][1,3]thiazol-2-yl)-5-bromopentanamide:

9-Fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine (150 mg, 0.640 mmole) was dissolved in 3 ml of dry THF, mixed with triethylamine (135 µl, 0.970 mmole), cooled in an ice bath and treated with 5-bromovaleryl chloride (90 µl, 0.67 mmole). The mixture was stirred at room temperature for 3 days. TLC analysis of the mixture showed 50% completion. Hence, a catalytic amount of DMAP was added and stirring was continued for 16 hours. There was no change in the TLC of the mixture. 5-Bromovaleryl chloride (90 µl) was added and stirred for 3 days. The reaction mixture was partitioned between ethyl acetate and saturated sodium carbonate and the aqueous phase separated and extracted with 2X3 ml ethyl acetate. The combined ethyl acetate extracts were washed with saturated sodium chloride solution, concentrated and chromatographed (silica, Ethyl acetate/hexanes) to give 179 mg of the desired product as a white foam (70%): 397, 399 (ESMS, MH+).

1-5-[(9-Fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-[d][1,3]-thiazol-2-yl)amino]-5-oxopentyl-1,2-triazadien-2-ium:

N1-(9-Fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-[d][1,3]thiazol-2-yl)-5-bromopentanamide (179 mg, 0.45

117

mmole) was dissolved in 1.5 ml of dry DMSO and treated with 72 mg of ammonium chloride (1.35 mmole) and 59 mg of sodium azide (0.91 mmole). After 5 hours, the mixture was heated at 60 °C for 16 hours. The reaction mixture was cooled, quenched with ice-water, extracted with ethyl acetate (3x3 ml), and the combined ethyl acetate extracts washed with water, and concentrated to give 145 mg of a yellow solid (90%): 360 (ESMS, MH⁺).

10 N1-(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-
[d][1,3]thiazol-2-yl)-5-aminopentanamide:

3907-50-2 1-5-[(9-Fluoro-5,6-dihydro-4H-benzo[6,7]-
cyclohepta[d][1,3]thiazol-2-yl)amino]-5-oxopentyl-1,2-
triazadien-2-ium (145 mg, 0.40 mmole) was dissolved in 2
15 ml of THF and treated with P(CH₃)₃ in THF (1M, 0.80 ml, 0.8
mmol) and water (22 µL, 1.22 mmole). The resulting
mixture was stirred at room temperature for 16 hours and
the solvent evaporated. The residue was dissolved in
ethyl acetate and chromatographed over silica gel (14 g)
20 eluting with ethyl acetate and then with EtOAc/7M NH₃ in
MeOH (20:3) to give an oil (115 mg).

N1-(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-
[d][1,3]thiazol-2-yl)-5-[(methylsulfonyl)amino]-
25 pentanamide:

N1-(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-
[d][1,3]thiazol-2-yl)-5-aminopentanamide (115 mg, 0.34
mmole) was dissolved in 2 ml of dry THF, cooled with an
ice bath, triethylamine was added (96 µl, 0.69 mmole)
30 followed by addition of 27 µl of methane sulfonyl chloride
(0.35 mmole). The reaction mixture was stirred at room
temperature for 6 hours, quenched with saturated sodium
carbonate solution and extracted with 3x3 ml of ethyl
acetate. The combined ethyl acetate extracts were washed

118

with saturated sodium chloride solution, concentrated and chromatographed (12 g of silica gel, EtOAc/hexanes/7N NH₃ in MeOH, 50:50:2) to give 53 mg of the desired product as a white solid (38%): m.p. 145-147 °C; 412 (ESMS, MH⁺).

Example 46

10 *trans*-N2-(4-Aminosulfonylaminomethyl)cyclohexyl-4,5-dihydro-benzo[2,3]oxepino[4,5-d][1,3]thiazol-2-amine

a. 2,3,4,5-Tetrahydro-1-benzoxepin-5-one To a solution of P₂O₅ (62 g) in methanesulfonic acid (415 ml) was added a solution of 4-phenoxybutyric acid (16 g) in methanesulfonic acid (120 ml) in 4 portions every 2 hours. After the last addition, the mixture was allowed to stand for another 2 hours before it was poured onto ice (3 Kg). The mixture was neutralized with 50% NaOH solution and the product extracted into chloroform. Purification by flash chromatography over silica gel (EtOAc/hexane 1:10) afforded an oil (5.06 g, 35%): ¹H NMR (300 MHz, CDCl₃) δ 2.22 (2H, m), 2.91 (2H, t, J = 7.0 Hz), 4.25 (2H, t, J = 6.6 Hz), 7.10 (2H, m), 7.43 (1H, dt, J = 1.8, 7.5 Hz), 7.77 (1H, dd, J = 1.6, 7.7 Hz).

25 b. 4-Bromo-2,3,4,5-tetrahydro-1-benzoxepin-5-one: To a solution of 2,3,4,5-tetrahydro-1-benzoxepin-5-one (5.06 g, 31.2 mmol) in acetic acid (50 ml) cooled by an ice water bath was added bromine (1.75 ml) dropwise. The solvent was evaporated to give the product as a brown oil: ¹H NMR (300 MHz, CDCl₃) δ 2.40-2.60 (1H, m), 2.90-3.00 (1H, m), 4.08-4.18 (1H, m), 4.42-4.49 (1H, m), 4.99 (1H, dd, J = 8.1, 7.0 Hz), 7.07 (1H, d, J = 8.3 Hz), 7.12 (1H, t, J =

119

7.5 Hz), 7.45 (1H, dt, $J = 1.6, 7.7$ Hz), 7.75 (1H, dd, $J = 1.7, 7.8$ Hz).

c *trans*-N2-(4-(*t*-Butyloxycarbonylaminomethyl)-cyclohexyl-4,5-dihydro-benzo[2,3]oxepino[4,5-*d*][1,3]thiazol-2-amine:

A mixture of 4-bromo-2,3,4,5-tetrahydro-1-benzoxepin-5-one (7.6 mmol), *trans*-1-(aminocarbothioylamino)-4-(*t*-butyloxycarbonylaminomethyl)cyclohexane (6.9 mmol) and *N,N*-diisopropylethylamine (2.5 mL) in ethanol (48 mL) was heated at reflux for 6 hours. The solvent was evaporated and the residue flash chromatographed over silica gel (EtOAc/hexane 1:3) to give a yellow solid (1.97 g, 66%): ^1H NMR (300 MHz, CDCl_3) δ 1.00-1.15 (4H, m), 1.44 (10H, s), 1.82 (2H, d, $J = 11.8$ Hz), 2.21 (2H, d, $J = 11.9$ Hz), 2.99 (2H, t, $J = 6.4$ Hz), 3.14 (2H, t, $J = 5.1$ Hz), 3.2-3.25 (1H, m), 4.32 (2H, t, $J = 5.2$ Hz), 4.64 (1H, t, $J = 4.7$ Hz), 4.94 (1H, d, $J = 8.0$ Hz), 6.98 (1H, dd, $J = 1.3, 7.8$ Hz), 7.07 (1H, dt, $J = 1.4, 7.5$ Hz), 7.14 (1H, dt, $J = 1.8, 7.5$ Hz), 8.24 (1H, dd, $J = 1.8, 7.8$ Hz).

d. *trans*-N2-(4-Aminomethyl)cyclohexyl-4,5-dihydro-benzo[2,3]oxepino[4,5-*d*][1,3]thiazol-2-amine

dihydrochloride: *trans*-N2-(4-(*t*-butyloxy-carbonylamino-methyl)cyclohexyl-4,5-dihydro-benzo-[2,3]oxepino[4,5-

d][1,3]thiazol-2-amine (1.97 g, 4.58 mmol) in EtOAc (50 mL) and 4M HCl in dioxane (12 mL) was stirred at room temperature overnight. The solvent was evaporated to give a yellow solid (1.85 g, 99%): m.p. 245 °C (dec.); ^1H NMR (300 MHz, CD_3OD) δ 1.20-1.30 (2H, m), 1.45-1.60 (2H, m), 1.70 (1H, m), 1.95 (2H, d, $J = 9.8$ Hz), 2.21 (2H, d, $J = 10.8$ Hz), 2.83 (2H, d, $J = 6.9$ Hz), 3.17 (2H, t, $J = 5.4$ Hz), 3.50-3.70 (1H, m), 4.33 (2H, t, $J = 5.5$ Hz), 7.11 (1H, d, $J = 8.1$ Hz), 7.21 (1H, t, $J = 7.5$ Hz), 7.36 (1H, t, $J = 8.1$ Hz), 7.70 (1H, d, $J = 7.8$ Hz).

e. *trans*-N2-(4-Aminosulfonylaminomethyl)cyclohexyl-4,5-dihydro-benzo[2,3]oxepino[4,5-d][1,3]thiazol-2-amine: A mixture of *trans*-N2-(4-aminomethyl)-cyclohexyl-4,5-dihydro-benzo[2,3]oxepino-[4,5-d][1,3]thiazol-2-amine dihydrochloride (402 mg, 1 mmol), sulfamide (288 mg, 3 mmol) and triethylamine (0.56 ml) in ethanol (2.5 ml) and water (2.5 ml) was heated at reflux for 3 hours before it was poured into water (50 ml). The product was extracted into chloroform and then purified by flash chromatography over silica gel (EtOAc/hexane 2:1) to afford the desired product as a white solid (362 mg, 89%): m.p. 170-173 °C; CIMS m/e = 409 (MH⁺); ¹H NMR (300 MHz, CDCl₃) δ 1.0-1.3 (4H, m), 1.5-1.7 (1H, m), 1.89 (2H, d, *J* = 12.3 Hz), 2.26 (2H, d, *J* = 11.5 Hz), 3.01 (2H, t, *J* = 6.5 Hz), 3.15 (2H, t, *J* = 5.2 Hz), 3.20-3.40 (1H, m), 4.33 (2H, t, *J* = 5.1 Hz), 4.43 (1H, t, *J* = 6.1 Hz), 4.59 (1H, s, broad), 4.92 (1H, s, broad), 6.99 (1H, d, *J* = 7.9 Hz), 7.08 (1H, t, *J* = 7.4 Hz), 7.16 (1H, dt, *J* = 1.7, 7.2 Hz), 8.23 (1H, dd, *J* = 1.7, 7.8 Hz).

Example 47

trans-N2-(4-Methylsulfonylaminomethyl)cyclohexyl-4,5-dihydro-benzo[2,3]oxepino[4,5-d][1,3]thiazol-2-amine: A mixture of *trans*-N2-(4-aminomethyl)cyclohexyl-4,5-dihydro-benzo[2,3]oxepino[4,5-d][1,3]thiazol-2-amine dihydrochloride (345 mg, 0.87 mmol) and triethylamine (1.1 ml) in THF (5 ml) was cooled in an ice water bath and treated with methanesulfonyl chloride (0.087 ml). After stirring for 30 minutes at room temperature, the mixture was poured into water (50 ml) containing potassium carbonate (1 g). The product was extracted into chloroform and purified by flash chromatography

121

(EtOAc/hexane 2:1) to give a yellow solid (308 mg, 88% yield): m.p. 67-69 °C; CIMS m/e = 408 (MH⁺); ¹H NMR (300 MHz, CDCl₃) δ 0.90-1.40 (5H, m), 1.76 (2H, d, J = 11.7 Hz), 2.15 (2H, d, J = 10.5 Hz), 2.80-3.00 (5H, m), 3.14 (3H, m), 4.30 (2H, t, J = 4.7 Hz), 5.25 (1H, t, J = 4.7 Hz), 5.35 (1H, d, J = 7.9 Hz), 6.98 (1H, d, J = 7.8 Hz), 7.06 (1H, t, J = 7.5 Hz), 7.14 (1H, t, J = 7.4 Hz), 8.19 (1H, d, J = 7.6 Hz).

10

Example 48

trans-1-Aza-4,5-dihydro-2-{4-(2-methoxy-5-methyl)phenyl-sulfonylaminomethyl}cyclohexylamino-6-oxa-3-thia-benzo[e]azulene: 91% yield; 514 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 8.20 (dd, 1H, J=1.6, 7.8 Hz), 7.68 (d, 1H, J=2.1 Hz), 7.30 (dd, 1H, J=2.1, 8.4 Hz), 7.10 (apparent dt, 1H, J=1.1, 7.4 Hz), 7.01 (apparent dt, 1H, J=1.1, 7.5 Hz), 6.96 (d, 1H, J=7.1 Hz), 6.90 (d, 1H, J=8.5 Hz), 5.27 (t, 2H, J=3.8 Hz), 4.29 (t, 2H, 5.2 Hz), 3.90 (s, 3H), 3.18 (m, 1H), 3.11 (t, 2H, J= 5.1 Hz), 2.66 (t, 2H, J=6.5 Hz), 2.31 (s, 3H), 1.95 (ABm, 4H), 1.40-0.80 (m, 8H); m.p. (HCl salt) 100-105 °C (decomp.)

25

Example 49

N1-(9-Fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-[d][1,3]-thiazol-2-yl)-5-[(2-methoxy-5-methylphenyl)-sulfonyl]-aminopentanamide

N1-(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d]-[1,3]-thiazol-2-yl)-5-aminopentanamide (42 mg, 0.13 mmole) was dissolved in 2 ml of dry THF, cooled in an ice bath, triethylamine was added (36 µl, 0.26 mmole) followed by addition of 9 mg of o-methoxy-m-toluenesulfonyl chloride (0.13 mmole). The reaction mixture was stirred at room

122

temperature for 6 hours, quenched with saturated sodium carbonate solution and extracted with 3x3 ml of ethyl acetate. The combined ethyl acetate extracts were washed with saturated sodium chloride solution, concentrated and chromatographed (13 g of silica gel, EtOAc/hexanes/7N NH₃ in MeOH, 50:50:2) to give the desired product: m.p. 70-75 °C; 518 (ESMS, MH⁺)

Example 50

trans-N2-(4-Methylsulfonylamino)cyclohexylmethyl-4,5-dihydro-benzo[2,3]oxepino[4,5-*d*][1,3]thiazol-2-amine as a yellow solid: m.p. 64-67 °C; 408 (ESMS, MH⁺); ¹H NMR (300 MHz, CDCl₃) δ 1.0-1.4 (4H, m), 1.92 (2H, d, *J* = 12.7 Hz), 2.12 (2H, d, *J* = 11.7 Hz), 2.98 (3H, s), 3.16 (4H, m), 3.25 (1H, m), 4.12 (1H, d, *J* = 7.5 Hz), 4.33 (2H, t, *J* = 5.0 Hz), 5.1 (1H, broad), 7.07 (1H, d, *J* = 7.2 Hz), 7.11 (1H, t, *J* = 7.9 Hz), 7.16 (1H, t, *J* = 7.0 Hz), 8.24 (1H, d, *J* = 7.5 Hz).

Example 51

trans-1-Aza-4,5-dihydro-2-{4-(2-methoxy-5-methyl-phenyl)-sulfonylamino}cyclohexylmethylamino-6-oxa-3-thia-benzo[*e*]azulene in 41% yield: m.p. 220-224 °C; 514 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 8.22 (dd, 1H, *J*=1.6, 7.8 Hz), 7.71 (d, 1H, *J*=2.1 Hz), 7.32 (dd, 1H, *J*=2.1, 8.4 Hz), 7.18 (apparent dt, 1H, *J*=1.1, 7.4 Hz), 7.05 (apparent dt, 1H, *J*=1.1, 7.5 Hz), 6.97 (d, 1H, *J*=7.1 Hz), 6.91 (d, 1H, *J*=8.5 Hz), 5.13 (t, 1H, *J*=6.0 Hz), 4.83 (d, 1H, *J*=7.2 Hz), 4.31 (t, 2H, *J*=5.2 Hz), 3.93 (s, 3H), 3.13 (t, 2H, *J*=5.2 Hz), 3.05 (t, 2H, *J*=6.3 Hz), 3.02 (m, 1H), 2.34 (s, 3H), 1.90-0.80 (m, 12H).

Example 52

trans-N2-(4-Ethylsulfonylamino)cyclohexylmethyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine as a white solid: m.p. 140-142 °C; CIMS *m/e* = 424 (MH⁺); ¹H NMR (300 MHz, CDCl₃) δ 1.00-1.40 (4H, m), 1.37 (3H, t, *J* = 7.4 Hz), 1.91 (2H, d, *J* = 12.5 Hz), 2.00-2.20 (4H, m), 2.79 (2H, t, *J* = 5.5 Hz), 2.86 (2H, t, *J* = 6.9 Hz), 3.04 (2H, q, *J* = 7.4 Hz), 3.13 (2H, t, *J* = 6.3 Hz), 3.10-3.30 (1H, m), 4.04 (1H, d, *J* = 7.8 Hz), 5.09 (1H, broad), 6.84 (1H, dt, *J* = 2.8, 8.2 Hz), 7.06 (1H, dd, *J* = 5.9, 8.3 Hz), 7.68 (1H, dd, *J* = 2.8, 10.8 Hz).

Example 53

trans-1-Aza-9-fluoro-4,5-dihydro-2-{4-isopropylsulfonylamino}cyclohexylmethylamino-3-thia-benzo[e]-azulene in 15% yield: 452 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 7.76 (dd, 1H, *J*=2.7, 10.6 Hz), 7.05 (dd, 1H, *J*=6.0, 8.2 Hz); 6.82 (apparent dt, 1H, *J*=2.8, 8.3 Hz), 5.69 (m, 1H), 4.56 (d, 1H, *J*=8.4 Hz), 3.20-2.90 (m, 4H); 2.83 (t, 2H, *J*=7.0 Hz), 2.78 (m, 2H), 2.20-0.80 (m, 12H), 1.48 (d, 6H, *J*=7.0 Hz).

Example 54

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-(3-pyridylsulfonylamino)cyclohexyl)amino-3-thia-benzo[e]azulene (hydrogen chloride salt) was obtained as a white solid in 65% yield: m.p. (HCl salt) 165-167 °C; ¹H NMR (CD₃OD) δ 9.27 (s, 1H), 9.02 (m, 1H), 8.86 (d, 1H, *J* = 8.0 Hz), 8.15 (m, 1H), 7.34 (m, 2H), 7.09 (td, 1H, *J* = 2.3, 8.6 Hz), 3.58 (b, 1H), 3.27 (b, 1H), 2.76 (m, 4H), 2.21-2.07 (m, 4H), 1.92 (m, 2H), 1.47 (m, 4H); 473 (ESMS, MH⁺).

Example 55

1-Aza-9-fluoro-4,5-dihydro-2-(5-(3-pyridyl)sulfonyl-
5 amino)pentylamino-3-thia-benzo[e]azulene (hydrogen
chloride salt) was obtained as a white solid in 66% yield:
m.p. (HCl salt) 84-86 °C; ¹H NMR (free base, CD₃OD) δ 9.25
(broad, 1H), 9.05 (b, 1H), 8.85 (d, 1H, J = 8.1 Hz), 8.18
(m, 1H), 7.36 (m, 2H), 7.09 (td, 1H, J = 2.4, 8.4 Hz),
10 3.47 (t, 2H, J = 6.9 Hz), 3.02 (t, 2H, J = 6.6 Hz), 2.78
(m, 4H), 2.17 (m, 2H), 1.75 (m, 2H), 1.64-1.48 (m, 4H);
461 (ESMS, MH⁺).

Example 56

15 1-Aza-9-fluoro-4,5-dihydro-2-(4-(3-pyridyl)sulfonyl-
amino)butylamino-3-thia-benzo[e]azulene (hydrogen chloride
salt) was obtained as a light yellow solid in 56% yield:
m.p. (HCl salt) 101-103 °C; ¹H NMR (free base, CD₃OD) δ
20 9.13 (m, 1H), 8.91 (m, 1H), 8.59 (m, 1H), 7.94 (m, 1H),
7.36 (m, 2H), 7.10 (td, 1H, J = 2.7, 8.4 Hz), 3.48 (t, 2H,
J = 6.9 Hz), 3.02 (t, 2H, J = 6.5 Hz), 2.78 (m, 4H), 2.19
(m, 2H), 1.80 (m, 2H), 1.65 (m, 2H); 447 (ESMS, MH⁺).

Example 57

25 1-Aza-9-fluoro-4,5-dihydro-2-{2-(2-methylsulfonyl-
amino)ethoxy}ethylamino-3-thia-benzo[e]azulene in 76%
yield: m.p. 159-162 °C; ¹H NMR (CDCl₃) δ 7.75 (dd, 1H,
30 J=2.5, 10.7 Hz), 7.06 (dd, 1H, J=6.0, 8.1 Hz), 6.83
(apparent dt, 1H, J=2.8, 8.3 Hz), 5.42 (broad, 1H), 5.14
(b, 1H), 3.69 (t, 2H, J=4.9 Hz), 3.61 (t, 2H, J=4.8 Hz),
3.51 (broad q, 2H, J=4.3 Hz), 3.31 (q, 2H, J=4.8 Hz), 2.98
(s, 3H), 2.86 (t, 2H, J=6.9 Hz), 2.07 (m, 2H).

Example 58

1-Aza-9-fluoro-4,5-dihydro-2-{2-[2-(2-methoxy-5-methylphenyl)sulfonylamino]ethoxy}ethylamino-3-thia-
benzo[e]azulene in 87% yield: 506 (ESMS, MH⁺); ¹H NMR
(CDCl₃) δ 7.73-7.68 (m, 2H), 7.34 (dd, 1H, J=2.1, 10.1 Hz),
6.94 (d, 1H, J=8.4 Hz), 6.84 (apparent dt, 1H, J=2.8, 8.2
Hz), 5.43 (t, 1H, J=6.0 Hz), 5.23 (m, 1H), 3.94 (s, 3H),
3.51-3.43 (m, 4H), 3.09 (q, 2H, J=5.5 Hz), 2.87 (t, 2H,
J=6.9 Hz), 2.82-2.78 (m, 2H), 2.12-2.00 (m, 2H); m.p. (HCl
salt) 56-60 °C.

Example 59

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-(3-pyridyl)-
sulfonylaminomethyl)cyclohexylamino-3-thia-benzo[e]-
azulene (hydrogen chloride salt) was obtained as a light
yellow solid in 69% yield: m.p. (HCl salt) 127-129 °C; ¹H
NMR (free base, CD₃OD) δ 9.08 (d, 1H, J = 1.9 Hz), 8.81 (d,
1H, J = 4.8 Hz), 8.15 (m, 1H), 7.47 (m, 2H), 7.05 (dd, 1H,
J = 6.0, 8.2 Hz), 6.80 (td, 1H, J = 2.7, 8.2 Hz), 5.98 (m,
1H), 5.33 (m, 1H), 3.14 (m, 1H), 2.83 (t, 2H, J = 6.9 Hz),
2.76 (m, 4H), 2.10 (m, 4H), 1.73 (m, 2H), 1.23 (m, 1H),
1.10 (m, 2H), 0.95 (m, 2H); 487 (ESMS, MH⁺).

Example 60

trans-N2-(4-Ethylsulfonylamino)cyclohexylmethyl-8-methoxy-
4,5-dihydro-benzo[2,3]oxepino[4,5-d][1,3]thiazol-2-amine
as a yellow solid: m.p. 170-172 °C; 452 (ESMS, MH⁺); ¹H NMR
δ 1.20 (4H, m), 1.37 (3H, t, J = 7.4 Hz), 1.90 (2H, m),
2.10 (2H, m), 3.04 (2H, q, J = 7.4 Hz), 3.13 (4H, m), 3.25
(1H, broad), 3.80 (3H, s), 3.96 (1H, d, J = 7.7 Hz), 4.33

126

(2H, t, $J = 5.1$ Hz), 5.02 (1H, broad), 6.55 (1H, d, $J = 2.5$ Hz), 6.66 (1H, dd, $J = 2.6, 8.6$ Hz), 8.15 (1H, d, $J = 8.8$ Hz).

5

Example 61

trans-1-Aza-4,5-dihydro-8-methoxy-2-{4-methyl-sulfonyl-amino)cyclohexylmethylamino-6-oxa-3-thia-benzo[e]azulene in 59% yield: m.p. 65-70 °C; 438 (ESMS, MH^+); 1H NMR (CDCl₃) δ 8.15 (d, 1H, $J=8.9$ Hz), 6.66 (dd, 1H, $J=2.5, 8.8$ Hz), 6.66 (d, 1H, $J=2.5$ Hz), 5.08 (b, 1H), 4.33 (t, 2H, $J=5.1$ Hz), 4.08 (d, 1H, $J=7.3$ Hz), 3.80 (s, 3H), 3.25 (m, 1H), 3.13 (q, 2H, $J=5.3$ Hz), 2.98 (s, 3H), 2.02-1.00 (m, 9H).

15

Example 62

trans-1-Aza-9-fluoro-4,5-dihydro-2-{4-(3-pyridyl)sulfonylamino)cyclohexylmethylamino-3-thia-benzo[e]azulene in 63% yield: m.p. 158-160 °C; 487 (ESMS, MH^+); 1H NMR (CDCl₃) δ 9.10 (d, 1H, $J=2.2$ Hz), 8.81 (dd, 1H, $J=1.5, 4.8$ Hz), 8.18 (apparent dt, 1H, $J = 4.1, 1.5$ Hz), 7.65 (dd, 1H, $J=2.8, 10.8$ Hz), 7.48 (dd, 1H, $J=4.8, 5.4$ Hz), 7.05 (dd, 1H, $J=5.9, 8.3$ Hz), 6.83 (apparent dt, 1H, $J=2.8, 8.2$ Hz), 5.06 (b, 1H), 4.59 (d, 1H, $J=7.7$ Hz), 3.19-3.00 (m, 1H), 3.10 (t, 2H, $J=6.1$ Hz), 2.85 (t, 2H, $J=3.4$ Hz), 2.80-2.76 (m, 2H), 2.10-0.80 (m, 9H).

25

Example 63

30

trans-1-Aza-4,5-dihydro-9-methoxy-2-{4-methylsulfonyl-amino)cyclohexylmethylamino-6-oxa-3-thia-benzo[e]azulene in 54% yield: 438 (ESMS, MH^+); 1H NMR (CDCl₃) δ 7.80 (d, 1H, $J=3.2$ Hz), 6.92 (d, 1H, $J=8.7$ Hz), 6.72 (dd, 1H,

127

J=3.2, 8.8 Hz), 5.04 (broad m, 1H), 4.28 (t, 2H, J=5.1 Hz), 4.12 (q, 2H, J=7.2 Hz), 4.11 (m, 1H), 3.83 (s, 3H), 3.40-3.13 (m, 5H), 2.99 (s, 3H), 2.20-1.00 (m, 9H); m.p. (HCl salt) 200-215 °C.

5

Example 64

trans-N2-(4-Ethylsulfonylamino)cyclohexylmethyl-9-methoxy-4,5-dihydro-benzo[2,3]oxepino[4,5-d][1,3]thiazol-2-amine

10 as a gray solid: m.p. 165-170 °C; 452 (ESMS, MH⁺); ¹H NMR δ 1.1-1.3 (4H, m), 1.37 (3H, t, J = 7.4 Hz), 1.92 (2H, m), 2.12 (2H, m), 3.04 (2H, q, J = 7.4 Hz), 3.14 (4H, t, J = 5.2 Hz), 3.25 (1H, m), 3.83 (3H, s), 3.95 (1H, d, J = 8.1 Hz), 4.28 (2H, t, J = 5.2 Hz), 5.07 (1H, broad), 6.72 (1H, dd, J = 3.1, 8.7 Hz), 6.92 (1H, d, J = 8.7 Hz), 7.79 (1H, d, J = 3.1 Hz).

15

Example 65

20 *trans*-N2-(4-Methylsulfonylamino)cyclohexylmethyl-7-methoxy-4,5-dihydro-benzo[2,3]oxepino[4,5-d][1,3]-thiazol-2-amine hydrochloride as a beige solid: m.p. 225-230 °C; 438 (ESMS, MH⁺); ¹H NMR (free base) δ 1.10-1.30 (4H, m), 1.93 (2H, m), 2.15 (2H, m), 2.98 (3H, s), 3.16 (4H, m), 3.28 (1H, m), 3.88 (3H, s), 4.40 (2H, t, J = 5.2 Hz), 5.02 (1H, broad), 6.82 (1H, dd, J = 1.5, 8.0 Hz), 7.02 (1H, t, J = 8.1 Hz), 7.85 (1H, dd, J = 1.5, 8.1 Hz).

25

Example 66

30 *trans*-1-Aza-4,5-dihydro-7-methoxy-2-{4-dimethylamino-sulfonylamino}cyclohexylmethylamino-6-oxa-3-thia-benzo[e]azulene in 27% yield: 467 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 7.84 (dd, 1H, J=1.4, 8.1 Hz), 7.02 (apparent t,

30

1H, J=8.1 Hz), 6.82 (dd, ¹²⁸1H, J=1.4, 8.0 Hz), 5.11 (m, 1H), 4.41-4.33 (m, 4H), 3.96 (d, 1H, J=8.0 Hz), 3.88 (s, 3H), 3.19-3.10 (m, 4H), 2.78 (s, 6H), 2.20-1.00 (m, 8H); m.p. (HCl salt) 206-208 °C.

5

Example 67

trans-N2-(4-Dimethylphosphonylamino)cyclohexylmethyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine as a yellow solid: m.p. 58-62 °C; CIMS m/e = 454 (MH⁺); ¹H NMR (300 MHz, CDCl₃) δ 1.04 (4H, m), 1.48 (1H, broad), 1.80 (2H, m), 2.00 (2H, m), 2.10 (2H, m), 2.80 (6H, m), 3.04 (2H, m), 3.67 (3H, s), 3.71 (3H, s), 5.62 (1H, broad), 6.81 (1H, m), 7.04 (1H, m), 7.66 (1H, dd, J = 2.6, 10.8 Hz).

15

Example 68

trans-N2-(4-Ethoxycarbonylamino)cyclohexylmethyl-9-fluoro-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine hydrochloride as a white solid: m.p. (HCl salt) 94-97 °C; ¹H NMR (free base, 300 MHz, CDCl₃) δ 1.10 (4H, m), 1.23 (3H, t, J = 7.3 Hz), 1.57 (1H, m), 1.86 (2H, m), 2.04 (4H, m), 2.79 (2H, m), 2.86 (2H, t, J = 6.9 Hz), 3.11 (2H, t, J = 6.2 Hz), 3.44 (1H, m), 4.11 (2H, q, J = 7.2 Hz), 4.50 (1H, broad), 5.24 (1H, broad), 6.83 (1H, dt, J = 2.8, 8.2 Hz), 7.06 (1H, dd, J = 5.9, 8.3 Hz), 7.69 (1H, dd, J = 2.8, 10.8 Hz).

25

30

Example 69

1-Aza-9-fluoro-4,5-dihydro-2-(2-(2-isopropylsulfonylamino)-ethoxy)ethylamino-3-thia-benzo[e]-azulene in 60% yield: m.p. (HCl salt) 50-55 °C; ¹H NMR (300 MHz,

129

CDCl₃) δ 1.00 1.37 (d, 6H, $J = 6.9$ Hz), 2.08 (m, 2H),
2.75-2.90 (m, 4H), 3.18 (septet, 1H, $J = 6.9$ Hz), 3.33
(apparent q, 2H, $J = 5.0$ Hz), 3.53 (apparent q, 2H, $J = 5.0$
Hz), 3.61 (t, 2H, $J = 5.0$ Hz), 3.71 (t, 2H, $J = 4.8$ Hz), 4.68
5 (t, 1H, d, $J = 5.5$ Hz), 5.38 (1H, broad), 6.84 (1H, dt, J
= 2.8, 8.2 Hz), 7.07 (1H, dd, $J = 5.9, 8.3$ Hz), 7.70 (1H,
dd, $J = 2.8, 10.8$ Hz).

10

Example 70

15

2-(4-Methylsulfonylaminomethyl)cyclohexylamino-4H-
chromeno[4,3-d]thiazole (hydrogen chloride salt) was
obtained as a blue-green solid in 64% yield: m.p. (HCl
salt) 140-144 °C; ¹H NMR (CDCl₃) δ 7.56 (dd, 1H, $J = 1.3,$
7.6 Hz), 7.13 (m, 1H), 6.96 (td, 1H, $J = 0.9, 7.4$ Hz),
6.88 (d, 1H, $J = 7.4$ Hz), 5.31 (m, 3H), 4.78 (t, 1H, $J =$
6.5 Hz), 3.27 (m, 1H), 2.99 (t, 2H, $J = 6.6$ Hz), 2.97 (s,
3H) 2.23 (m, 2H), 1.87 (m, 2H), 1.49 (m, 1H), 1.30-1.06
20 (m, 4H); 394 (ESMS, MH⁺).

Example 71

25

30

trans-1-Aza-4,5-dihydro-8-methoxy-2-(4-methylsulfonyl-
amino)cyclohexylmethylamino-3-thia-benzo[e]azulene
(hydrogen chloride salt) was obtained as a yellow solid in
50% yield: m.p. (hydrogen chloride salt) 94-97 °C; ¹H NMR
(CD₃OD) δ 7.88 (d, 1H, $J = 8.6$ Hz), 6.78 (dd, 1H, $J = 2.7,$
8.8 Hz), 6.68 (d, 1H, $J = 2.6$ Hz), 5.21 (m, 1H), 4.23 (m,
1H), 3.81 (s, 3H), 3.25 (m, 1H), 3.11 (m, 2H), 2.98 (s,
3H), 2.83 (t, 2H, $J = 7.0$ Hz), 2.81 (m, 2H), 2.10 (m, 4H),
1.89 (m, 2H), 1.59 (m, 1H), 1.31-1.07 (m, 4H); 436 (ESMS,
MH⁺).

Example 72

trans-1-Aza-4,5-dihydro-8-methoxy-2-(4-methylsulfonylamino-methyl)cyclohexylamino-3-thia-
benzo[e]azulene was obtained as a yellow solid in 20%
yield: m.p. 224-225 °C; ¹H NMR δ 7.87 (d, 1H, *J* = 8.7 Hz),
6.78 (dd, 1H, *J* = 2.7, 8.5 Hz), 6.68 (d, 1H, *J* = 2.5 Hz),
4.94 (m, 1H), 4.44 (m, 1H), 3.81 (s, 3H), 3.26 (m, 1H),
2.98 (m, 2H), 2.96 (s, 3H), 2.83 (m, 4H), 2.27 (m, 2H),
2.10 (m, 2H), 1.88 (m, 2H), 1.51 (m, 1H), 1.27-1.04 (m,
4H); 436 (ESMS, MH⁺).

Example 73

trans-1-Aza-4,5-dihydro-2-(4-isopropylsulfonylaminomethyl)-cyclohexylamino-8-methoxy-3-thia-benzo[e]-
azulene (hydrogen chloride salt) was obtained as a light
brown solid in 3.4% yield: m.p. (HCl salt) 87-89 °C; ¹H NMR
(free base, CDCl₃) δ 7.87 (d, 1H, *J* = 8.4 Hz), 6.8 (dd, 1H,
J = 2.7, 8.4 Hz), 6.69 (d, 1H, *J* = 2.7 Hz), 5.10 (b, 1H),
4.19 (m, 1H), 3.81 (s, 3H), 3.29-3.12 (m, 2H), 2.98 (t,
2H, *J* = 6.6 Hz), 2.82 (m, 4H), 2.25 (m, 2H), 2.09 (m, 2H),
1.91 (m, 2H), 1.50 (m, 1H), 1.38 (d, 6H, *J* = 6.6 Hz),
1.30-1.03 (m, 4H); 464 (ESMS, MH⁺).

Example 74

trans-1-Aza-4,5-dihydro-2-(4-methylsulfonylamino-methyl)-
cyclohexylamino-7-methoxy-3-thia-benzo[e]-azulene was
obtained as a light yellow solid in 59% yield: ¹H NMR δ
7.75 (dd, 1H, *J* = 1.2, 8.0 Hz), 7.01 (t, 1H, *J* = 8.1 Hz),
6.81 (dd, 1H, *J* = 1.2, 7.0 Hz), 4.81 (m, 1H), 4.40 (t, 2H,

$J = 5.1$ Hz), 4.34 (m, 1H), ¹³¹ 3.88 (s, 3H), 3.34 (m, 1H), 3.18 (t, 2H, $J = 5.0$ Hz), 3.03 (t, 2H, $J = 7.0$ Hz), 2.97 (s, 3H), 2.28 (m, 2H), 1.91 (m, 2H), 1.50 (m, 1H), 1.30-1.08 (m, 4H); 438 (ESMS, MH⁺).

5

Example 75

trans-1-Aza-4,5-dihydro-2-(4-ethylcarbonylamino-methyl)-
10 cyclohexylamino-9-fluoro-3-thia-benzo[e]-azulene was
obtained as a light yellow solid in 87% yield: ¹H NMR δ
7.69 (dd, 1H, $J = 2.7, 10.8$ Hz), 7.06 (dd, 1H, $J = 5.9,$
8.4 Hz), 6.83 (td, 1H, $J = 2.8, 8.2$ Hz), 5.50 (m, 1H),
4.82 (m, 1H), 3.29 (m, 1H), 3.15 (t, 2H, $J = 6.5$ Hz), 2.86
15 (t, 2H, $J = 6.9$ Hz), 2.79 (m, 2H), 2.21 (m, 4H), 2.07 (m,
2H), 1.84 (m, 2H), 1.51 (m, 1H), 1.29-1.03 (m, 4H), 1.17
(t, 3H, $J = 7.5$ Hz); 402 (ESMS, MH⁺).

Example 76

20 trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-(4-morpholinyl)-
sulfonylaminomethyl)cyclohexylamino-3-thia-benzo[e]azulene
(hydrogen chloride salt) was obtained as a light yellow
solid in 28% yield: ¹H NMR (free base) δ 7.67 (dd, 1H, $J =$
25 2.8, 10.8 Hz), 7.06 (dd, 1H, $J = 5.9, 8.0$ Hz), 6.83 (td,
1H, $J = 2.7, 8.1$ Hz), 4.96 (d, 1H, $J = 8.0$ Hz), 4.59 (t,
1H, $J = 6.1$ Hz), 3.74 (m, 4H), 3.23 (m, 5H), 2.93 (t, 2H,
 $J = 4.8$ Hz), 2.86 (t, 2H, $J = 6.9$ Hz), 2.79 (m, 2H), 2.24
(m, 2H), 2.07 (m, 2H), 1.84 (m, 2H), 1.46 (m, 1H), 1.30-
30 1.02 (m, 4H); 495 (ESMS, MH⁺).

Example 77

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-(2-methoxy)-ethoxy-carbonylaminoethyl)cyclohexylamino-3-thia-benzo[e]azulene
5 2-methoxyethyl N-({4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[d][1,3]thiazol-2-yl)amino]cyclohexyl)methyl}-carbamate:

tert-butyl N-[(4-{[(benzoylamino)carbothioyl]amino}-cyclo-
10 hexyl)methyl]carbamate: A solution of tert-butyl N-[(4-aminocyclohexyl)methyl]carbamate (4.683 g, 20.5 mmol) and benzoyl isothiocyanate (3.77 g, 23.1 mmol) in 180 ml THF was stirred at room temperature for 20 hours. The solvent was removed *in vacuo* to yield a golden viscous oil. The
15 oil was triturated with 250 ml hexane and the resulting solid was collected by filtration and washed with hexane. The off-white solid was washed several times with hexane and was dried under vacuum to yield 91% (7.32 g) of the desired product as an off-white solid: ¹H-NMR (CDCl₃) δ
20 1.20 (6H, m), 1.44 (9H, s), 1.83 (2H, m), 2.25 (2H, m), 3.01 (2H, t, J = 6.4 Hz), 4.20 (1H, m), 4.62 (1H, m), 7.51 (2H, t, J = 7.9 Hz), 7.60 (1H, t, J = 7.4 Hz), 7.81 (2H, d, J = 7.2 Hz), 8.93 (1H, s); 392 (ESMS, MH⁺).

tert-butyl-N-({4-[(aminocarbothioyl)amino]-cyclohexyl}-
25 methyl)carbamate : A solution of tert-butyl-N-[(4-{[(benzoylamino)carbothioyl]amino}-cyclohexyl)methyl]carbamate (7.30 g, 18.6 mmol) in 110 ml MeOH was prepared. Then, a solution of potassium carbonate (5.15 g, 37.3 mmol) in 50 ml water was added to
30 the stirred solution. The solution was stirred at room temperature for 18 hours. The solvent was removed *in vacuo* and the residue was extracted several times with EtOH. The combined EtOH extracts were evaporated to

133

dryness to yield 6.15 g of an off-white solid. The solid was re-dissolved in 200 ml acetone and stirred for 20 minutes. The solution was filtered by suction to remove the insoluble salts and the filtrate was evaporated to yield 98% (5.23g) of the desired product as a slightly yellow solid: $^1\text{H-NMR}$ (CDCl_3) δ 1.10 (6H, m), 1.43 (9H, s), 1.80 (2H, m), 2.10 (2H, m), 2.95 (2H, t, $J = 6.4$ Hz), 4.75 (1H, broad), 6.03 (2H, s), 6.64 (1H, broad); 288 (ESMS, MH^+).

6-Bromo-3-fluoro-6,7,8,9-tetrahydro-5H-

benzo[a]cyclohepten-5-one: A solution of 8-fluoro-1-benzosuberone (20.00g, 112.2 mmol) in 125 ml glacial acetic acid was prepared. Bromine (5.80 ml, 113 mmol) was added dropwise with stirring. After addition was complete, the solution was poured into 400 ml water and extracted with EtOAc (500 ml). The EtOAc layer was separated and washed with 500 ml water. The organic layer was dried over MgSO_4 , filtered and evaporated to yield 100% (28.9 g) of the desired product as a viscous oil: $^1\text{H-NMR}$ (CDCl_3) δ 2.00 (2H, m), 2.33 (2H, m), 2.84 (1H, m), 3.00 (1H, m), 4.83 (1H, m), 7.13 (2H, m), 7.29 (1H, dd).

tert-Butyl-N-({4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]-cyclohepta-[d][1,3]thiazol-2-yl)amino]-cyclohexyl}methyl)-carbamate: A solution of tert-butyl-N-({4-[(aminocarbothioyl)amino]cyclohexyl}-methyl)carbamate (22.4 g, 78.0 mmol), 6-bromo-3-fluoro-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one (20.1 g, 78.0 mmol), and 22.5 ml diisopropylethylamine in 260 ml absolute ethanol was heated at reflux for 3 hours. The solvent was removed in vacuo with gentle heating (~ 40 C). The resulting viscous golden oil was taken up into 400 ml EtOAc and washed with water (600 ml). The EtOAc layer was separated and

134

evaporated to yield 39.0 g of a viscous golden oil. The oil was chromatographed on silica-gel eluting with EtOAc to yield 34.5 g of the desired product (99% yield): ¹H-NMR (CDCl₃) δ 1.18 (6H, h, J = 12.2 Hz), 1.44 (9H, s), 1.81 (2H, m), 2.22 (2H, m), 2.79 (2H, m), 2.86 (2H, t, J = 7.1 Hz), 3.00 (2H, t, J = 6.3 Hz), 3.25 (1H, broad), 4.62 (1H, broad), 4.86 (1H, broad), 6.83 (1H, dt, J = 2.9 Hz), 7.04 (1H, m), 7.70 (1H, dd, J = 2.8 Hz).

10 trans-N2-[4-(Aminomethyl)cyclohexyl]-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine: A solution of trans-tert-butyl N-({4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-[d][1,3]thiazol-2-yl)amino]cyclohexyl}-methyl)-carbamate (34.5 g, 77.0 mmol) in 200 ml EtOAc and 100 ml 4N HCl in dioxane was stirred at room temperature for 2 hours. The resulting solid was collected by filtration and the solid product washed with Et₂O and dried under vacuum to yield 25.9 g of the desired product as a pale yellow solid. An additional 1.43g of the product was obtained by allowing the filtrate to stand in a closed flask for several days. The overall yield was 85%: ¹H-NMR (CD₃OD) δ 1.25 (2H, m), 1.51 (2H, m), 1.70 (1H, broad), 1.95 (2H, m), 2.18 (4H, m), 2.80 (6H, m), 3.63 (1H, broad), 7.11 (1H, dt, J = 2.5Hz, 8.4 Hz), 7.36 (2H, m); 346 (ESMS, MH⁺).

trans-1-Aza-9-fluoro-4,5-dihydro-2-(4-(2-methoxy)-ethoxy-carbonylaminoethyl)cyclohexylamino-3-thia-benzo[e]azulene
2-methoxyethyl N-({4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-2-yl)amino]cyclohexyl}-methyl)carbamate: A solution trans-N2-[4-(aminomethyl)-cyclohexyl]-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-[d][1,3]thiazol-2-amine (0.30 g, 0.72 mmol) in anhydrous THF (3 ml) and triethylamine (1.0 ml)

135

was flushed with nitrogen. The solution was stirred for 5 minutes and 0.095 ml chloroformic acid 2-methoxyethyl ester (0.114g, 0.82 mmol) was added dropwise to the stirring solution via syringe. The solution was stirred at room temperature for 2 hours. The solvent was removed *in vacuo* and the residue was chromatographed on silica-gel eluting with 1:1 EtOAc/Hexane to yield 25% (80 mg) of the desired product as a colorless viscous oil: ^1H -NMR (CDCl_3) δ 1.11 (4H, septet, $J = 9.8$ Hz), 1.42 (1H, broad), 1.80 (2H, m), 2.04 (2H, m), 2.20 (2H, m), 2.77 (2H, m), 2.84 (2H, t, $J = 6.9$ Hz), 3.04 (2H, t, $J = 6.4$ Hz), 3.24 (1H, broad), 3.38 (3H, s), 3.58 (2H, m), 4.22 (2H, m), 4.90 (1H, broad), 5.00 (1H, d), 6.81 (1H, dt, $J = 2.8$ Hz), 7.03 (1H, m), 7.68 (1H, dd, $J = 2.7$ Hz, 10.9 Hz); 448 (ESMS, MH^+).

Example 78

trans-N2-(4-(1-Morpholinylsulfonylaminomethyl)-cyclohexyl-8-methoxy-5,6-dihydro-4*H*-benzo[6,7]-cyclohepta[d][1,3]thiazol-2-amine hydrochloride: Prepared from *trans*-N2-(4-aminomethyl)cyclohexyl-8-methoxy-5,6-dihydro-4*H*-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine and 1-morpholinylsulfonyl chloride as a light brown solid: m.p. 130-132 °C; 507 (ESMS, MH^+); ^1H NMR (free base) δ 7.88 (dd, 1H, $J = 0.4$, 8.7 Hz), 6.78 (dd, 1H, $J = 2.4$, 8.8 Hz), 6.68 (d, 1H, $J = 2.3$ Hz), 4.85 (m, 1H), 4.25 (m, 1H), 3.81 (s, 3H), 3.73 (m, 4H), 3.30 (m, 1H), 3.25 (m, 4H), 2.96 (m, 2H), 2.81 (m, 4H), 2.26 (m, 2H), 2.10 (m, 2H), 1.87 (m, 2H), 1.49 (m, 1H), 1.30-1.03 (m, 4H).

Example 79

3 - ({4 - [(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-
[d][1,3]thiazol-2-yl)amino]cyclohexyl)methyl)-1,3-
5 oxazolan-2-one:

2-chloroethyl-N-({4 - [(9-fluoro-5,6-dihydro-4H-benzo[6,7]-
cyclohepta[d][1,3]thiazol-2-yl)amino]-cyclohexyl)methyl)-
carbamate: A solution of trans-N2-[4-
10 (aminomethyl)cyclohexyl]-9-fluoro-5,6-dihydro-4H-benzo-
[6,7]cyclohepta[d][1,3]thiazol-2-amine (5.00 g, 11.95
mmol), 20 ml anhydrous pyridine, and 8 ml
diisopropylethylamine was stirred under N₂. Then 2-
chloroethyl chloroformate (1.40 ml, 13.6 mmol) was added
15 dropwise to the stirring solution. The solution was
stirred at room temperature for 2 hours and then poured
into 250 ml water and extracted with 250 ml EtOAc. The
extract was separated and evaporated to yield 100% (5.40
g) of the desired product as a highly viscous golden oil:
20 ¹H-NMR (CDCl₃) δ 1.13 (3H, m), 1.43 (1H, m), 1.82 (2H, m),
2.06 (2H, m), 2.22 (2H, m), 2.81 (4H, m), 3.06 (2H, t, J =
6.4 Hz), 3.25 (1H, m), 3.70 (3H, m), 4.31 (2H, t, J = 5.7
Hz), 4.89 (2H, m), 6.82 (1H, t, J = 2.8 Hz), 7.04 (1H, m),
7.70 (1H, dd, J = 2.7, 10.8 Hz); 452 (ESMS, MH⁺).

25 3 - ({4 - [(9-Fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-
[d][1,3]thiazol-2-yl)amino]cyclohexyl)methyl)-1,3-
oxazolan-2-one: To a stirred solution of 2-chloroethyl-N-
({4 - [(9-fluoro-5,6-dihydro-4H-benzo[6,7]-
30 cyclohepta[d][1,3]thiazol-2-yl)amino]-
cyclohexyl)methyl)carbamate (5.4 g, 11.95 mmol) in 100 ml
anhydrous DMF was added NaH (60% in mineral oil) (0.6 g,
15.0 mmol). TLC (EtOAc) indicated that the reaction had
proceeded to approximately 40% completion after 15

137

minutes. An additional 0.58g (14.5 mmol) NaH (60% in mineral oil) was added and the solution was stirred at room temperature for another 15 minutes. TLC indicated the reaction was complete. The solution was poured into water (500 ml) and was extracted with EtOAc. The extract was separated and evaporated to yield a dark viscous oil. The oil was chromatographed on silica-gel eluting with 2:1 EtOAc/hexane to yield a white solid. The solid was dissolved in 40 ml CHCl₃ and 10 ml 1N HCl in Et₂O was added to the solution. The volatiles were removed and the resulting solid was triturated with Et₂O. The solution was filtered and the slightly tan solid washed with ether and dried to yield 52% (2.83 g) of the desired product as the HCl salt: ¹H-NMR (Free-base) (CDCl₃) δ 1.16 (4H, m), 1.60 (1H, m), 1.80 (2H, m), 2.05 (2H, m), 2.22 (2H, m), 2.81 (4H, m), 3.12 (2H, d, J = 7.3Hz), 3.27 (1H, m), 3.57 (2H, t, J = 8.2 Hz), 4.33 (2H, t, J = 8.0 Hz), 5.05 (1H, d, J = 7.7 Hz), 6.83 (1H, dt, J = 2.7, 8.2 Hz), 7.05 (1H, m), 7.68 (1H, dd, J = 2.7, 10.8 Hz). m.p. (HCl salt) 223-225 °C. Anal. Calcd. For C₂₂H₂₇N₃ClFO₂S: C, 58.5; H, 6.0; N, 9.3. Found: C, 58.43; H, 6.06; N, 9.38; 416 (ESMS, MH⁺).

Example 80

N1-({4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[d][1,3]thiazol-2-yl)amino]cyclohexyl}-methyl)-2-methoxyacetamide: To a stirred, septum capped solution of *trans*-N2-[4-(aminomethyl)-cyclohexyl]-9-fluoro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[d][1,3]thiazol-2-amine (5.00g, 11.95 mmol) in anhydrous pyridine (20 ml) and diisopropylethylamine (10 ml) was added methoxyacetyl chloride (1.20 ml, 13.1 mmol) dropwise. The solution was stirred at room temperature for 2 hours and was then poured into 250 ml water and extracted with 250 ml EtOAc.

138

The extract was evaporated to yield a golden viscous oil which was chromatographed on silica-gel eluting with EtOAc to yield 88% (4.40 g) of the desired product as a slightly yellow solid. A small portion of the product was recrystallized from EtOAc (m.p. 95-97 °C). The free-base was dissolved in 60 ml CHCl₃ and converted to its HCl salt by adding 14.0 ml 1.0 N HCl in ether followed by evaporation of the solvent. The resulting foam was stirred with ether (75 ml), filtered and washed with ether to yield the HCl salt as a solid: ¹H-NMR (Free-base) (CDCl₃) δ 1.12 (4H, m), 1.45 (1H, m), 1.81 (2H, m), 2.05 (2H, m), 2.21 (2H, m), 2.80 (4H, m), 3.16 (2H, t, J = 6.5 Hz), 3.25 (1H, m), 3.41 (3H, s), 3.89 (2H, s), 5.03 (1H, d, J = 8.0 Hz), 6.62 (1H, broad t), 6.81 (1H, dt, J = 2.9, 8.3 Hz), 7.03 (1H, m), 7.69 (1H, dd, J = 2.8, 10.8 Hz). m.p. 63-65 °C. Anal. Calcd. For C₂₂H₂₉N₃O₂SFCl₃+0.5CHCl₃: C, 52.61; H, 5.79; N, 8.18. Found: C, 52.61; H, 5.92; N, 8.08; 418 (ESMS, MH⁺).

Example 81

N1-({4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]-cyclohepta-[d][1,3]thiazol-2-yl)amino]cyclohexyl}-methyl)acetamide:

To a stirred, septum capped solution of *trans*-N2-[4-(aminomethyl)cyclohexyl]-9-fluoro-5,6-dihydro-4H-benzo-[6,7]cyclohepta-[d][1,3]thiazol-2-amine (0.165 g, 0.394 mmol) in 3.0 ml anhydrous pyridine was added 0.034 ml (0.038 g, 0.48 mmol) acetyl chloride dropwise. Stirring at room temperature was continued for 90 minutes and the solution was then poured into 50 ml water and extracted with EtOAc (2X50 ml). The combined extracts were evaporated to yield 0.126 g of the crude product as a viscous golden oil. The oil was chromatographed on preparative TLC eluting with EtOAc to yield 28% (0.042 g)

139
of the desired product as a white solid: ^1H -NMR (CDCl_3)
 δ 1.13 (4H, m), 1.43 (1H, m), 1.80 (2H, m), 1.98 (3H, s),
2.05 (2H, m), 2.21 (2H, m), 2.80 (4H, m), 3.10 (2H, t, J =
6.5 Hz), 3.24 (1H, m), 5.00 (1H, d, J = 7.8 Hz), 5.75 (1H,
5 m), 6.81 (1H, dt, J = 2.8, 8.2 Hz), 7.04 (1H, m), 7.69
(1H, dd, J = 2.7, 10.8 Hz); m.p. 88-91°C.

Example 82

10 *trans*-1-Aza-9-fluoro-4,5-dihydro-2-(4-(N-propyl-
formamido)-methyl)cyclohexylamino-3-thia-benzo[e]-azulene
(hydrogen chloride salt) was obtained as a yellow solid in
54% yield: ^1H NMR (free base) δ 8.10 (s, 0.5H), 8.00 (s,
0.5H), 7.68 (m, 1H), 7.04 (m, 1H), 6.83 (m, 1H), 5.07 (m,
15 1H), 3.31-3.02 (m, 5H), 2.85 (m, 4H), 2.23 (m, 2H), 2.05
(m, 2H), 1.77 (m, 2H), 1.58 (m, 3H), 1.26-0.87 (m, 7H);
416 (ESMS, MH^+).

Example 83

20 *trans*-1-Aza-9-fluoro-4,5-dihydro-2-(4-(N-isopropyl-
formamido)methyl)cyclohexylamino-3-thia-benzo[e]-azulene
(hydrogen chloride salt) was obtained as a yellow solid in
48% yield: m.p. (HCl salt) 106-107 °C; ^1H NMR (CD_3OD) δ
25 8.22 (s, 0.6H), 7.98 (s, 0.4H), 7.37 (m, 2H), 7.12 (m,
1H), 4.18 (m, 0.4H), 3.73 (m, 0.6H), 3.60 (m, 1H), 3.17
(m, 2H), 2.78 (m, 4H), 2.19 (m, 4H), 1.86 (m, 2H), 1.74
(m, 1H), 1.41 (m, 2H), 1.32 (d, 4H), 1.31 (d, 2H), 1.17
(m, 2H); 416 (ESMS, MH^+).

30

Example 84

N1-{4-[(4,5-dihydrobenzo[2,3]oxepino[4,5-*d*][1,3]thiazol-2-
ylamino)methyl]cyclohexyl}-2-methoxyacetamide:

Benzyl-N-[4-({[(benzoylamino)carbothioyl]amino}-methyl)-cyclohexyl]carbamate: A solution of benzyl N-[4-(aminomethyl)cyclohexyl]carbamate HCl (3.46 g, 11.6 mmol) and diisopropylethylamine (6.0 ml) in anhydrous THF (90 ml) was stirred at room temperature for 20 minutes. To this solution was added benzoylisothiocyanate (2.00 g, 12.3 mmol) and the reaction mixture stirred at room temperature for 14 hours. The solution was poured into water (200 ml) and extracted with Et₂O (2X150 ml). The organic layer was dried over MgSO₄, filtered and evaporated to yield a solid. The solid was triturated with hexane and filtered to yield 99% (5.023g) of the desired product.

Benzyl-N-(4-{[(aminocarbothioyl)amino]methyl}-cyclohexyl)-carbamate: To a solution of benzyl N-[4-({[(benzoylamino)carbothioyl]amino}-methyl)-cyclohexyl]carbamate (5.023 g, 11.4 mmol) in MeOH (110 ml) was added a solution of K₂CO₃ (3.5 g, 25.3 mmol) in water (30 ml). The solution was stirred for 24 hours at room temperature and the solvent was removed in vacuo with repeated additions of MeOH to azeotropically remove water. The resulting solid was treated with acetone (60 ml) and filtered. The filtrate was evaporated to yield 95% (3.47 g) of a yellow solid.

Benzyl-N-{4-[(4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]-thiazol-2-ylamino)methyl]cyclohexyl}-carbamate: A solution containing benzyl N-(4-{[(aminocarbothioyl)amino]methyl}-cyclohexyl)-carbamate (3.45 g, 10.73 mmol), 4-bromo-2,3,4,5-tetrahydro-1-benzoxepin-5-one (2.60 g, 10.78 mmol), diisopropylethylamine (2.50 ml) and EtOH (50 ml) was heated at reflux for 4 hours. The solvent was removed and

141

the residue was chromatographed on silica-gel eluting with 1:3 EtOAc/Hexane to yield 65% (3.2 g) of the product as a white solid: m.p. 124-127 °C; ¹H-NMR (CDCl₃) δ 1.09 (4H, pentet, J = 10.4 Hz), 1.55 (1H, broad), 1.83 (2H, m), 2.04 (2H, broad), 3.10 (2H, t, J = 6.2 Hz), 3.15 (2H, t, J = 5.2 Hz), 3.43 (1H, broad) 4.33 (2H, t, J = 5.2 Hz), 4.62 (1H, d, J = 8.0 Hz), 5.08 (2H, s), 5.31 (1H, m), 7.00 (1H, dd, J = 1.2 Hz, 7.8 Hz), 7.12 (2H, m), 7.35 (5H, s), 8.25 (1H, dd, J = 1.7 Hz, 7.8 Hz); 464 (ESMS, MH⁺).

N2-[(4-aminocyclohexyl)methyl]-4,5-dihydrobenzo[2,3]-oxepino[4,5-d][1,3]thiazol-2-amine: A solution containing benzyl-N-{4-[(4,5-dihydrobenzo [2,3]oxepino[4,5-d]-[1,3]-thiazol-2-ylamino)methyl]-cyclohexyl}carbamate (3.00 g, 6.47 mmol), glacial acetic acid (50 ml) and conc. HCl (20 ml) was heated to a gentle reflux for 45 minutes. The solvent was removed on the rotor-vap with gentle heating and the residue diluted with 150 ml water. The resulting solution was made basic with NaOH (pH~11). At this point, much of the product precipitated out of solution and stuck to the sides of the flask. EtOAc (200 ml) was added and the solution was shaken vigorously. The phases were separated and the aqueous phase extracted with EtOAc. The combined extracts were washed with water and evaporated to yield 2.30 g of a brown viscous oil. The oil was taken up into 40 ml EtOAc and 5 ml 4M HCl in dioxane added. The resulting dihydrochloride salt was obtained by filtration and washed with EtOAc and then several times with ether. Upon drying, 100% (2.6 g) of the salt was obtained as a slightly tan free-flowing solid: ¹H-NMR (d₄-MeOH) δ 1.21 (2H, m), 1.43 (2H, m), 1.79 (1H, m), 1.99 (2H, m), 2.10 (2H, m), 3.09 (1H, m), 3.18 (2H, t, J = 5.0 Hz), 3.35 (2H, d, J = 6.7 Hz), 4.32 (2H, t, J = 5.3 Hz), 7.12 (1H, dd, J

142

= 1.1, 8.0 Hz), 7.22 (1H, m), 7.33 (1H, m), 7.70 (1H, dd, J = 1.3, 8.0 Hz); 330 (ESMS, MH⁺).

To a stirred solution of N2-[(4-aminocyclohexyl)methyl]-4,5-dihydrobenzo[2,3]oxepino-[4,5-d][1,3]thiazol-2-amine (0.600 g, 1.49 mmol), anhydrous pyridine (3 ml), and diisopropylethylamine (3 ml) was added methoxyacetyl chloride (0.163 ml, 1.78 mmol) dropwise. The solution was stirred for 30 minutes at room temperature and then poured into water (60 ml) and extracted with EtOAc. The extract was evaporated and the residue chromatographed on silica-gel eluting with EtOAc to yield 79% (470 mg) of the product as a white solid. The product was dissolved in chloroform and converted to its HCl salt with 1N HCl in Et₂O to yield 522 mg of the salt as a white solid: m.p. (HCl salt) 234-237°C; ¹H-NMR (free base, CDCl₃) δ 1.20 (4H, m), 1.80 (2H, broad), 2.00 (4H, m), 3.13 (4H, t, J = 5.0 Hz), 3.40 (3H, s), 3.79 (1H, broad), 3.86 (2H, s), 4.34 (2H, t, J = 5.1 Hz), 6.35 (1H, d), 7.05 (1H, dd, J = 1.3 Hz, 7.7 Hz), 7.28 (2H, m), 7.99 (1H, d, J = 8.3 Hz); 402 (ESMS, MH⁺).

Example 85

N-{[4-(4,5-Dihydrobenzo[2,3]oxepino[4,5-d][1,3]-thiazol-2-ylamino)cyclohexyl)methyl}-N-propylformamide:

N1-{[4-(4,5-Dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-ylamino)cyclohexyl)methyl}-propanamide: To a stirred solution of N2-[4-(aminomethyl)cyclohexyl]-4,5-dihydrobenzo[2,3]-oxepino[4,5-d][1,3]thiazol-2-amine (0.600 g, 1.49 mmol) in anhydrous pyridine (3 ml) and diisopropylethylamine (3 ml) was added propionyl chloride (0.144 ml, 1.66 mmol) dropwise. The solution was stirred

143

for 20 minutes at room temperature and then poured into water (50 ml) and extracted with EtOAc. The solvent was removed and the residue chromatographed on silica-gel eluting with EtOAc to yield 70% (402 mg) of the product as a light yellow solid: $^1\text{H-NMR}$ (CDCl_3) δ 1.12 (6H, m), 1.50 (1H, m), 1.65 (1H, m), 1.85 (2H, d, $J = 12.1$ Hz), 2.22 (4H, m), 3.16 (4H, t, $J = 5.3$ Hz), 3.32 (1H, m), 4.33 (2H, t, $J = 5.2$ Hz), 4.81 (1H, m), 5.49 (1H, m), 7.00 (1H, d, $J = 8.5$ Hz), 7.15 (2H, m), 8.25 (1H, d, $J = 8.1$ Hz). m.p. 56-60°C. FAB+ $m/z = 386$ (MH^+).

N2-{4-[(Propylamino)methyl]cyclohexyl}-4,5-dihydrobenzo-[2,3]oxepino[4,5-d][1,3]thiazol-2-amine: To a solution of N1-{[4-(4,5-dihydrobenzo[2,3]-oxepino[4,5-d][1,3]thiazol-2-ylamino)cyclohexyl]-methyl}propanamide (0.380 g, 0.944 mmol) in anhydrous THF (7 ml) was added 1M BH_3 in THF (2.0 ml, 2.0 mmol). The solution was stirred for several hours at room temperature. TLC (EtOAc) indicated reaction to be incomplete and an additional 2.5 ml 1M BH_3 in THF was added to the solution which was then stirred overnight. 4N HCl (25 ml) was slowly added to the stirred solution and the solution was washed with EtOAc. The aqueous layer was made basic by addition of 20% NaOH. The resulting solution was extracted with EtOAc and the extract washed with water. Upon removal of the solvent, 72% (254 mg) of the product was obtained as a white solid: $^1\text{H-NMR}$ (CDCl_3) δ 0.88 (3H, t, $J = 7.3$ Hz), 1.12 (4H, m), 1.43 (3H, m), 1.75 (2H, m), 2.14 (2H, m), 2.41 (2H, d, $J = 6.6$ Hz), 2.51 (2H, t, $J = 7.4$ Hz), 2.60 (1H, m), 3.10 (2H, t, $J = 5.0$ Hz), 3.22 (1H, m), 4.27 (2H, t, $J = 5.0$ Hz), 5.35 (1H, m), 6.95 (1H, d, $J = 7.6$ Hz), 7.06 (2H, m), 8.20 (1H, d, $J = 7.3$ Hz); 372 (ESMS, MH^+).

N-{[4-(4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-ylamino)cyclohexyl]methyl}-N-propylformamide: To a solution of N2-{4-[(propyl-amino)methyl]cyclohexyl}-4,5-dihydrobenzo[2,3]-oxepino[4,5-d][1,3]thiazol-2-amine (0.254 g, 0.68 mmol) in anhydrous THF (4 ml) was added 1H-benzotriazole-1-carboxaldehyde (0.110 g, 0.75 mmol). The solution was stirred at room temperature for 1 hour and then poured into 2N NaOH (50 ml) and extracted with EtOAc (60 ml). The extract was separated and evaporated to yield 295 mg of a viscous oil. Chromatography on silica-gel eluting with 1:1 EtOAc/hexane yielded 56% (152 mg) of the product as a white solid: ¹H-NMR (CDCl₃) δ 0.90 (3H, m), 1.14 (4H, m), 1.58 (2H, m), 1.78 (2H, m), 2.23 (2H, m), 3.20 (8H, m), 4.32 (2H, t, J = 4.9 Hz), 5.03 (1H, m), 6.99 (1H, m), 7.12 (2H, m), 8.01 (0.5H, s), 8.11 (0.5H, s), 8.22 (1H, d); m.p. 53-55 °C; 400 (ESMS, MH⁺).

Example 86

N-{4-[(4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-ylamino)methyl]cyclohexyl}-N-(2-methoxyethyl)formamide

N2-({4-[(2-methoxyethyl)amino]cyclohexyl}methyl)-4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-amine: To a stirred solution of N1-{4-[(4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-ylamino)-methyl]cyclohexyl}-2-methoxyacetamide (free-base) (0.450g, 1.12 mmol) in anhydrous THF (5 ml) was added 5.0 ml 1M BH₃ in THF (5.0 mmol). The solution was stirred at room temperature for 26 hours. 4N HCl (12 ml) was slowly added to the stirred solution and stirred continued for 1 hour. The solution was made basic with 2N NaOH (pH~9) and was extracted with EtOAc. Upon evaporation of the solvent,

145

90% (390 mg) of the desired amine was obtained as a highly viscous colorless oil. The compound was converted to its dihydrochloride salt in chloroform with 1N HCl in Et₂O. Evaporation of the solvent gave 460 mg of the salt as a slightly hygroscopic white solid: m.p. 243-245°C (dec); ¹H-NMR (CDCl₃) δ 1.08 (4H, t, J = 10.8 Hz), 1.6 (2H, broad), 1.91 (4H, m), 2.41 (1H, broad), 2.80 (2H, t, J = 4.9 Hz), 3.15 (4H, m), 3.35 (3H, s), 3.49 (2H, t, J = 5.3 Hz), 4.32 (2H, t, J = 5.4 Hz), 5.13 (1H, broad t), 6.98 (1H, dd, J = 1.3 Hz, 7.7 Hz), 7.12 (2H, m), 8.24 (1H, dd, J = 1.6 Hz, 7.9 Hz); 388 (ESMS, MH⁺).

N-{4-[(4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]-thiazol-2-ylamino)methyl]cyclohexyl}-N-(2-methoxy-ethyl)formamide:
To a stirred solution of N2-({4-[(2-methoxyethyl)amino]cyclohexyl)methyl}-4,5-dihydro-benzo-[2,3]oxepino[4,5-d][1,3]thiazol-2-amine (free-base) (0.356 g, 0.92 mmol) in anhydrous THF (4 ml) was added 1H-benzotriazole-1-carboxaldehyde (0.144 g, 0.98 mmol). The solution was stirred for 5 minutes at room temperature and TLC (EtOAc) indicated reaction to be complete. The solution was poured into 2N NaOH (50 ml) and extracted with EtOAc (60 ml). The extract was evaporated to yield 420 mg of the crude product as a white solid. The product was chromatographed on silica-gel eluting with 3:1 EtOAc/hexane to yield 78% (300 mg) of a white solid. The material was converted to its HCl salt in chloroform with 1M HCl in Et₂O. Upon evaporation of the solvent, a white hygroscopic solid was obtained. The solid was triturated with hexane, filtered and dried to yield 330 mg of a non-hygroscopic free flowing white solid: m.p. 164-166°C; ¹H-NMR (CDCl₃) δ 1.11 (2H, p), 1.53 (3H, m), 1.90 (5H, m), 3.14 (4H, m), 3.32 (3H, s), 3.43 (4H, m), 4.34 (2H, t),

5.43 (1H, broad), 7.00 (1H, ¹⁴⁶d), 7.12 (2H, m), 8.08 (0.4H, s), 8.18 (0.6H, s), 8.24 (1H, dd); 416 (ESMS, MH⁺).

Example 87

5

trans-1-Aza-2-(4-(n-(ethyl)formamido)cyclohexyl)methyl-amino-4,5-dihydro-6-oxa-3-thia-benzo[e]azulene:

10

trans-2-(4-Acetamido)cyclohexylmethylamino-1-aza-4,5-dihydro-6-oxa-3-thia-benzo[e]azulene:

15

Benzyl-N-[4-({[(benzoylamino)carbothioyl]amino}-methyl)-cyclohexyl]carbamate: A solution of benzyl N-[4-(aminomethyl)cyclohexyl]carbamate HCl (3.46 g, 11.58 mmol) and diisopropylethylamine (6.0 ml) in anhydrous THF (90 ml) was stirred at room temperature for 20 minutes. To this solution was added benzoylisothiocyanate (2.00 g, 12.3 mmol). The solution was stirred at room temperature for 14 hours, poured into water (200 ml) and then extracted with Et₂O (2X150 ml). The organic layer was dried over MgSO₄, filtered and evaporated to yield a solid. The solid was triturated with hexane and filtered to yield 99% (5.023g) of the desired product.

20

25

Benzyl-N-(4-{[(aminocarbothioyl)amino]methyl}-cyclohexyl)-carbamate: To a solution of benzyl N-[4-({[(benzoylamino)carbothioyl]amino}methyl)cyclohexyl]carbamate (5.02 g, 11.4 mmol) in MeOH (110 ml) was added a solution of K₂CO₃ (3.5 g, 25.3 mmol) in water (30 ml). The solution was stirred for 24 hours at room temperature. The solvent was removed in vacuo with repeated additions of MeOH to azeotropically remove water. The resulting solid was treated with acetone (60 ml) and filtered. The filtrate was evaporated to yield 95% (3.47 g) of a yellow solid.

30

35

Benzyl-N-{4-[(4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]-thiazol-2-ylamino)methyl]cyclohexyl}-carbamate:

A solution containing benzyl N-(4-
5 {[(aminocarbothioyl)amino]methyl}-cyclohexyl)-carbamate
(3.45 g, 10.73 mmol), 4-bromo-2,3,4,5-tetrahydro-1-benzoxepin-5-one (2.60 g, 10.78 mmol), and diisopropylethylamine (2.50 ml) in EtOH (50 ml) was heated at reflux for 4 hours. The solvent was removed and the
10 residue was chromatographed on silica gel eluting with 1:3 EtOAc/hexane to yield 65% (3.2 g) of the product as a white solid: m.p. 124-127°C; ¹H-NMR (CDCl₃) δ 1.09 (4H, pentet, J = 10.4 Hz), 1.55 (1H, broad), 1.83 (2H, m), 2.04 (2H, broad), 3.10 (2H, t, J = 6.2 Hz), 3.15 (2H, t, J =
15 5.2 Hz), 3.43 (1H, broad) 4.33 (2H, t, J = 5.2 Hz), 4.62 (1H, d, J = 8.0 Hz), 5.08 (2H, s), 5.31 (1H, m), 7.00 (1H, dd, J = 1.2 Hz, 7.8 Hz), 7.12 (2H, m), 7.35 (5H, s), 8.25 (1H, dd, J = 1.7 Hz, 7.8 Hz). ESIMS m/z = 464 (MH⁺).

20 N2-[(4-aminocyclohexyl)methyl]-4,5-dihydrobenzo[2,3]-oxepino[4,5-d][1,3]thiazol-2-amine: A solution of benzyl-N-{4-[(4,5-dihydrobenzo-[2,3]oxepino-[4,5-d][1,3]-thiazol-2-ylamino)methyl]cyclohexyl}carbamate (3.00 g, 6.47 mmol) in glacial acetic acid (50 ml) and conc. HCl (20 ml) was
25 heated to a gentle reflux for 45 minutes. The solvent was removed on the rotor-vap with gentle heating and the residue diluted with water (150 ml). The resulting solution was made basic with NaOH (pH~11). EtOAc (200 ml) was added and the solution was shaken vigorously. The
30 phases were separated and the aqueous phase was extracted with EtOAc. The combined extract was washed with water and evaporated to yield 2.30 g of a brown viscous oil. The oil was taken up into 40 ml EtOAc and 5 ml 4N HCl in dioxane was added. The resulting dihydrochloride salt was

148

obtained by filtration and washed with EtOAc and ether. Upon drying, 100% (2.6 g) of the salt was obtained as a tan solid: $^1\text{H-NMR}$ ($\text{d}_4\text{-MeOH}$) δ 1.21 (2H, m), 1.43 (2H, m), 1.79 (1H, m), 1.99 (2H, m), 2.10 (2H, m), 3.09 (1H, m), 3.18 (2H, t, $J = 5.0$ Hz), 3.35 (2H, d, $J = 6.7$ Hz), 4.32 (2H, t, $J = 5.3$ Hz), 7.12 (1H, dd, $J = 1.1, 8.0$ Hz), 7.22 (1H, m), 7.33 (1H, m), 7.70 (1H, dd, $J = 1.3, 8.0$ Hz). ESIMS $m/z = 330$ (MH^+).

10 N1-{4-[(4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]-thiazol-2-ylamino)methyl]cyclohexyl}acetamide: To a stirred solution of N2-[(4-aminocyclohexyl)methyl]-4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-amine (0.600 g, 1.49 mmol) in anhydrous pyridine (3 ml) and diisopropylethylamine was added acetyl chloride (0.130 ml, 1.83 mmol) dropwise. The solution was stirred for 30 minutes at room temperature and then poured into water (60 ml). The mixture was extracted with EtOAc, and the organic phase evaporated to afford a residue that was chromatographed on silica-gel eluting with EtOAc to yield 20 72% (400 mg) of the desired product as a light yellow solid: m.p. 69-72°C. $^1\text{H-NMR}$ (CDCl_3) δ 1.11 (4H, m), 1.60 (1H, broad), 1.73 (1H, broad), 1.83 (2H, m), 1.95 (3H, s), 2.05 (2H, m), 3.14 (4H, m), 3.73 (1H, broad), 4.33 (2H, t, $J = 4.9$ Hz), 5.30 (1H, broad), 7.00 (1H, dd, $J = 0.9$ Hz, 7.7 Hz), 7.12 (2H, m), 8.24 (1H, dd, $J = 0.9$ Hz, 7.8 Hz). ESIMS $m/z = 372$ (MH^+).

30 N2-{[4-(Ethylamino)cyclohexyl]methyl}-4,5-dihydro-benzo-[2,3]oxepino[4,5-d][1,3]thiazol-2-amine: To a stirred solution of N1-{4-[(4,5-dihydro-benzo[2,3]oxepino[4,5-d][1,3]thiazol-2-ylamino)-methyl]cyclohexyl}acetamide (0.380 g, 1.02 mmol) in anhydrous THF (4 ml) was added 1M BH_3 in THF (5.0 ml, 5.0 mmol). The solution was stirred at

149

room temperature overnight. 4N HCl (12 ml) was added slowly and stirring continued for 15 minutes. The resulting solution was made basic (pH-9) with 2N NaOH. The cloudy solution was extracted with EtOAc and the organic layer washed with water. Removal of the solvent in vacuo, gave 69% (252 mg) of the product as a solid which was used without further purification: $^1\text{H-NMR}$ (CDCl_3) δ 0.96 (4H, m), 1.08 (3H, t, $J = 7.1$ Hz), 1.47 (1H, m), 1.76 (2H, d, $J = 12.1$ Hz), 1.90 (2H, m), 2.37 (2H, m), 2.63 (2H, q, $J = 7.0$ Hz), 3.05 (2H, t, $J = 5.7$ Hz), 3.11 (2H, t, $J = 5.1$ Hz), 4.28 (2H, t, $J = 5.2$ Hz), 5.73 (1H, broad t), 6.96 (1H, dd, $J = 0.9$ Hz, 7.7 Hz), 7.10 (2H, m), 8.22 (1H, dd, $J = 1.3$ Hz, 7.6 Hz); 358 (ESMS, MH^+).

N-{4-[(4,5-Dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-ylamino)methyl]cyclohexyl}-N-ethylformamide: To a stirred solution of N2-{[4-(ethylamino)cyclohexyl]methyl}-4,5-dihydrobenzo[2,3]-oxepino-[4,5-d][1,3]-thiazol-2-amine (0.252 g, 0.70 mmol) in anhydrous THF (3.5 ml) was added 1H-benzotriazole-1-carboxaldehyde (0.112 g, 0.76 mmol). The solution was stirred at room temperature for 1 hour and then poured into 2N NaOH (60 ml). The mixture was extracted with EtOAc and evaporated to afford 268 mg of the desired crude product as a solid. The product was chromatographed on silica-gel eluting with 70% EtOAc/30% hexane to yield 56% (150 mg) of a white solid. The compound was dissolved in chloroform and converted to its HCl salt by addition of 1N HCl in ether. The solvents were removed in vacuo and the residue triturated with hexane/ether and filtered to afford the product as a white solid: m.p. 210-213°C; $^1\text{H-NMR}$ (CDCl_3) δ 1.13 (4H, m), 1.59 (2H, m), 1.88 (6H, m), 3.17 (4H, m), 3.24 (3H, m), 4.32 (2H, t), 5.40 (1H, broad), 7.00 (1H, d), 7.13 (2H,

150

m), 8.09 (0.4H, s), 8.12 (0.6H, s), 8.23 (1H, dd);
386 (ESMS, MH⁺).

Example 88

5

N-{4-[(4,5-Dihydrobenzo[2,3]oxepino[4,5-d][1,3]-thiazol-2-ylamino)methyl]cyclohexyl}-N-propyl-formamide:

10

N1-{4-[(4,5-Dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-ylamino)methyl]cyclohexyl}-propanamide: To a stirred solution of N2-{(4-aminocyclohexyl)methyl}-4,5-dihydrobenzo[2,3]oxepino-[4,5-d][1,3]thiazol-2-amine (0.600 g, 1.49 mmol) in anhydrous pyridine (3 ml) and diisopropylethylamine (3 ml) was added propionyl chloride (0.155 ml, 1.78 mmol) dropwise. The solution was stirred for 1 hour at room temperature and poured into water (60 ml). The solution was extracted with EtOAc and the solvent removed in vacuo to yield a viscous golden oil. The oil was chromatographed on silica-gel eluting with 70% EtOAc/30% hexane to yield 74% (425 mg) of an orange solid: m.p. 70-73°C. ¹H-NMR (CDCl₃) δ 1.10 (6H, m), 1.60 (1H, broad), 1.83 (3H, m), 2.00 (3H, m), 2.17 (2H, q, J = 7.6 Hz), 3.13 (4H, q, J = 5.1 Hz), 3.73 (1H, broad), 4.32 (2H, t, J = 5.3 Hz), 5.32 (1H, broad), 7.00 (1H, dd, J = 6.4 Hz). ESIMS m/z = 386 (MH⁺).

25

N2-{[4-(propylamino)cyclohexyl]methyl}-4,5-dihydrobenzo-[2,3]oxepino[4,5-d][1,3]thiazol-2-amine: To a stirred solution of N1-{4-[(4,5-dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-ylamino)methyl]cyclohexyl}propanamide (0.405 g, 1.05 mmol) in anhydrous THF (4 ml) was added 1M BH₃ in THF (5.5 ml, 5.5 mmol). The solution was stirred at room temperature for 52 hours. 4N HCl (12 ml) was added slowly and stirring continued for 30 minutes thereafter.

30

151

The solution was poured into 2N NaOH (60 ml) and extracted with EtOAc. The extract was washed with 10% brine, separated and evaporated to yield 77% (300 mg) of the amine as a viscous golden oil: $^1\text{H-NMR}$ (CDCl_3) δ 0.88 (3H, t), 1.06 (2H, m), 1.38 (2H, m), 1.49 (2H, m), 1.80 (6H, m), 2.40 (1H, m), 2.58 (2H, t), 3.15 (2H, m), 3.63 (2H, t), 4.30 (2H, t), 5.12 (1H, m), 6.98 (1H, d), 7.11 (2H, m), 8.23 (1H, dd); 400 (ESMS, MH^+).

10 N-{4-[(4,5-Dihydrobenzo[2,3]oxepino[4,5-d][1,3]thiazol-2-ylamino)methyl]cyclohexyl}-N-propylformamide: To a stirred solution of N2-{[4-(propylamino)cyclohexyl]methyl}-4,5-dihydrobenzo-[2,3]-oxepino[4,5-d][1,3]thiazol-2-amine (0.300 g, 0.81 mmol) in
15 anhydrous THF (4 ml) was added 1H-benzotriazole-1-carboxaldehyde (0.126 g, 0.86 mmol). The solution was stirred for 1 hour at room temperature and poured into 2N NaOH (50 ml). The mixture was extracted with EtOAc (60 ml). The extract was evaporated to yield 320 mg of the
20 crude product as a viscous yellow oil. The oil was chromatographed on silica-gel eluting with 70% EtOAc/30% hexane to afford 125 mg of the product as a colorless oil. The oil was dissolved in chloroform and converted to its HCl salt with 1N HCl in Et_2O . Upon evaporation of the
25 solvent, a solid was obtained. The solid was triturated with EtOAc, filtered and washed with hexane to give 25% (87mg) of the desired product as a white solid: m.p. (HCl salt) 195-197°C; $^1\text{H-NMR}$ (CDCl_3) δ 0.89 (3H, t), 1.05 (2H, p), 1.50 (5H, m), 1.79 (4H, m), 3.10 (6H, m), 4.30 (2H, m), 5.93 (2H, m), 7.06 (3H, m), 8.02 (0.4H, s), 8.10
30 (0.6H, s), 8.22 (1H, dd); 400 (ESMS, MH^+).

Example 89

N1-{4-[(9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-
[d][1,3]thiazol-2-yl)amino]benzyl}-2-methoxyacetamide: To
5 a stirred solution of N2-[4-(aminomethyl)phenyl]-9-fluoro-
5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine
(See example 90) (1.50 g, 3.64 mmol) in anhydrous pyridine
(10 ml) and diisopropylethylamine (5.0 ml) was added
methoxyacetyl chloride (0.40 ml, 4.38 mmol) dropwise. The
10 solution was stirred overnight, poured into water (200 ml)
and extracted with EtOAc (200 ml). The extract was
evaporated to yield 1.8 g of a viscous golden oil. The
oil was chromatographed on silica-gel eluting with EtOAc
to yield 80% (1.2 g) of the desired product as a foam.
15 The product was converted to its HCl salt in chloroform
using 1M HCl in Et₂O: m.p. (HCl salt) 175-178 °C; ¹H-NMR
(CDCl₃) δ 2.15 (2H, m), 2.86 (4H, m), 3.42 (3H, s), 3.96
(2H, s), 4.50 (2H, d, J = 6.1 Hz), 7.00 (1H, t), 7.18-7.40
(7H, m), 7.61 (1H, dd, J = 2.5 Hz, 9.5 Hz); 412 (ESMS,
20 MH⁺).

Example 90

N-{4-[(9-Fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta-
25 [d][1,3]thiazol-2-yl)amino]benzyl}methanesulfonamide:

N2-[4-(Aminomethyl)phenyl]-9-fluoro-5,6-dihydro-4H-
benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine:

30 6-Bromo-3-fluoro-6,7,8,9-tetrahydro-5H-
benzo[a]cyclohepten-5-one: To a stirred solution of 8-
fluoro-1-benzosuberone (20.00g, 112.2 mmol) in glacial
acetic acid (125 ml) was added bromine (5.8 ml, 112.6
mmol) dropwise. Once the addition was complete, the

153

solution was poured into water (400 ml) and extracted with EtOAc (500 ml). The extract was separated and washed with water (500 ml). The organic layer was dried over MgSO_4 , filtered and evaporated to yield 100% (28.9 g) of the desired product as a viscous oil: ^1H -NMR (CDCl_3) δ 2.00 (2H, m), 2.33 (2H, m), 2.84 (1H, m), 3.00 (1H, m), 4.83 (1H, m), 7.13 (2H, m), 7.29 (1H, dd).

tert-Butyl N-(4-Aminobenzyl)carbamate: To a stirred solution of 4-amino-benzylamine (7.5 g, 61.39 mmol) in anhydrous THF (100 ml) under N_2 was added di-t-butyl dicarbonate (13.5 g, 61.9 mmol) in portions. The solution was stirred at room temperature under N_2 for 90 minutes. The solvent was removed in vacuo to yield 100% (13.7 g) of the Boc-protected amine as a white solid: m.p. 70-72 °C; ^1H -NMR (CDCl_3) δ 1.43 (9H, s), 3.62 (2H, broad), 4.18 (2H, d), 4.77 (1H, broad), 6.63 (2H, d), 7.06 (2H, d).

tert-Butyl N-(4-{[(Benzoylamino)carbothioyl]amino}benzyl)-carbamate: To a stirred solution of tert-butyl N-(4-aminobenzyl)carbamate (6.00 g, 27.0 mmol) in anhydrous THF (50 ml) was added benzoylisothiocyanate (4.43 g, 27.0 mmol). The solution was stirred for 2 hours and the solvent removed in vacuo to yield 100% (10.4 g) of the protected thiourea as a yellow solid: m.p. 161-163 °C; ^1H -NMR (CDCl_3) δ 1.45 (9H, s), 4.33 (2H, d), 4.85 (1H, broad), 7.34 (2H, d), 7.60 (5H, m), 7.90 (2H, d), 9.13 (1H, s), 12.58 (1H, s); 386 (ESMS, MH^+).

tert-Butyl N-{4-[(Aminocarbothioyl)amino]benzyl}-carbamate: To a stirred near solution of tert-butyl N-(4-{[(benzoylamino)carbothioyl]amino}benzyl)-carbamate (10.4 g, 27.0 mmol) in MeOH (150 ml) was added a solution of K_2CO_3 (8.5 g, 61.5 mmol) in 15 ml water. The solution was

154

stirred at room temperature for 24 hours and filtered to remove a white precipitate. The filtrate was evaporated in vacuo to yield 100% (7.6 g) of the desired product as an off-white solid.

5

tert-Butyl N-{4-[(9-Fluoro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[d][1,3]thiazol-2-yl)amino]benzyl}-carbamate: A solution containing tert-butyl N-{4-[(aminocarbothioyl)-amino]benzyl}carbamate (7.60 g, 27.0 mmol), 6-bromo-3-fluoro-6,7,8,9-tetrahydro-5H-benzo[a]cyclohepten-5-one (6.95 g, 27.0 mmol) and diisopropylethylamine (5.0 ml) in EtOH (60 ml) was heated at reflux for 2.5 hours. The solvent was removed in vacuo and the residue dissolved in EtOAc (125 ml). The resulting solution was filtered to remove the insoluble (i-pr)₂NEt•HBr. The filtrate was transferred to a 250 ml round bottom flask and 4N HCl in dioxane (35 ml) was added with stirring. The solution was stirred for 90 minutes and then filtered to collect the hydrochloride salt. The solid was washed with EtOAc and then with Et₂O. Upon drying, 86% (11.1 g) of the Boc-protected amine hydrochloride was obtained as a pale yellow solid: ¹H-NMR (d₄-MeOH) δ 1.43 (9H, s), 2.18 (2H, m), 2.80 (4H, m), 4.25 (2H, s), 7.12 (1H, dt), 7.33 (2H, m), 7.43 (4H, s).

25

N2-[4-(aminomethyl)phenyl]-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]thiazol-2-amine: To a stirred near solution of tert-butyl [4-(4H-chromeno[4,3-d][1,3]thiazol-2-ylamino)cyclohexyl]-methylcarbamate (11.1 g, 23.3 mmol) in MeOH (150 ml) was added 4N HCl in dioxane (40 ml). The solution was brought to a brief gentle reflux and the solvent then removed in vacuo. The resulting solid was re-suspended in Et₂O, filtered and dried to yield 90% (9.25 g) of the desired di-

30

155

hydrochloride salt as a white solid: m.p. 236-239 °C; $^1\text{H-NMR}$ (CDCl_3) δ 1.75 (2H, broad), 2.06 (2H, m), 2.77 (2H, m), 2.83 (2H, t), 3.79 (2H, s), 6.82 (1H, t), 7.04 (1H, m), 7.22 (4H, m), 7.79 (1H, dd), 8.8 (1H, broad); 340 (ESMS, MH^+).

To a stirred solution of N2-[4-(aminomethyl)phenyl]-9-fluoro-5,6-dihydro-4H-benzo[6,7]cyclohepta[d][1,3]-thiazol-2-amine (1.50 g, 3.64 mmol) in anhydrous pyridine (10 ml) and diisopropylethylamine (5.0 ml) was added methanesulfonyl chloride (0.38 ml, 4.91 mmol) dropwise. The solution was stirred at room temperature overnight, poured into water (250 ml) and extracted with EtOAc. The extract was evaporated to yield 1.7 g of the crude product as a viscous orange oil. The oil was chromatographed on silica-gel eluting with EtOAc (R_f of product ~ 0.9) to yield 66% (1.0 g) of the desired product as a solid. The solid was triturated with Et_2O /hexane (1:1) and filtered to give 0.93 g of the product as a pale yellow solid: m.p. 148-150 °C; $^1\text{H-NMR}$ (CDCl_3) δ 2.13 (2H, m), 2.79 (2H, m), 2.89 (3H, s), 2.91 (2H, m), 4.24 (2H, d, $J = 4.3$ Hz), 5.06 (1H, m), 6.88 (1H, t), 7.13 (1H, t), 7.30 (4H, m), 7.48 (1H, m), 7.72 (1H, dd, $J = 2.7$ Hz, 10.6 Hz); 418 (ESMS, MH^+).

I. Synthetic Methods for ExamplesB. Bicyclic Compounds5 **General Procedures Relating to Examples:**

For the formation of 2-aminothiazoles from 2-haloketones and thioureas, see, for example, Kearney, P.C., et al., 1998; Di Fabio, R. and Pentassuglia, G., 1998; De Kimpe, N., et al., 1996; Plazzi, P.V., et al., 1995; and
10 Novikova, A. P., 1991.

For the formation of thiazoles from 2-haloketones and thioamides, see, for example, Critcher, D. J. and Pattenden, G., 1996; and Friedman, B. S., et al., 1937.

15 For the formation of 2-aminoimidazoles from 2-haloketones and guanidines, see, for example, Little, T. L. and Webber, 1994; and Chabaka, L.M., et al., 1994.

20 For the formation of imidazoles from 2-haloketones and amidines, see, for example, Demchenko, A. M., et al., 1997; and Nagao, Y., et al., 1996.

25 For the synthesis of 2-aminooxazoles from 2-haloketones and ureas, see, for example, Pathak, V.N., et al., 1993; Crangk, G. and Foulis, M.J., 1971; and Marchetti, E., et al., 1968.

30 For the formation of oxazoles from 2-haloketones and amides, see, for example, Hammar, W.J. and Rustad, M.A., 1981; and Zhao, Z., et al., 1991.

157

Benzotriazole-1- carboxaldehyde was purchased from Aldrich Chemical Company and is recommended for the formation of formamides from amines.

5 All reactions were performed under an inert atmosphere (Argon) and the reagents, neat or in appropriate solvents, were transferred to the reaction vessel via syringe and cannula techniques. Anhydrous solvents were purchased from Aldrich Chemical Company and used as received. The
10 examples 1-9 described in the application were named using ACD/Name program (version 2.51, Advanced Chemistry Development Inc., Toronto, Ontario, M5H2L3, Canada).

¹H and ¹³C spectra were recorded at 300 and 75 MHz (QE
15 Plus) with CDCl₃ as solvent (unless otherwise noted) and tetramethylsilane as internal standard. s = singlet; d = doublet; t = triplet; q = quartet; p = pentet; sextet; septet; b = broad; m = multiplet. Elemental analyses were performed by Robertson Microlit Laboratories, Inc. Low-
20 resolution electrospray MS spectra were measured (ESMS, MS) and MH⁺ is reported. Thin-layer chromatography (TLC) was carried out on glass plates precoated with silica gel 60 F₂₅₄ (0.25 mm, EM Separations Tech.). Preparative thin-
25 layer chromatography was carried out on glass sheets precoated with silica gel GF (2 mm, Analtech). Flash column chromatography was performed on Merck silica gel 60 (230 - 400 mesh). Melting points were determined in open capillary tubes on a Med-Temp apparatus and are uncorrected.

30

General Procedure for the Synthesis of Bromoketones:

To a cooled solution of the ketone (1 equivalent) in acetic acid or an appropriate solvent, cooled in a water bath, was slowly added bromine or a brominating agent such as tetrabutylammonium perbromide (1 equivalent). The reaction mixture was stirred at room temperature. The solvent was evaporated and the residue was dissolved in dichloromethane, and washed with saturated sodium bicarbonate and water. The organic phase was dried over sodium sulfate. Evaporation of the combined decolorized organic phase afforded a light yellow oil. In some cases, the desired product precipitated upon concentration of the reaction mixture. Many of the bromoketones used in this application were commercially available.

General Procedure for the Synthesis of Thioureas:

A protected diamine such as N-Boc-1,4-diaminobutane or N-Boc-1,5-diaminopentane (1 equivalent) was dissolved in tetrahydrofuran and stirred at room temperature. Benzoyl isothiocyanate (1.1 equivalent) was added dropwise to the aforementioned solution. The resulting mixture was stirred at room temperature for 24 hours. The solvent was removed under reduced pressure to give a yellow oil. The yellow oil (1 equivalent) was dissolved in methanol, an aqueous potassium carbonate (3 equivalents) solution was added, and the mixture stirred for 48 hours. Water was added to the reaction mixture and extracted with 2x75 ml ethyl acetate. The combined ethyl acetate extracts were washed with water, dried with anhydrous sodium sulfate, filtered and the solvent removed under reduced pressure to give the desired thiourea.

The following procedure is typical:

a. ***t*-Butyl-*trans*-*N*-(4-(benzoylaminocarbothioylamino)-cyclohexyl)methyl-carbamate:** A solution of *t*-butyl *trans*-*N*-(4-aminocyclohexyl)methylcarbamate (4.68 g, 20.5 mmol) and benzoylisothiocyanate (3.77 g, 23.1 mmol) in THF (180 ml) was stirred at room temperature for 20 hours. The solvent was removed *in vacuo* to yield a golden viscous oil. The oil was triturated with hexanes (250 ml). The resulting solid was collected by filtration and washed with hexanes to yield 91% (7.32 g) of the product as an off-white solid: ^1H NMR (CDCl_3) δ 1.20 (6H, m), 1.44 (9H, s), 1.83 (2H, m), 2.25 (2H, m), 3.01 (2H, t, $J = 6.4$ Hz), 4.20 (1H, m), 4.62 (1H, m), 7.51 (2H, t, $J = 7.9$ Hz), 7.60 (1H, t, $J = 7.4$ Hz), 7.81 (2H, d, $J = 7.2$ Hz), 8.93 (1H, s); ESMS $m/e = 392$ (MH^+).

b. ***t*-Butyl *trans*-*N*-(4-(aminocarbothioylamino)-cyclohexyl)-methylcarbamate:** To a solution of *t*-butyl *trans*-*N*-(4-(benzoylaminocarbothioylamino)-cyclohexyl)methyl-carbamate (7.30 g, 18.6 mmol) in MeOH (110 ml) was added an aqueous solution of potassium carbonate (5.15 g, 37.3 mmol in 50 ml). The reaction mixture was stirred at room temperature for 18 hours. The solvent was removed *in vacuo* and the residue was extracted several times with EtOH. The combined EtOH extracts were evaporated to dryness to yield 6.15 g of an off-white solid. The solid was dissolved in acetone (200 ml) and stirred for 20 minutes. The solution was filtered by suction to remove any insoluble salts and the filtrate was evaporated to yield 98% (5.23 g) of the desired product as a slightly yellow solid: ^1H NMR (CDCl_3) δ 1.10 (6H, m), 1.43 (9H, s), 1.80 (2H, m), 2.10 (2H, m), 2.95 (2H, t, $J =$

6.4 Hz), 4.75 (1H, broad), 6.03 (2H, s), 6.64 (1H, broad); ESMS m/e = 288 (MH⁺).

tert-Butyl 5-[(aminocarbothioyl)amino]pentylcarbamate was obtained as a light yellow wax from tert-butyl 5-
5 {[(benzoylamino)carbothioyl]amino}-pentylcarbamate: ¹H NMR (CD₃OD) δ 3.44 (m, 1H), 3.10 (m, 1H), 3.01 (t, 2H, J = 6.7 Hz), 1.60-1.31 (m, 6H), 1.41 (s, 9H); 262 (ESMS, MH⁺).

tert-Butyl-5-{[(benzoylamino)carbothioyl]amino}-pentyl-
10 carbamate was obtained as a light yellow solid in 79% yield from N-BOC-1,5-diaminopentane and benzoyl isothiocyanate: m.p. 90-93 °C.

trans-tert-Butyl-{4-[(aminocarbothioyl)amino]-
15 cyclohexyl}methylcarbamate was obtained as a light yellow wax from trans-tert-butyl-(4-{[(benzoylamino)-carbothioyl]amino}-cyclohexyl)-methylcarbamate: ¹H NMR (CD₃OD) δ 3.92 (m, 1H), 2.86 (m, 2H), 2.00 (m, 2H), 1.76 (m, 2H), 1.41 (s, 9H), 1.37 (m, 1H), 1.06 (m, 4H); 288
20 (ESMS, MH⁺).

trans-tert-Butyl-(4-{[(benzoylamino)carbothioyl]-
25 amino}cyclohexyl)methylcarbamate was obtained as a yellow solid in 97% yield from tert-butyl 4-aminocyclohexylmethylcarbamate and benzoyl isothiocyanate.

trans-tert-Butyl 4-aminocyclohexylmethylcarbamate was
30 obtained in more than 95% yield by hydrogenation of benzyl 4-{[(tert-butoxycarbonyl)amino]methyl}-cyclocarbamate.

Benzyl-4-[[[tert-

butoxycarbonyl]amino]methyl]

cyclohexylcarbamate: To a stirred suspension of

4-[[[(tert-butoxycarbonyl)amino]methyl]

5 cyclohexanecarboxylic acid (Maybridge Chemical Co., Ltd.)
(45g) and diphenylphosphoryl azide (44 ml) in toluene (600
ml) was added triethylamine (32 ml) over a period of 20
min whilst maintaining the internal temperature at -10-
0 C. The mixture was slowly warmed and then stirred at
10 70 C for 4h. After cooling to 40 C, benzyl alcohol (36
ml) was added and the reaction mixture heated at reflux
for 20 h. The cold reaction mixture was washed with water
and brine and dried over anhydrous magnesium sulfate.
Removal of the solvent and recrystallization of the
15 organic residue from ethyl acetate and diethyl ether gave
the title compound, benzyl-4-[[[tert-butoxycarbonyl]
amino]methyl]cyclohexylcarbamate as a white solid, m.p.
129-131 C.

20 *trans*-Benzyl-4-{[(aminocarbothioyl)amino]methyl}-
cyclohexyl-carbamate was obtained as a yellow solid in 71%
yield from *trans*-benzyl 4-
({[(benzoylamino)carbothioyl]amino}methyl)-
cyclohexylcarbamate: 322 (ESMS, MH⁺).

25 *trans*-Benzyl-4-({[(benzoylamino)carbothioyl]amino}-
methyl)-cyclohexylcarbamate was obtained as a yellow solid
from benzyl 4-(aminomethyl)cyclohexylcarbamate and benzoyl
isothiocyanate.

30 *trans*-Benzyl 4-(aminomethyl)cyclohexylcarbamate was
obtained as a white solid in more than 95% yield by
stirring benzyl 4-{[(tert-butoxycarbonyl)amino]-methyl}-

162

cyclocarbamate in 2N HCl (made from 1 : 1 of EtOAc and 4N HCl in dioxane).

General Procedure for the Synthesis of Bicyclic Thiazoles:

5

A mixture of a bromoketone (1 equivalent), thiourea (1 equivalent), and diisopropylethylamine (2 equivalents) in acetone or anhydrous ethanol was heated at reflux temperature overnight. The solvent was evaporated, the residue dissolved in dichloromethane and washed with saturated aqueous sodium bicarbonate solution. The aqueous phase was extracted with dichloromethane three times, the combined organic extracts were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude product was purified by flash column chromatography (silica gel, hexanes : ethyl acetate).

10

15

The following procedure is typical:

20

c. *trans*-2-(4-(*t*-Butoxycarbonylaminomethyl)-cyclohexyl)-amino-4-(2-methoxy-phenyl)1,3-thiazole: A mixture of 2-bromoacetylanisole (1.5 g, 6.5 mmol), *trans*-4-(aminocarbothioylamino)-1-(*t*-butoxycarbonyl-amino)methyl-cyclohexane (1.88 g, 6.5 mmol) and *N,N*-diisopropylethylamine (2.3 ml, 13.1 mmol) in acetone (20 ml) was heated at reflux temperature overnight. The solvent was evaporated, the residue was dissolved in dichloromethane and washed with saturated aqueous sodium bicarbonate solution. The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude product was purified by flash column chromatography over silica gel (hexanes/ethyl acetate 5:2) to give the desired product as a light brown oil (1.2 g, 44% yield): ESMS *m/e* = 418 (*MH*⁺): ¹H NMR δ 8.08 (dd, 1H, *J*

25

30

163

= 1.6, 7.7 Hz), 7.24 (m, 1H), 7.18 (s, 1H), 7.01 (t, 1H, $J = 7.6$ Hz), 6.95 (d, 1H, $J = 8.2$ Hz), 4.96 (d, 1H, $J = 8.1$ Hz), 4.60 (m, 1H), 3.93 (s, 3H), 3.30 (m, 1H), 3.01 (t, 2H, $J = 6.4$ Hz), 2.25 (m, 2H), 1.84 (m, 2H), 1.47 (m, 1H), 1.45 (s, 9H), 1.30-1.03 (m, 4H).

tert-Butyl-(4-{[4-(2-methoxyphenyl)-1,3-thiazol-2-yl]amino}cyclohexyl)methylcarbamate was obtained as a light brown oil in 95% yield from 2-bromo-2'-methoxyacetophenone and tert-butyl {4-[(aminocarbothioyl)amino]cyclohexyl}-methylcarbamate: ^1H NMR δ 8.08 (dd, 1H, $J = 1.6, 7.7$ Hz), 7.24 (m, 1H), 7.18 (s, 1H), 7.01 (t, 1H, $J = 7.6$ Hz), 6.95 (d, 1H, $J = 8.2$ Hz), 4.96 (d, 1H, $J = 8.1$ Hz), 4.60 (m, 1H), 3.93 (s, 3H), 3.30 (m, 1H), 3.01 (t, 2H, $J = 6.4$ Hz), 2.25 (m, 2H), 1.84 (m, 2H), 1.47 (m, 1H), 1.45 (s, 9H), 1.30-1.03 (m, 4H); ESIMS $m/e = 418$ (MH^+).

tert-Butyl-(4-{[4-(3-methoxyphenyl)-1,3-thiazol-2-yl]amino}cyclohexyl)methylcarbamate was obtained as a yellow oil in 86% yield from 2-bromo-3'-methoxyacetophenone and tert-butyl {4-[(aminocarbothioyl)amino]cyclohexyl}-methylcarbamate: ^1H NMR δ 7.36 (m, 2H), 7.30 (m, 1H), 6.83 (m, 1H), 6.68 (s, 1H), 5.05 (d, 1H, $J = 7.9$ Hz), 4.60 (m, 1H), 3.85 (s, 3H), 3.29 (m, 1H), 3.01 (t, 2H, $J = 6.4$ Hz), 2.26 (m, 2H), 1.86 (m, 2H), 1.46 (m, 1H), 1.45 (s, 9H), 1.30-1.03 (m, 4H); ESIMS $m/e = 418$ (MH^+).

tert-Butyl-(4-{[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]amino}cyclohexyl)methylcarbamate was obtained as a yellow oil in 86% yield from 2-bromo-4'-methoxyacetophenone and tert-butyl-{4-[(aminocarbothioyl)amino]cyclohexyl}methylcarbamate: ^1H NMR δ 7.71 (d, 2H, $J = 8.4$ Hz), 6.90 (d, 2H,

164

$J = 8.4$ Hz), 6.55 (s, 1H), 5.01 (m, 1H), 4.60 (m, 1H), 3.83 (s, 3H), 3.29 (m, 1H), 3.01 (t, 2H, $J = 6.3$ Hz), 2.25 (m, 2H), 1.86 (m, 2H), 1.46 (m, 1H), 1.45 (s, 9H), 1.28-1.02 (m, 4H); ESIMS $m/e = 418$ (MH^+).

5

tert-Butyl-[4-({4-[4-(aminocarbonyl)phenyl]-1,3-thiazol-2-yl}amino)cyclohexyl]methylcarbamate was obtained as a yellow solid in more than 95% yield from 4-(2-bromoacetyl)benzamide (1 equivalent) and tert-butyl-{4-[(aminocarbothioyl)amino]cyclohexyl}-methylcarbamate (1 equivalent, no diisopropylethylamine was added for this reaction): 1H NMR (CD_3OD) δ 7.94 (d, 2H, $J = 8.4$ Hz), 7.79 (d, 2H, $J = 8.4$ Hz), 7.07 (s, 1H), 3.57 (m, 1H), 2.91 (d, 2H, $J = 6.6$ Hz), 2.17 (m, 2H), 1.86 (m, 2H), 1.42 (s, 9H), 1.39 (m, 3H), 1.10 (m, 2H); ESIMS $m/e = 431$ (MH^+).

15

General Procedure for the Deprotection of the Boc-bicyclic Thiazoles Intermediates.

20

The Boc-protected 2-amino-1,3-thiazole intermediate was treated with 2N hydrogen chloride in 1,4-dioxane and ethyl acetate (prepared from 4N HCl in dioxane) at room temperature for 2 hours or longer as needed. The desired compound was collected by filtration.

25

N-[4-(Aminomethyl)cyclohexyl]-4-(2-methoxyphenyl)-1,3-thiazol-2-amine dihydrogen chloride was obtained as a light yellow solid in more than 95% yield from tert-butyl-(4-{[4-(2-methoxyphenyl)-1,3-thiazol-2-yl]amino}-cyclohexyl)methyl-carbamate: 1H NMR (CD_3OD) δ 7.54-7.45 (m, 2H), 7.16 (d, 1H, $J = 8.4$ Hz), 7.06 (m, 1H), 7.01 (s, 1H), 3.92 (s, 3H), 3.63 (d, 2H, $J = 1.2$ Hz), 3.50 (m, 1H), 2.83 (m, 2H), 2.21 (m, 2H), 1.96 (m, 2H), 1.69 (m, 1H), 1.49 (m, 2H), 1.22 (m, 2H); ESI $m/e = 318$ (MH^+).

30

N-[4-(Aminomethyl)cyclohexyl]-4-(3-methoxyphenyl)-1,3-thiazol-2-amine dihydrogen chloride was obtained as a pale solid in more than 95% yield from *tert*-butyl 4-{[4-(3-methoxyphenyl)-1,3-thiazol-2-yl]-amino}cyclohexyl)methyl-carbamate.

N-[4-(Aminomethyl)cyclohexyl]-4-(4-methoxyphenyl)-1,3-thiazol-2-amine dihydrogen chloride was obtained as a pale solid in more than 95% yield from *tert*-butyl-4-{[4-(4-methoxyphenyl)-1,3-thiazol-2-yl]-amino}cyclohexyl)methyl-carbamate: ^1H NMR (CD_3OD) δ 7.60 (d, 2H, $J = 8.6$ Hz), 7.02 (d, 2H, $J = 8.6$ Hz), 6.89 (s, 1H), 3.83 (s, 3H), 3.63 (m, 1H), 2.85 (m, 2H), 2.20 (m, 2H), 1.95 (m, 2H), 1.72 (m, 1H), 1.52 (m, 2H), 1.28 (m, 2H).

trans-4-(2-{[4-(Aminomethyl)cyclohexyl]amino}-1,3-thiazol-4-yl)benzamide dihydrogen chloride was obtained as a yellow solid in more than 95% yield from *tert*-butyl [4-({4-[4-(aminocarbonyl)phenyl]-1,3-thiazol-2-yl}amino)cyclohexyl]-methylcarbamate: ^1H NMR (CD_3OD) δ 7.99 (d, 2H, $J = 8.5$ Hz), 7.78 (d, 2H, $J = 8.5$ Hz), 7.19 (s, 1H), 3.69 (m, 1H), 2.84 (m, 1H), 2.21 (m, 2H), 1.95 (m, 2H), 1.71 (m, 1H), 1.53 (m, 2H), 1.08 (m, 2H).

General Procedure for the Derivatization of Amines with Carboxylic Acid and Sulfonic Acid Derivatives:

An amine such as N1 - (4,5-Dihydrobenzo[2,3]thiepino-[4,5-d][1,3]thiazol-2-yl)-1,6-hexanediamine or N2-[4-(aminomethyl)cyclohexyl]-4,5-dihydrobenzo[2,3]-thiepino[4,5-d][1,3]thiazol-2-amine (0.305 mmol) was dissolved in 2 ml CH_2Cl_2 containing 2 equivalents of diisopropylethylamine. A sulfonyl chloride, sulfamoyl

166

chloride, acid chloride or carbamoyl chloride (1-3 equivalents) was added dropwise. The reaction mixture was stirred at room temperature for 1-3 days, quenched with water, washed with 10% NaHCO₃, dried over Na₂SO₄ and chromatographed using column chromatography, preparative TLC or reverse phase HPLC.

10

General Procedure for the Formation of Formamides:

tert-Butyl-*N*-[4-(isopropylamino)cyclohexyl]methylcarbamate:

15 Isopropyl iodide (2 equivalents) was added dropwise to a suspension of *tert*-butyl *N*-[4-aminocyclohexyl]-methylcarbamate (1 equivalent) and diisopropylethyl amine (3 equivalents) in THF. The resulting mixture was stirred for 1 day. TLC analysis showed some starting amine.

20 Additional isopropyl iodide (1 equivalent) and diisopropylethyl amine (3 equivalents) were added to the reaction mixture and heated at 40 °C for 1 day. The reaction mixture was concentrated and chromatographed to give *tert*-butyl *N*-[4-

25 (isopropylamino)cyclohexyl]methylcarbamate: 22% yield, 271 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 4.65 (broad, 1H), 2.91 (m, 3H), 2.42 (m, 1H), 1.80 (ABm, 4H), 1.38 (s, 9H), 0.98 (d, 6H, J=6.3 Hz), 1.32-0.85 (m, 5H).

30

Similarly, *tert*-butyl-*N*-[4-(2-methoxyethylamino)-cyclohexyl]methylcarbamate was obtained (2-methoxyethyl bromide and n-Bu₄NI were used): 35% yield, 378 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 4.64 (broad, 1H), 3.44 (m, 2H), 3.31 & 3.30 (two s, 3H), 2.92 (m, 2H), 2.74 (m, 2H), 2.33 (m,

167

1H), 1.81 (ABm, 4H), 1.39 & 1.38 (two s, 9H), 1.34 (m, 1H), 0.98 (m, 4H).

tert-butyl N-[4-(isopropylformylamino)cyclohexyl]-methylcarbamate: A solution of a tert-butyl N-[4-(isopropylamino)cyclohexyl]methylcarbamate (7.89 mmol, 1 equivalent) in 5 ml of THF was added dropwise to a solution of 1H-benzotriazole-1-carboxaldehyde (8.68 mmol, 1.2 equivalent) in 10 ml of THF at room temperature. The reaction mixture was stirred overnight and then heated at reflux temperature for two hours. 1H-benzotriazole-1-carboxaldehyde (additional 1 equivalent) was added to the reaction mixture and stirred overnight. The solvent was removed in vacuo and dichloromethane was added to the residue. The organic phase was washed with 2N NaOH solution, saturated NaCl solution, dried over Na₂SO₄, and the solvent removed in vacuo. The residue was then chromatographed to give tert-butyl N-[4-(isopropylformylamino)cyclohexyl]methylcarbamate: 100% yield, 299 (ESMS, MH⁺); ¹H NMR (CD₃OD) δ 8.22 & 8.18 (two s, 1H), 4.63 (broad, 1H), 4.30 & 3.60 (two m, 1H), 3.76 (m, 1H), 2.99 (m, 2H), 1.44 (s, 9H), 1.27 (d, 3H, J=6.5 Hz), 1.21 (d, 3H, J=6.5 Hz), 1.91-0.82 (m, 9H).

Similarly, N-[4-(2-methoxyethylformylamino)-cyclohexyl]-methylcarbamate was prepared: 58% yield; 315 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 8.25 & 8.16 (two s, 1H), 4.80 (broad, 1H), 4.07 & 3.23 (two m, 1H), 3.50 (m, 2H), 3.40-3.33 (m, 2H), 3.31 (s, 3H), 2.99 (m, 2H), 1.46 (s, 9H), 1.86-0.95 (m, 9H).

N-[4-(Aminomethyl)cyclohexyl]-N-isopropylformamide:

Dioxane containing HCl (10 ml of 4N HCl solution) was added to a solution of tert-butyl N-[4-

168

(isopropylformylamino)-cyclohexyl]methylcarbamate dissolved in 10 ml Et₂O, stirred at room temperature for 2 hours, and the solvent removed to obtain *N*-[4-(aminomethyl)cyclohexyl]-*N*-isopropylformamide: 100% yield, 199 (ESMS, MH⁺); ¹H NMR (CD₃OD) δ 8.16 (s, 1H), 4.16 & 3.57 (two m, 1H), 3.70 (m, 1H), 2.79 (m, 2H), 1.36 (m, 6H), 1.91-1.06 (m, 9H).

Similarly, *N*-[4-(aminomethyl)cyclohexyl]-*N*-(2-methoxyethyl)-formamide was obtained: 100% yield; 215 (ESMS, MH⁺); ¹H NMR (CD₃OD) δ 8.44 & 8.03 4.65 (two s, 1H), 3.79-3.36 (m, 7H), 3.71 (s, 3H), 2.12-1.13 (m, 9H).

N-Benzoyl-*N'*-[4-(isopropylformylamino)cyclohexyl]-methylthiourea: A mixture of *N*-[4-(aminomethyl)-cyclohexyl]-*N*-isopropylformamide salt (4.55 mmol, 1 equivalent), benzoyl isothiocyanate (5.46 mmol, 1.2 equivalents) and triethylamine (5.46 mmol, 1.2 equivalents) in THF (50 ml) were stirred at room temperature overnight. The removal of solvent and chromatography afforded the desired product: 39% yield, 362 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 10.87 (broad, 1H), 9.20 (broad, 1H), 8.20 & 8.18 (two s, 1H), 7.83 (d, 2H, J=7.7 Hz), 7.60 (m, 1H), 7.49 (m, 2H), 4.26 (m, 1H), 3.76 & 3.08 (two m, 1H), 3.57 (m, 2H), 1.25 (d, 3H, J=6.8 Hz), 1.19 (d, 3H, J=6.8 Hz), 1.97-1.03 (m, 9H).

Similarly, *N*-Benzoyl-*N'*-[4-(2-methoxyethylformyl-amino)-cyclohexyl]methylthiourea was obtained: 100% yield, 378 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 10.85 (broad, 1H), 9.03 (broad, 1H), 8.18 & 8.08 (two s, 1H), 7.84 (d, 2H, J=7.9 Hz), 7.64 (m, 1H), 7.52 (d, 2H, J=7.8 Hz), 3.63-3.24 (m, 7H), 3.34 & 3.33 (two m, 3H), 2.03-1.13 (m, 9H).

N-[4-

(isopropylformylamino)cyclohexyl]methylthiourea: K_2CO_3 (2 equivalents) in water was added to a solution of *N*-benzoyl-*N'*-[4-(isopropylformylamino)-cyclohexyl]methylthiourea in MeOH and stirred at room temperature overnight. The solvent was removed in vacuo and the residue was dissolved in EtOH. The solution was filtered to remove a white precipitate and the filtrate concentrated. The crude product was chromatographed to yield the desired product: 100% yield; 100% yield; 258 (ESMS, MH^+); 1H NMR (CD_3OD) δ 8.15 & 8.13 (two s, 1H), 4.15 & 3.73 (two m, 1H), 3.34 & 2.97 (two m, 1H), 3.29 (m, 2H), 1.26 (d, 3H, $J=6.7$ Hz), 1.23 (d, 3H, $J=6.7$ Hz), 1.91-1.03 (m, 9H).

Similarly, *N*-[4-(2-methoxyethylformylamino)-cyclohexyl]methylthiourea was prepared: 77% yield, 274 (ESMS, MH^+); 1H NMR (CD_3OD) δ 8.15 & 8.00 (two s, 1H), 7.55 & 7.43 (two m, 1H), 3.90 & 2.97 (two m, 1H), 3.46-3.28 (m, 10H), 1.90-0.99 (m, 9H).

Formation of 2-amino-1,3-thiazoles containing a formamide:

A thiourea such as *N*-[4-(isopropylformylamino)cyclohexyl]methylthiourea (0.029 mmol, 1 equivalent), a bromoketone such as 2-bromoacetylbenzene (0.044 mmol, 1.5 equivalents) and 2 equivalents of diisopropylethyl amine in 10 ml of EtOH were heated at reflux temperature for 2 days. The reaction mixture was concentrated and the crude product was chromatographed (silica) to afford the desired product. This sequence of reactions were utilized for the preparation of examples 98-99.

Alternatively, the terminal amino functionality of a primary amine such as *N*-[4-(aminomethyl)cyclohexyl]-4-(2-methoxyphenyl)-1,3-thiazol-2-amine was alkylated using alkyl halides or a sequence of acylation followed by reduction to give a secondary amine. The secondary amine was, in turn, used in the formation of the formamide utilizing a formylating reagent such as benzotriazole-1-carboxaldehyde. This sequence of reactions were utilized in the preparation of examples 96-97.

N-{4-[(isopropylamino)methyl]cyclohexyl}-4-(2-methoxyphenyl)-1,3-thiazol-2-amine: The reaction mixture of *N*-[4-(aminomethyl)cyclohexyl]-4-(2-methoxyphenyl)-1,3-thiazol-2-amine (1 equivalent), 2-iodopropane (1 equivalent), and potassium carbonate (2 equivalents) in DMF was heated at about 50 °C overnight. Water was then added to the reaction mixture and extracted with diethyl ether (3 times). Evaporation of the combined organic phase afforded a yellow oil which was purified by preparative TLC plates. The desired product was obtained as a yellow oil in 53% yield: ESIMS *m/e* = 360 (MH⁺).

N-(4-{[2-Methoxyethyl]amino}methyl)cyclohexyl)-4-(2-methoxyphenyl)-1,3-thiazol-2-amine was obtained as a light yellow oil in 82% yield from 2-methoxy-*N*-[(4-{[4-(2-methoxyphenyl)-1,3-thiazol-2-yl]amino}-cyclohexyl)methyl]-acetamide and borane (4 equivalents, 1M in THF): ESIMS *m/e* = 376 (MH⁺).

2-Methoxy-*N*-[(4-{[4-(2-methoxyphenyl)-1,3-thiazol-2-yl]amino}cyclohexyl)-methyl]acetamide was obtained as a yellow oil in 87% yield from *N*-[4-(aminomethyl)cyclohexyl]-4-(2-methoxyphenyl)-1,3-thiazol-2-amine and methoxyacetyl chloride.

A sequence of reactions depicted in schemes 7-11 were utilized for the preparation of examples 91-95.

5

Example 91

trans-4-(2-Methoxy)phenyl-2-(4-methylsulfonylamino-methyl)-cyclohexylamino-1,3-thiazole hydrochloride. Obtained as a white solid in 43% yield from *trans*-2-(4-aminomethyl)cyclohexylamino-4-(2-methoxyphenyl)-1,3-thiazole dihydrochloride and methanesulfonyl chloride: ESMS $m/e = 396$ (MH^+); 1H NMR (free base) δ 8.06 (dd, 1H, $J = 1.6, 7.7$ Hz), 7.25 (m, 1H), 7.17 (s, 1H), 7.03-6.94 (m, 2H), 5.09 (d, 1H, $J = 7.7$ Hz), 4.59 (m, 1H), 3.92 (s, 3H), 3.31 (m, 1H), 2.99 (t, 2H, $J = 6.5$ Hz), 2.96 (s, 3H), 2.26 (m, 2H), 1.88 (m, 2H), 1.49 (m, 1H), 1.29-1.07 (m, 4H).

15

Example 92

trans-2-(4-Ethoxycarbonylaminomethyl)cyclohexylamino-4-(2-methoxy)phenylthiazole hydrogen chloride was obtained as a light yellow solid in 62% from *N*-[4-(aminomethyl)cyclohexyl]-4-(2-methoxyphenyl)-1,3-thiazol-2-amine dihydrogen chloride and ethyl chloroformate: 1H NMR ($CDCl_3$, free base) δ 8.07 (dd, 1H, $J = 1.3, 7.7$ Hz), 7.24 (m, 1H), 7.18 (s, 1H), 7.03-6.93 (m, 2H), 5.01 (m, 1H), 4.72 (m, 1H), 4.11 (q, 2H, $J = 7.0$ Hz), 3.93 (s, 3H), 3.31 (m, 1H), 3.06 (m, 2H), 2.26 (m, 2H), 1.85 (m, 2H), 1.49 (m, 1H), 1.24 (t, 3H, $J = 7.0$ Hz), 1.30-1.05 (m, 4H); ESIMS $m/e = 390$ (MH^+).

25

30

Example 93

172

trans-4-(3-Methoxy)phenyl- 2-(4-methylsulfonylamino-methyl)-cyclohexylaminothiazole was obtained as a light yellow solid in 84% from *N*-[4-(aminomethyl)cyclohexyl]-4-(3-methoxyphenyl)-1,3-thiazol-2-amine dihydrogen chloride and methanesulfonyl chloride: ^1H NMR δ 7.36 (m, 2H), 7.27 (m, 1H), 6.83 (m, 1H), 6.69 (s, 1H), 5.08 (m, 1H), 4.33 (m, 1H), 3.85 (s, 3H), 3.35 (m, 1H), 3.02 (t, 2H, $J = 6.6$ Hz), 2.97 (s, 3H), 2.28 (m, 2H), 1.91 (m, 2H), 1.58 (m, 1H), 1.28-1.05 (m, 4H); ESIMS $m/e = 396$ (MH^+).

10

Example 94

trans-2-(4-Ethoxycarbonylaminomethyl)cyclohexylamino-4-(3-methoxy)phenylthiazole hydrogen chloride was obtained as a yellow solid in 81% from *N*-[4-(aminomethyl)cyclohexyl]-4-(3-methoxyphenyl)-1,3-thiazol-2-amine dihydrogen chloride and ethyl chloroformate: ^1H NMR (free base) δ 7.36 (m, 2H), 7.27 (m, 1H), 6.83 (m, 1H), 6.69 (s, 1H), 5.01 (m, 1H), 4.71 (m, 1H), 4.12 (m, 2H), 3.85 (s, 3H), 3.30 (m, 1H), 3.07 (t, 2H, $J = 6.5$ Hz), 2.25 (m, 2H), 1.86 (m, 2H), 1.49 (m, 1H), 1.30-1.08 (m, 7H); ESIMS $m/e = 390$ (MH^+).

25

30

General Procedure for the Demethylation of Methoxyphenylthiazoles Using Boron Tribromide:

5 Boron tribromide (3 equivalents, 1M in CH_2Cl_2) was added dropwise to a cooled solution of methoxyphenylthiazole (1 equivalent) in dichloromethane. The reaction mixture was stirred at -78°C for ten minutes, then warmed up to room temperature and stirred for an additional hour. Water was
10 added to destroy the boron complex and 2M sodium hydroxide was added to basify the mixture to pH 13-14. The resulting mixture was extracted with chloroform (3 times) and the combined organic extracts were dried and concentrated in vacuo. The crude product was purified by flash column
15 (silica gel, hexane : ethyl acetate). In most cases, a light yellow oil was obtained which was converted to a hydrogen chloride salt.

Example 95

trans-4-(2-Hydroxy)phenyl-2-(4-methylsulfonylamino-methyl)-cyclohexylaminothiazole was obtained as a light brown solid in 50% yield from *trans*-4-(2-methoxy)phenyl-2-(4-methylsulfonylaminomethyl)-cyclohexylaminothiazole hydrogen chloride and boron tribromide: ^1H NMR δ 11.01 (b, 1H), 8.10 (dd, 1H, $J = 1.4, 7.9$ Hz), 7.25 (m, 2H), 6.89 (m, 2H), 5.18 (d, 1H, $J = 7.5$ Hz), 4.46 (m, 1H), 3.39 (m, 1H), 3.00 (t, 2H, $J = 6.5$ Hz), 2.96 (s, 3H), 2.22 (m, 2H), 1.91 (m, 2H), 1.60 (m, 1H), 1.30-1.04 (m, 4H) ; ESIMS $m/e = 382$ (MH^+).

Example 96

trans-2-(4-(*N*-Isopropyl)formamidomethyl)cyclohexyl-amino-4-(2-methoxy)phenylthiazole hydrogen chloride was obtained as a yellow solid in 70% from *N*-{4-[(isopropylamino)methyl]-cyclohexyl}-4-(2-methoxy-phenyl)-1,3-thiazol-2-amine and 1*H*-benzotriazole-1-carboxaldehyde: ^1H NMR (CDCl_3 , free base) δ 8.25 (s, 0.6 H), 8.08 (dd, 1H, $J = 1.7, 7.7$ Hz), 7.99 (s, 0.4H), 7.24 (m, 1H), 7.18 (s, 1H), 7.03-6.93 (m, 2H), 5.04 (t, 1H, $J = 8.7$ Hz), 4.31 (p, 0.4H, $J = 6.8$ Hz), 3.92 (s, 3H), 3.64 (p, 0.6H, $J = 6.8$ Hz), 3.31 (m, 1H), 3.11 (d, 1.3H, $J = 7.3$ Hz), 3.01 (d, 0.7 Hz, $J = 7.3$ Hz), 2.24 (m, 2H), 1.79 (m, 2H), 1.69 (m, 0.6H), 1.51 (m, 0.4H), 1.30-1.00 (m, 10H); ESIMS $m/e = 388$ (MH^+).

Example 97

trans-2-(4-(*N*-(2-Methoxyethoxy)formamido)methyl)-cyclohexylamino-4-(2-methoxy)phenylthiazole hydrogen chloride was obtained as a yellow syrup in 70% from *N*-(4-{[2-methoxyethyl)amino]methyl}cyclohexyl)-4-(2-methoxyphenyl)-1,3-thiazol-2-amine and 1*H*-benzo-triazole-1-carboxaldehyde: ¹H NMR (CDCl₃, free base) δ 8.11 (s, 0.4H), 8.08 (dd, 1H, *J* = 1.6, 7.8 Hz), 8.02 (s, 0.6 H), 7.24 (m, 1H), 7.18 (s, 1H), 7.00 (t, 1H, *J* = 7.5 Hz), 6.94 (d, 1H, *J* = 8.2 Hz), 5.01 (m, 1H), 3.92 (s, 3H), 3.56-3.14 (m, 7H), 2.25 (m, 2H), 1.76 (m, 2H), 1.66 (m, 1H), 1.29-0.97 (m, 4H); ESIMS *m/e* = 404 (MH⁺).

15

Example 98

trans-N-(2-Methoxyethyl)-*N*-(4-[(4-phenyl-1,3-thiazol-2-yl)amino]methylcyclohexyl)formamide: 29% yield, 374 (ESMS, MH⁺): ¹H NMR (CDCl₃) δ 8.18 & 8.08 (two s, 1H), δ 7.78 (d, 2H, *J*=8.6 Hz), 7.37 (apparent t, 2H, *J*=7.7 Hz), 7.29 (d, 1H, *J*=7.0 Hz), 6.69 (s, 1H), 5.49 (b, 1H), 4.09 (m, 1H), 3.49-3.13 (m, 7H), 3.34 & 3.33 (two s, 3H), 2.00-0.85 (m, 9H).

20

25

Example 99

trans-N-Isopropyl-*N*-(4-[(4-phenyl-1,3-thiazol-2-yl)amino]methylcyclohexyl)formamide: 84% yield, 358 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ 8.21 & 8.17 (two s, 1H), δ 7.78 (d, 2H, *J*=8.7 Hz), 7.37 (apparent t, 2H, *J*=7.7 Hz), 7.28 (d, 1H, *J*=7.0 Hz), 6.68 (s, 1H), 5.63 & 5.77 (two b, 1H), 4.28 & 3.58 (two m, 1H), 3.78 & 3.03 (two m, 1H), 3.14 (m, 2H), 1.27 (d, 3H, *J*=6.9 Hz), 1.21 (d, 3H, *J*=6.9 Hz), 1.96-1.03 (m, 9H).

30

I. Synthetic Methods for ExamplesC. Tricyclic Compounds5 **General Procedures Relating to Examples:**

For the formation of 2-aminothiazoles from 2-haloketones and thioureas, see, for example, Kearney, P.C., et al., 1998; Di Fabio, R. and Pentassuglia, G., 1998; De Kimpe, N., et al., 1996; Plazzi, P.V., et al., 1995; and
10 Novikova, A. P., 1991.

For the formation of thiazoles from 2-haloketones and thioamides, see, for example, Critcher, D. J. and Pattenden, G., 1996; and Friedman, B. S., et al., 1937.

15 For the formation of 2-aminoimidazoles from 2-haloketones and guanidines, see, for example, Little, T. L. and Webber, 1994; and Chabaka, L.M., et al., 1994.

20 For the formation of imidazoles from 2-haloketones and amidines, see, for example, Demchenko, A. M., et al., 1997; and Nagao, Y., et al., 1996.

25 For the synthesis of 2-aminooxazoles from 2-haloketones and ureas, see, for example, Pathak, V.N., et al., 1993; Crangk, G. and Foulis, M.J., 1971; and Marchetti, E., et al., 1968.

30 For the formation of oxazoles from 2-haloketones and amides, see, for example, Hammar, W.J. and Rustad, M.A., 1981; and Zhao, Z., et al., 1991.

All reactions were performed under an inert atmosphere (Argon) and the reagents, neat or in appropriate solvents,

were transferred to the reaction vessel via syringe and cannula techniques. The parallel synthesis reaction arrays were performed in vials (without an inert atmosphere) using J-KEM heating shakers (Saint Louis, MO). Unless stated otherwise all solvents were AR grade and used as supplied. Anhydrous solvents were purchased from Aldrich Chemical Company and used as received. Examples 1-7 described in this application were named using ACD/Name program (version 2.51, Advanced Chemistry Development Inc., Toronto, Ontario, M5H2L3, Canada).

^1H and ^{13}C spectra were recorded at 300 and 75 MHz (QE Plus) with CDCl_3 as solvent (unless otherwise noted) and tetramethylsilane as internal standard. s = singlet; d = doublet; t = triplet; q = quartet; p = pentet; sextet; septet; b = broad; m = multiplet. Elemental analyses were performed by Robertson Microlit Laboratories, Inc. Low-resolution electrospray MS spectra were measured (ESMS, MS) and MH^+ is reported. Thin-layer chromatography (TLC) was carried out on glass plates precoated with silica gel 60 F₂₅₄ (0.25 mm, EM Separations Tech.). Preparative thin-layer chromatography was carried out on glass sheets precoated with silica gel GF (2 mm, Analtech). Flash column chromatography was performed on Merck silica gel 60 (230 - 400 mesh). Melting points were determined in open capillary tubes on a Med-Temp apparatus and are uncorrected.

General Procedure for the Synthesis of Benzothiepin-5-ones:

2,3,4,5-Tetrahydro-1-benzothiepin-5-one

5

Step 1.

4-(phenylsulfanyl)butanoic acid:

10 Sodium methoxide (1.2 equivalent) was added to 60 ml of ethanol, in one portion, and the suspension was stirred at room temperature. Thiophenol (1 equivalent) was added to the above suspension and stirred at room temperature for an additional 30 minutes. Butyrolactone (1.1 equivalent) was then added and the resulting mixture was stirred at
15 reflux temperature for 6 hours, cooled to room temperature and concentrated in vacuo. The residual solid was washed with 200 ml hexane/ether 2:1, v/v. The solid was suspended in ice cold 2N HCl and stirred for 15 minutes. The solid was then filtered, washed with 100 ml hexane/ether and
20 dried under reduced pressure at room temperature to give 4-(phenylsulfanyl)butanoic acid as a tan solid: 52% yield; ^1H NMR (CDCl_3) δ 7.32-7.12 (m, 5H), 2.94 (t, 2H, $J=7.2$ Hz), 2.41 (t, 2H, $J=7.2$ Hz), 1.85 (p, 2H, $J=7.2$ Hz); Anal. Calc. For $\text{C}_{10}\text{H}_{12}\text{S}_1\text{O}_2$: C, 61.22; H, 6.12. Found: C, 61.16; H,
25 6.28.

A similar procedure was used for the synthesis of 4-(4-fluorophenylsulfanyl)butanoic acid: 60% yield; ^1H NMR (CDCl_3) δ 7.34 (m, 2H, 7.00 (m, 2H), 2.94 (t, 2H, $J=7.2$ Hz), 2.51 (t, 2H, $J=7.2$ Hz), 1.93 (p, 2H, $J=7.2$ Hz); Anal. Calc. For $\text{C}_{10}\text{H}_{11}\text{F}_1\text{S}_1\text{O}_2$: C, 56.07; H, 5.14. Found: C, 55.80; H, 5.19.

30

Step 2.

35

Benzothiepin-5-ones:

Polyphosphoric acid (6 equivalents) was heated to 80°C under argon. 4-(Phenylsulfanyl)butanoic acid, (1 equivalent) was added in portions and the mixture heated at 100°C for 2 hours. The reaction mixture was cooled, poured into ice cold water and then extracted with 2X100 ml ethyl acetate. The combined ethyl acetate extracts were washed with 100 ml water, 100 ml saturated sodium bicarbonate, and 100 ml water. The ethyl acetate extract was dried (anhydrous sodium sulfate), filtered and the solvent removed in vacuo to give a tan solid. This solid was dried to give 2,3,4,5-tetrahydro-1-benzothiepin-5-one: 52% yield; ^1H NMR (CDCl_3) δ 7.824 (dd, 1H, $J=0.9, 7.5$ Hz), 7.45 (dd, 1H, $J=0.6, 6.9$ Hz), 7.34-7.21 (m, 2H), 3.05 (t, 2H, $J=6.6$ Hz), 2.97 (t, 2H, $J=6.6$ Hz), 2.29 (p, 2H, $J=6.6$ Hz).

The above described procedure was also used to afford 7-fluoro-2,3,4,5-tetrahydro-1-benzothiepin-5-one: 60% yield; ^1H NMR (CDCl_3) δ 7.51 (dd, 1H, $J=3.0, 9.3$ Hz), 7.41 (dd, 1H, $J=8.7, 5.1$ Hz), 7.04 (apparent dt, 1H, $J=3.0, 4.8$ Hz), 3.06 (t, 2H, $J=6.6$ Hz), 2.96 (t, 2H, $J=6.6$ Hz), 2.64 (t, 2H, $J=6.9$ Hz); Anal. Calc. For $\text{C}_{10}\text{H}_{10}\text{S}_1\text{O}_1$: C, 67.41; H, 5.61. Found: C, 67.48; H, 5.68.

General Procedure for the Synthesis of Bromoketones:

To a cooled solution of the ketone (1 equivalent) in acetic acid, was slowly added bromine (1 equivalent). The reaction mixture was stirred at room temperature for 3 hours. The solvent was evaporated, the residue dissolved in dichloromethane, and the resulting solution washed with saturated sodium bicarbonate and water. The organic phase was dried over sodium sulfate. Evaporation of the combined, decolorized organic phase gave a light yellow oil in > 80% yield in most cases.

7-Fluoro-2,3,4,5-tetrahydro-1-benzothiepin-5-one was brominated according to the procedure described below to give 4-bromo-7-fluoro-2,3,4,5-tetrahydro-1-benzothiepin-5-one. A similar procedure was also used to brominate 2,3,4,5-tetrahydro-1-benzothiepin-5-one.

4-Bromo-7-fluoro-2,3,4,5-tetrahydro-1-benzothiepin-5-one:
7-Fluoro-2,3,4,5-tetrahydro-1-benzothiepin-5-one (1 equivalent) was also dissolved in glacial acetic acid and stirring continued at room temperature. Bromine (2.5 equivalents) was added dropwise and stirring continued at room temperature for 4 hours. Water was added to the reaction mixture was then extracted with 2x25 ml ethyl acetate. The combined ethyl acetate extracts were washed with water, saturated sodium bicarbonate, and water. The combined extract was dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo to give a solid which was recrystallized from the ethyl acetate/hexane 1:1 v/v to afford 4-bromo-7-fluoro-2,3,4,5-tetrahydro-1-benzothiepin-5-one: ^1H NMR (CDCl_3) δ 7.55 (dd, 1H, $J=2.7$, 9.0 Hz), 7.44 (dd, 1H, $J=8.7$, 5.1 Hz), 7.11 (Apparent dt,

1H, J=2.7, 4.8 Hz), 5.34¹⁸¹ (dd, 1H, J=5.7, 10.2 Hz),
3.20-2.50 (m, 4H).

4-Bromo-2,3,4,5-tetrahydro-1-benzothiepin-5-one was
5 similarly prepared from the corresponding ketone: ¹H NMR
(CDCl₃) δ 7.83 (d, 1H, J=7.8 Hz), 5.35 (dd, 1H, J=5.7, 10.5
Hz), 3.30-2.50 (m, 4H).

General Procedure for the Synthesis of Boc Protected 10 Thioureas:

A protected diamine such as N-Boc-1,4-diaminobutane or N-
Boc-1,5-diaminopentane (1 equivalent) was dissolved in
tetrahydrofuran and stirred at room temperature. Benzoyl
15 thioisocyanate (1 equivalent) was added dropwise to the
aforementioned solution. The resulting mixture was stirred
at room temperature for 24 hours. The solvent was removed
under reduced pressure to give a yellow oil.

20 The yellow oil (1 equivalent) was then dissolved in
methanol, an aqueous potassium carbonate (3 equivalents)
solution added, and the mixture stirred for 48 hours.
Water was added to the reaction mixture, which was then
extracted with 2x75 ml ethyl acetate. The combined
25 extracts were then washed with water, dried over anhydrous
sodium sulfate, filtered and the solvent removed under
reduced pressure to give the desired thiourea.

tert-Butyl 5-[(aminocarbothioyl)aminol]pentylcarbamate was
30 obtained as a light yellow wax from tert-butyl 5-
{[(benzoylamino)carbothioyl]amino}-pentylcarbamate. ¹H NMR
(CD₃OD) δ 3.44 (m, 1H), 3.10 (m, 1H), 3.01 (t, 2H, J = 6.7
Hz), 1.60-1.31 (m, 6H), 1.41 (s, 9H); 262 (ESMS, MH⁺).

182

tert-Butyl

5-

{[(benzoylamino)carbothioyl]amino}-pentylcarbamate was obtained a light yellow solid in 79% yield from *N*-BOC-1,5-diaminopentate and benzoyl isothiocyanate; m.p. 90-93 °C.

5

tert-Butyl 4-[(aminocarbothioyl)amino]butylcarbamate was obtained as a light yellow wax from tert-butyl 4-{[(benzoylamino)carbothioyl]amino}-butylcarbamate. ¹H NMR (CD₃OD) δ 3.48 (m, 1H), 3.10 (m, 1H), 3.05 (t, 2H, *J* = 6.5 Hz), 1.60 (m, 4H), 1.42 (s, 9H); 248 (ESMS, MH⁺).

10

tert-Butyl 4-{[(benzoylamino)carbothioyl]-amino}butylcarbamate was obtained as a light brown oil in 93% yield from *N*-BOC-1,4-diaminobutane and benzoyl isothiocyanate.

15

trans-tert-Butyl {4-[(aminocarbothioyl)amino]-cyclohexyl}methylcarbamate was obtained as a light yellow wax from *trans*-tert-butyl (4-{[(benzoylamino)carbothioyl]amino} cyclohexyl)-methylcarbamate. ¹H NMR (CD₃OD) δ 3.92 (m, 1H), 2.86 (m, 2H), 2.00 (m, 2H), 1.76 (m, 2H), 1.41 (s, 9H), 1.37 (m, 1H), 1.06 (m, 4H); 288 (ESMS, MH⁺).

20

trans-tert-Butyl- (4-{[(benzoylamino)carbothioyl]-amino}cyclohexyl)methylcarbamate was obtained as a yellow solid in 97% yield from tert-butyl 4-aminocyclohexylmethylcarbamate and benzoyl isothiocyanate.

25

trans-tert-Butyl 4-aminocyclohexylmethylcarbamate was obtained in more than 95 % yield from hydrogenation of benzyl 4-{[(tert-butoxycarbonyl)amino]methyl}-cyclocarbamate.

30

Benzyl-4-[[[tert-

butoxycarbonyl]amino]methyl]

cyclohexylcarbamate: To a stirred suspension of

4-[[[(tert-butoxycarbonyl)amino]methyl]

5 cyclohexanecarboxylic acid (Maybridge Chemical Co., Ltd.) (45g) and diphenylphosphoryl azide (44 ml) in toluene (600 ml) was added triethylamine (32 ml) over a period of 20 min whilst maintaining the internal temperature at -10-0 C. The mixture was slowly warmed and then stirred at
10 70 C for 4h. After cooling to 40 C, benzyl alcohol (36 ml) was added and the reaction mixture heated at reflux for 20 h. The cold reaction mixture was washed with water and brine and dried over anhydrous magnesium sulfate. Removal of the solvent and recrystallization of the
15 organic residue from ethyl acetate and diethyl ether gave the title compound, benzyl-4-[[[tert-butoxycarbonyl]amino]methyl]cyclohexylcarbamate as a white solid, m.p. 129-131 C.

20 *trans*-*tert*-Butyl {4-[(aminocarbothioyl)amino]-cyclohexyl}carbamate was obtained as a yellow solid from *trans*-*tert*-butyl 4-{[(benzoylamino)carbothioyl]-amino}cyclohexyl)-carbamate: ¹H NMR (CD₃OD) δ 3.94 (m, 1H), 3.30 (m, 1H), 2.00 (m, 2H), 1.90 (m, 2H), 1.41 (s, 9H), 1.26
25 (m, 4H); 274 (ESMS, MH⁺).

trans-*tert*-Butyl 4-{[(Benzoylamino)carbothioyl]-amino}cyclohexyl)-carbamate was obtained as a white solid in 66% yield from *tert*-butyl 4-aminocyclohexylcarbamate
30 and benzoyl isothiocyanate.

trans-*tert*-Butyl 4-aminocyclohexylcarbamate was obtained as a light yellow wax in more than 95% yield from

184

hydrogenation of benzyl 4- [(tert-butoxycarbonyl)amino]cyclohexylcarbamate.

trans-Benzyl 4- {[(aminocarbothioyl) amino] - methyl} cyclohexylcarbamate was obtained as a yellow solid in 71% yield from trans-benzyl 4- {[(Benzoylamino) carbothioyl] amino} methyl - cyclohexylcarbamate; 322 (ESMS, MH⁺).

trans-Benzyl 4- {[(benzoylamino) carbothioyl] - amino} methyl - cyclohexylcarbamate was obtained as a yellow solid from benzyl 4- (aminomethyl) cyclohexylcarbamate and benzoyl isothiocyanate.

trans-Benzyl 4- (aminomethyl) cyclohexylcarbamate was obtained as a white solid in more than 95% yield by stirring benzyl 4- {[(tert-butoxycarbonyl) amino] methyl} cyclocarbamate in 2N HCl (made from 1 : 1 of EtOAc and 4N HCl in dioxane).

General Procedure for the Synthesis of the (4,5-dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-ylamino Template.

A mixture of bromoketone (1 equivalent), thiourea (1 equivalent), and diisopropylethylamine (2 equivalents) in anhydrous ethanol was stirred and heated at reflux temperature overnight. The solvent was evaporated, the brown residue dissolved in dichloromethane and washed with saturated aqueous sodium bicarbonate. The aqueous phase was extracted with dichloromethane three times. The combined extracts were dried over anhydrous sodium sulfate. The crude product was purified by flash column

chromatography (silica gel, hexane : ethyl acetate). An example of the aforementioned general procedure follows.

4-Bromo-2,3,4,5-tetrahydro-1-benzothiepin-5-one (1.2 equivalents, 29.76 mmol) and tert-butyl 5-[(aminocarbothioyl)amino]pentylcarbamate (1 equivalent, 24.8 mmol) were mixed with 2 equivalents of diisopropylethylamine in 200 ml of EtOH. The reaction mixture was heated at reflux overnight. The dark brown reaction mixture was concentrated and chromatographed (silica) to obtain tert-butyl N-{5-[(9-fluoro-4,5-dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-yl)amino]pentyl}carbamate as a light tan solid.

General Procedure for the Deprotection of BOC-Protected Amines.

tert-Butyl N-{[4-(4,5-dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-ylamino)cyclohexyl]methyl}carbamate or tert-butyl N-[6-(4,5-dihydrobenzo[2,3]thiepino-[4,5-d][1,3]thiazol-2-ylamino)hexyl]carbamate were dissolved in Et₂O. The same volume of 4N HCl in dioxane was added to make a 2N solution. The reaction mixture was stirred at room temperature overnight, and the solvent removed under reduced pressure to obtain the desired product as its HCl salt.

N1-(4,5-Dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-yl)-1,4-butanediamine: 45% yield; ¹H NMR (CDCl₃) δ 8.05 (dd, 1H, J= 0.56, 8.4 Hz), 7.33 (dd, 1H, J= 0.6, 8.4 Hz), 7.26 (t, 1H, J=6.5 Hz), 7.17 (t, 1H, J=6.5 Hz), 5.91 (broad, 1H), 3.20 (m, 6H), 2.69 (t, 2H, J=6.5 Hz), 1.61-1.27 (m, 6H).

N1- (4,5-

Dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-yl)-1,5-pentanediamine: 50% yield; ^1H NMR (CDCl_3) δ 8.03 (dd, 1H, $J=0.6, 8.4$ Hz), 7.49 (dd, 1H, $J=0.6, 8.4$ Hz), 7.28 (t, 1H, $J=6.5$ Hz), 7.16 (t, 1H, $J=6.5$ Hz), 5.92 (broad, 1H), 3.13 (m, 6H), 2.63 (t, 2H, $J=6.5$ Hz), 1.57-1.37 (m, 8H).

tert-Butyl N-{5-[(9-fluoro-4,5-dihydrobenzo[2,3]-thiepino[4,5-d][1,3]thiazol-2-yl)amino]pentyl}-carbamate: 60% yield; Anal. Calc. for $\text{C}_{21}\text{H}_{28}\text{N}_3\text{S}_2\text{O}_2 + 0.15 \text{CH}_2\text{Cl}_2$: C, 56.41; H, 6.33; N, 9.3. Found: C, 56.45; H, 6.17; N, 8.9; ^1H NMR (CDCl_3) δ 7.72 (dd, 1H, $J=1.15, 7.5$ Hz), 7.47-7.04 (m, 1H), 6.89-6.83 (m, 1H), 6.190-6.142 (m, 1H), 4.747-4.690 (m, 1H), 3.370-2.803 (m, 8H), 1.64-1.048 (m, 6H), 1.407 (s, 9H).

N2- [4-(Aminomethyl)cyclohexyl]-4,5-dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-amine: 73% yield, 346 (ESMS, MH^+); ^1H NMR (CDCl_3) δ 8.05 (dd, 1H, $J=1.2, 7.9$ Hz), 7.50 (dd, 1H, $J=1.2, 7.7$ Hz), 7.32 (apparent dt, 1H, $J=1.8, 7.2$ Hz), 7.15 (apparent dt, 1H, $J=1.7, 7.2$ Hz), 4.93 (b, 1H), 3.23 (m, 1H), 2.99 (t, 2H, $J=6.3$ Hz), 2.56 (d, 2H, $J=6.6$ Hz), 2.04 (ABM, 4H), 1.70-0.80 (m, 12H).

tert-Butyl N-[6-(4,5-dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-ylamino)hexyl]carbamate: 51% yield, 434 (ESMS, MH^+); ^1H NMR (CD_3OD) δ 7.92 (d, 1H, $J=7.5$ Hz), 7.48 (d, 1H, $J=7.6$ Hz), 7.30 (apparent dt, 1H, $J=1.2, 7.7$ Hz), 7.15 (apparent dt, 1H, $J=1.5, 7.5$ Hz), 3.30 (t, 2H, $J=1.6$ Hz), 3.16 (t, 2H, $J=6.3$ Hz), 3.05 (t, 2H, $J=5.9$ Hz), 3.01 (t, 2H, $J=6.5$ Hz), 1.63 (m, 2H), 1.42 (s, 9H), 1.51-1.28 (m, 6H).

N1- (4,5-

Dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-yl)-1,6-

hexanediamine: 75% yield, 334 (ESMS, MH⁺); ¹H NMR (CDCl₃) δ
8.05 (dd, 1H, J=1.0, 8.1 Hz), 7.51 (dd, 1H, J=1.1, 7.8
5 Hz), 7.32 (apparent dt, 1H, J=1.4, 7.4 Hz), 7.15,
(apparent dt, 1H, J=1.6, 7.6 Hz), 5.15 (broad, 1H),
3.23(m, 4H), 3.19 (s, 2H), 2.68 (t, 2H, J=5.7 Hz), 1.70-
1.21 (m, 8H).

10 tert-Butyl N-{[4-(4,5-Dihydrobenzo[2,3]thiepino[4,5-
d][1,3]thiazol-2-ylamino)cyclohexyl]methyl}carbamate: 44%
yield, 446 (ESMS, MH⁺); ¹H NMR (CD₃OD) δ 7.90 (dd, 1H, J=
1.2, 7.8 Hz), 7.49 (dd, 1H, J= 0.8, 7.8 Hz), 7.32
(apparent dt, 1H, J=1.4, 7.7 Hz), 7.16 (apparent dt, 1H,
15 J=1.3, 7.6 Hz), 3.41 (m, 1H), 3.30 (m, 2H), 3.19 (t, 2H,
J=6.5 Hz), 3.06, (t, 2H, J=5.8 Hz), 2.90 (d, 2H, J=7.0
Hz), 1.99 (ABm, 4H), 1.43 (s, 9H), 1.32-1.05 (m, 3H).

20 **General Procedure for the Derivatization of Amines with
Sulfamoyl Chlorides:**

An amine such as N1 -(4,5-Dihydrobenzo[2,3]thiepino[4,5-
d][1,3]thiazol-2-yl)-1,6-hexanediamine or N2-[4-
(aminomethyl)cyclohexyl]-4,5-
25 dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-amine
(0.305 mmol) was dissolved in 2ml CH₂Cl₂ with 2 equivalents
of diisopropylethylamine. A sulfamoyl chloride (1-3
equivalents) was added dropwise. The reaction solution was
stirred at room temperature for 3 days, quenched with
30 water, washed with 10% NaHCO₃, extracted with EtOAc, dried
over Na₂SO₄ and chromatographed using column chromatography
or preparative TLC.

Alternatively, a diamine such as 1,4-bis-aminomethylcyclohexane was derivatized with a sulfamyl chloride which was then derivatized to yield a sulfonyl urea followed by cyclization using a bromoketone as exemplified below:

N'-[4-(Aminomethyl)cyclohexyl]methyl-*N,N*-

dimethylsulfamide: A solution of 1,4-bis-aminomethylcyclohexane in 50 ml of dichloromethane was treated dropwise with 8.0 grams of neat dimethylaminosulfamoyl chloride. The resulting mixture was stirred at room temperature for 24 hours, washed with water and chromatographed (silica) to give 10 grams (25% yield) of *N'*-[4-(aminomethyl)cyclohexyl]methyl-*N,N*-dimethylsulfamide: $^1\text{H NMR}$ (CDCl_3) δ 3.00 (d, 2H, $J=7.5$ Hz), 2.78 (s, 6H), 3.60 (d, 2H, $J=7.5$), 1.85-1.22 (m, 8H), 1.00-0.80 (m, 1H).

N-Benzoyl-*N'*-[4-([(dimethylamino)sulfonyl]amino methyl)-cyclohexyl]methylthiourea and *N*-[4-([(dimethylamino)-sulfonyl]aminomethyl)cyclohexyl]methylthiourea: A solution of 1.0 g of *N'*-[4-(aminomethyl)cyclohexyl]-methyl-*N,N*-dimethylsulfamide in 25 ml of THF was treated with 0.7 g of benzoylisothiocyanate, stirred for 24 hours and the solvent removed to afford a crude product: $^1\text{H NMR}$ (CDCl_3) δ 8.20-7.60 (m, 5H), 3.80-3.30 (broad, 3H), 2.80-2.60 (m, 2H), 2.55-2.40 (m, 2H), 2.00-1.20 (m, 17H), 1.00-0.80 (m, 2H).

The crude product was treated with 15 ml of methanol and 15 ml of water containing 2.0 g of potassium carbonate. The mixture was stirred for 24 hours and then extracted with

ethyl acetate. The combined extracts were washed with saturated NaCl solution, dried (sodium sulfate) and the solvent removed under reduced pressure to give a yellow oil which was used in the next step without further purification.

N' - (4 - [(4,5-Dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-ylamino)methyl]cyclohexylmethyl) - *N,N*-dimethylsulfamide: 4-Bromo-7-fluoro-2,3,4,5-tetrahydro-1-benzothiepin-5-one (0.200 g, 0.720 mmole) and *N* - [4 - [(dimethylamino)sulfonyl]aminomethyl) - cyclohexyl]methylthiourea (0.250 g, 0.810 mmole) were heated at reflux for 18 hours, the solvent removed under reduced pressure and the crude product chromatographed (silica) to afford *N'* - (4 - [(4,5-dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-ylamino)methyl]cyclohexylmethyl) - *N,N*-dimethylsulfamide.

The following examples were prepared according to the reaction sequence shown in Schemes 12, 13 and 14 which describe the synthesis of carbamates and sulfonylureas:

Example 100

tert-Butyl *N* - { [4 - (4,5-Dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-ylamino)cyclohexyl]methyl } carbamate: 44% yield, 446 (ESMS, MH^+); 1H NMR (CD_3OD) δ 7.90 (dd, 1H, $J = 1.2, 7.8$ Hz), 7.49 (dd, 1H, $J = 0.8, 7.8$ Hz), 7.32 (apparent dt, 1H, $J = 1.4, 7.7$ Hz), 7.16 (apparent dt, 1H, $J = 1.3, 7.6$ Hz), 3.41 (m, 1H), 3.30 (m, 2H), 3.19 (t, 2H, $J = 6.5$ Hz), 3.06 (t, 2H, $J = 5.8$ Hz), 2.90 (d, 2H, $J = 7.0$ Hz), 1.99 (ABm, 4H), 1.43 (s, 9H), 1.32-1.05 (m, 3H).

Example 101

N' - [6- (4,5-Dihydrobenzo[2,3]thiepino[4,5-d] [1,3]thiazol-2-ylamino)hexyl] -N,N-diethylsulfamide:

5 72% yield, 470 (ESMS, MH^+); 1H NMR ($CDCl_3$) δ 8.05 (dd, 1H, $J=1.3$, 8.0 Hz), 7.52 (dd, 1H, $J=0.8$, 7.2 Hz), 7.33 (apparent dt, 1H, $J=0.8$, 8.0 Hz), 7.16 (apparent dt, 1H, $J=1.3$, 7.2 Hz), 5.05 (m, 1H), 4.02 (m, 1H), 3.26 (q, 4H, $J=7.1$ Hz), 3.29-3.22 (m, 2H), 3.20 (s, 2H), 2.97 (q, 2H, $J=6.7$ Hz), 1.67-1.37 (m, 10H), 1.81 (t, 6H, $J=7.1$ Hz).

Example 102

15 N' - { [4- (4,5-Dihydrobenzo[2,3]thiepino[4,5-d] [1,3]thiazol-2-ylamino)cyclohexyl]methyl} -N,N-diethylsulfamide: 84% yield, 481 (ESMS, MH^+); 1H NMR ($CDCl_3$) δ 8.05 (dd, 1H, $J=1.0$, 8.0 Hz), 7.52 (dd, 1H, $J=0.9$, 7.7 Hz), 7.33 (apparent dt, 1H, $J=1.0$, 7.7 Hz), 7.16 (apparent dt, 1H, $J=7.7$ Hz), 4.86 (m, 1H), 4.04 (m, 1H), 3.27 (q, 4H, $J=7.5$ Hz), 3.20 (m, 1H), 3.19 (s, 2H), 2.86 (t, 2H, $J=6.8$ Hz), 2.08 (ABm, 4H), 1.52 (m, 2H), 1.20 (t, 6H, $J=7.5$ Hz), 1.10 (m, 5H).

Example 103

25 N' - [5- (4,5-Dihydrobenzo[2,3]thiepino[4,5-d] [1,3]thiazol-2-ylamino)pentyl] -N,N-diethylsulfamide: 55% yield; Anal. Calc. for $C_{20}H_{30}N_4S_3O_2 + 0.1 CH_2Cl_2$: C, 52.12; H, 6.57; N, 12.10. Found: C, 52.13; H, 6.12; N, 11.96; 1H NMR ($CDCl_3$) δ 8.02 (dd, 1H, $J=0.6$, 7.5 Hz), 7.50 (dd, 1H, $J=0.6$, 7.5 Hz), 7.31 (t, 1H, $J=6.5$ Hz), 7.15 (t, $J=6.5$ Hz), 6.00 (broad, 1H), 4.85 (m, 1H), 3.27-3.11 (m, 10H), 2.87-2.85 (m, 2H), 1.52-1.18 (m, 12H).

Example 104

N' - [4 - (4,5-Dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-ylamino)butyl] -N,N-diethylsulfamide: 45% yield; Anal. Calc. for $C_{19}H_{28}N_4S_3O_2 + 0.2 CH_2Cl_2$: C, 50.39; H, 6.25; N 12.24. Found: C, 50.75; H, 6.17; N, 11.96. 1H NMR ($CDCl_3$) δ 7.99 (dd, 1H, J=0.6, 7.5 Hz), 7.50 (dd, 1H, J=0.6, 7.5 Hz), 7.31 (t, 1H, J=6.5 Hz), 7.15 (t, 1H, J=6.5 Hz), 6.17 (broad, 1H), 5.08 (m, 1H), 3.08-3.26 (m, 10H), 2.81 (m, 2H), 1.44-1.12 (m, 10H).

Example 105

tert-Butyl N-{5-[(9-fluoro-4,5-dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-yl)amino]pentyl}-carbamate: 60% yield; Anal. Calc. for $C_{21}H_{28}N_3F_2S_2O_2 + 0.15 CH_2Cl_2$: C, 56.41; H, 6.33; N, 9.30. Found : C, 56.45; H, 6.17; N, 8.90; 1H NMR ($CDCl_3$) δ 7.72 (dd, 1H, J=1.2, 7.5 Hz), 7.47-7.04 (m, 1H), 6.89-6.83 (m, 1H), 6.190-6.142 (m, 1H), 4.747-4.690 (m, 1H), 3.37-2.80 (m, 8H), 1.64-1.05 (m, 6H), 1.41 (s, 9H).

Example 106

N' - (4 - [(4,5-Dihydrobenzo[2,3]thiepino[4,5-d][1,3]thiazol-2-ylamino)methyl]cyclohexylmethyl) -N,N-dimethylsulfamide: 50% yield; Anal. Calc. For $C_{21}H_{29}F_1N_4S_3O_2$: C, 52.00; H, 6.0; N, 11.5. Found: C, 51.76; H, 6.41; N, 11.05; 1H NMR ($CDCl_3$) δ 7.70 (dd, 1H, J=4.3 Hz), 7.50-7.40 (m, 1H), 6.87-6.80 (m, 1H), 6.40-6.32 (broad, 1H), 5.20-5.05 (m, 1H), 3.20-3.00 (m, 6H), 2.90-2.82 (m, 2H), 2.78 (s, 6H), 1.80-1.22 (m, 8H), 1.00-0.78 (m, 2H).

II. Synthetic Methods for General Structures

A. Tricyclic Compounds

The examples described in Section IA are merely illustrative of the methods used to synthesize tricyclic compounds in Section A of the Summary Of The Invention and Detailed Description Of The Invention. Further compounds may be obtained utilizing methods shown in Schemes 1-6. The substituents in Schemes 1-6 are described in the Detailed Description.

It may be necessary to incorporate protection and deprotection strategies for substituents such as amino, amido, carboxylic acid, and hydroxyl groups in the synthetic methods described above to form tricyclic derivatives. Methods for protection and deprotection of such groups are well-known in the art, and may be found, for example in Green, T.W. and Wuts, P.G.M. (1991) Protection Groups in Organic Synthesis, 2nd Edition John Wiley & Sons, New York.

B. Bicyclic Compounds

The examples described in Section IB are merely illustrative of the methods used to synthesize bicyclic derivatives in Section B of the Summary Of The Invention and Detailed Description Of The Invention. Further derivatives may be obtained utilizing methods shown in Schemes 7-11. The substituents in Schemes 7-11 are described in the Detailed Description.

It may be necessary to incorporate protection and deprotection strategies for substituents such as amino, amido, carboxylic acid, and hydroxyl groups in the synthetic methods described above to form bicyclic derivatives. Methods for protection and deprotection of

such groups are well-known in the art, and may be found, for example in Green, T. W. and Wuts, P.G. M. (1991) Protection Groups in Organic Synthesis, 2nd Edition John Wiley & Sons, New York.

5

C. Tricyclic Compounds

The examples described in Section IC are merely illustrative of the methods used to synthesize tricyclic compounds in Section C of the Summary Of The Invention and Detailed Description Of The Invention. Further compounds may be obtained utilizing methods shown in Schemes 12-17. The substituents in Schemes 12-17 are described in the Detailed Description.

It may be necessary to incorporate protection and deprotection strategies for substituents such as amino, amido, carboxylic acid, and hydroxyl groups in the synthetic methods described above to form tricyclic derivatives. Methods for protection and deprotection of such groups are well-known in the art, and may be found, for example in Green, T.W. and Wuts, P.G.M. (1991) Protection Groups in Organic Synthesis, 2nd Edition John Wiley & Sons, New York.

III. Oral Compositions

As a specific embodiment of an oral composition of a compound of this invention, 100 mg of one of the compounds described herein is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.

IV. Pharmacological Evaluation of Compounds at Cloned Neuropeptide Y-type Receptors

The pharmacological properties of the compounds of the present invention were evaluated at one or more of the cloned human neuropeptide Y-type receptors Y1, Y2, Y4, and Y5, using protocols described below.

5

Cell Culture

COS-7 cells were grown on 150 mm plates in D-MEM with supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100 µg/ml streptomycin) at 37 °C, 5% CO₂. Stock plates of COS-7 cells were trypsinized and split 1:6 every 3-4 days. Human embryonic kidney 293 cells were grown on 150 mm plates in D-MEM with supplements (minimal essential medium) with Hanks' salts and supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100 µg/ml streptomycin) at 37 °C, 5% CO₂. Stock plates of 293 cells were trypsinized and split 1:6 every 3-4 days. Mouse fibroblast LM(tk-) cells were grown on 150 mm plates in D-MEM with supplements (Dulbecco's Modified Eagle Medium with 10% bovine calf serum, 4 mM glutamine, 100 units/mL penicillin/100 µg/mL streptomycin) at 37 °C, 5% CO₂. Stock plates of LM(tk-) cells were trypsinized and split 1:10 every 3-4 days.

25

LM(tk-) cells stably transfected with the human Y5 receptor were routinely converted from an adherent monolayer to a viable suspension. Adherent cells were harvested with trypsin at the point of confluence, resuspended in a minimal volume of complete DMEM for a cell count, and further diluted to a concentration of 10⁶ cells/ml in suspension media (10% bovine calf serum, 10% 10X Medium 199 (Gibco), 9 mM NaHCO₃, 25 mM glucose, 2 mM L-glutamine, 100 units/ml penicillin/100 µg/ml

30

195

streptomycin, and 0.05% methyl cellulose). The cell suspension was maintained in a shaking incubator at 37 °C, 5% CO₂ for 24 hours. Membranes harvested from cells grown in this manner may be stored as large, uniform batches in liquid nitrogen. Alternatively, cells may be returned to adherent cell culture in complete DMEM by distribution into 96-well microtiter plates coated with poly-D-lysine (0.01 mg/ml) followed by incubation at 37 °C, 5% CO₂ for 24 hours. Cells prepared in this manner yielded a robust and reliable NPY-dependent response in cAMP radio-immunoassays as further described hereinbelow.

Mouse embryonic fibroblast NIH-3T3 cells were grown on 150 mm plates in Dulbecco's Modified Eagle Medium (DMEM) with supplements (10% bovine calf serum, 4 mM glutamine, 100 units/ml penicillin/100 µg/ml streptomycin) at 37 °C, 5% CO₂. Stock plates of NIH-3T3 cells were trypsinized and split 1:15 every 3-4 days.

Sf9 and Sf21 cells were grown in monolayers on 150 mm tissue culture dishes in TMN-FH media supplemented with 10% fetal calf serum, at 27 °C, no CO₂. High Five insect cells were grown on 150 mm tissue culture dishes in Ex-Cell 400™ medium supplemented with L-Glutamine, also at 27 °C, no CO₂.

Transient Transfection

All receptor subtypes studied (human and rat Y1, human and rat Y2, human and rat Y4, human and rat Y5) were transiently transfected into COS-7 cells by the DEAE-dextran method, using 1 µg of DNA /10⁶ cells (Cullen, 1987). The human Y1 receptor was prepared using known methods (Larhammar, et al., 1992).

Stable Transfection

Human Y1, human Y2, and rat Y5 receptors were co-transfected with a G-418 resistant gene into the human embryonic kidney 293 cell line by a calcium phosphate transfection method (Cullen, 1987). Stably transfected cells were selected with G-418. Human Y4 and human Y5 receptors were similarly transfected into mouse fibroblast LM(tk-) cells and NIH-3T3 cells.

Binding of the compounds of the present invention to human Y1, Y2, Y4, and Y5 receptors was evaluated using stably transfected 293 or LM(tk-) cells as described above. Stably transfected cell lines which may be used for binding assays include, for example, for the human Y1 receptor, 293-hY1-5 (deposited June 4, 1996, under ATCC Accession No. CRL-12121), for the human Y2 receptor, 293-hY2-10 (deposited January 27, 1994, under ATCC Accession No. CRL-11537), for the human Y4 receptor, L-hY4-3 (deposited January 11, 1995, under ATCC Accession No. CRL-11779), and for human Y5 receptor, L-hY5-7 (deposited November 15, 1995, under ATCC Accession No. CRL-11995). These cell lines were deposited with the American Type Culture Collection (ATCC), 10801 University Blvd., Manassas, Virginia 20110-2209, U.S.A. under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

Membrane Harvest

Membranes were harvested from COS-7 cells 48 hours after transient transfection. Adherent cells were washed twice in ice-cold phosphate buffered saline (138 mM NaCl, 8.1 mM Na₂HPO₄, 2.5 mM KCl, 1.2 mM KH₂PO₄, 0.9 mM CaCl₂, 0.5 mM MgCl₂, pH 7.4) and lysed by sonication in ice-cold

197

sonication buffer (20 mM Tris-HCl, 5 mM EDTA, pH 7.7). Large particles and debris were cleared by low speed centrifugation (200 x g, 5 min, 4 °C). Membranes were collected from the supernatant fraction by centrifugation (32,000 x g, 18 min, 4 °C), washed with ice-cold hypotonic buffer, and collected again by centrifugation (32,000 x g, 18 min, 4 °C). The final membrane pellet was resuspended by sonication into a small volume of ice-cold binding buffer (-1 ml for every 5 plates: 10 mM NaCl, 20 mM HEPES, 0.22 mM KH₂PO₄, 1.26 mM CaCl₂, 0.81 mM MgSO₄, pH 7.4). Protein concentration was measured by the Bradford method (Bradford, 1976) using Bio-Rad Reagent, with bovine serum albumin as a standard. Membranes were held on ice for up to one hour and used fresh, or flash-frozen and stored in liquid nitrogen.

Membranes were prepared similarly from 293, LM(tk-), and NIH-3T3 cells. To prepare membranes from baculovirus infected cells, 2 x 10⁷ Sf21 cells were grown in 150mm tissue culture dishes and infected with a high-titer stock of hY5BB3. Cells were incubated for 2-4 days at 27 °C, no CO₂ before harvesting and membrane preparation as described above.

Membranes were prepared similarly from dissected rat hypothalamus. Frozen hypothalami were homogenized for 20 seconds in ice-cold sonication buffer with the narrow probe of a Virtishear homogenizer at 1000 rpm (Virtis, Gardiner, NY). Large particles and debris were cleared by centrifugation (200 x g, 5 min, 4 °C) and the supernatant fraction was reserved on ice. Membranes were further extracted from the pellet by repeating the homogenization and centrifugation procedure two more times. The supernatant fractions were pooled and subjected to high

198

speed centrifugation (100,000 x g, 20 min. 4 °C).
The final membrane pellet was resuspended by gentle
homogenization into a small volume of ice-cold binding
buffer (1 mL/gram wet weight tissue) and held on ice for
up to one hour, or flash-frozen and stored in liquid
nitrogen.

Radioligand Binding to Membrane Suspensions

Membrane suspensions were diluted in binding buffer
supplemented with 0.1% bovine serum albumin to yield an
optimal membrane protein concentration so that ^{125}I -PYY (or
alternative radioligand such as ^{125}I -NPY, ^{125}I -PYY₃₋₃₆, or
 ^{125}I -[Leu³¹Pro³⁴]PYY) bound by membranes in the assay was
less than 10% of ^{125}I -PYY (or alternative radioligand)
delivered to the sample (100,000 dpm/sample = 0.08 nM for
competition binding assays). ^{125}I -PYY (or alternative
radioligand) and peptide competitors were also diluted to
desired concentrations in supplemented binding buffer.
Individual samples were then prepared in 96-well
polypropylene microtiter plates by mixing ^{125}I -PYY (25 μL)
(or alternative radioligand), competing peptides or
supplemented binding buffer (25 μL), and finally, membrane
suspensions (200 μL). Samples were incubated in a 30 °C
water bath with constant shaking for 120 min. Incubations
were terminated by filtration over Whatman GF/C filters
(pre-coated with 1% polyethyleneimine and air-dried before
use), followed by washing with 5 mL of ice-cold binding
buffer. Filter-trapped membranes were impregnated with
MeltiLex solid scintillant (Wallac, Turku, Finland) and
counted for ^{125}I in a Wallac Beta-Plate Reader.
Alternatively, incubations were carried out in GF/C filter
plates (pre-coated with 1% polyethyleneimine and air-dried
before use), followed by vacuum filtration and three
washes of 300 μL of ice-cold binding buffer. 50 μL of

199

UltimaGold (Packard) scintillant were added and counted for ^{125}I in a Wallac MicroBeta Trilux. Non-specific binding was defined by 300 nM human NPY for all receptors except the Y4 subtypes; 100 nM human PP was used for the human Y4 and 100 nM rat PP for the rat Y4. Specific binding in time course and competition studies was typically 80%; most non-specific binding was associated with the filter. Binding data were analyzed using nonlinear regression and statistical techniques available in the GraphPAD Prism package (San Diego, CA).

Functional Assay: Radioimmunoassay of cAMP

Stably transfected cells were seeded into 96-well microtiter plates and cultured until confluent. To reduce the potential for receptor desensitization, the serum component of the media was reduced to 1.5% for 4 to 16 hours before the assay. Cells were washed in Hank's buffered saline, or HBS (150 mM NaCl, 20 mM HEPES, 1 mM CaCl_2 , 5 mM KCl, 1 mM MgCl_2 , and 10 mM glucose) supplemented with 0.1% bovine serum albumin plus 5 mM theophylline and pre-equilibrated in the same solution for 20 min at 37 °C in 5% CO_2 . Cells were then incubated 5 min with 10 μM forskolin and various concentrations of receptor-selective ligands. The assay was terminated by the removal of HBS and acidification of the cells with 100 mM HCl. Intracellular cAMP was extracted and quantified with a modified version of a magnetic bead-based radioimmunoassay (Advanced Magnetix, Cambridge, MA). The final antigen/antibody complex was separated from free ^{125}I -cAMP by vacuum filtration through a PVDF filter in a microtiter plate (Millipore, Bedford, MA). Filters were punched and counted for ^{125}I in a Packard gamma counter. Binding data were analyzed using nonlinear regression and

200

statistical techniques available in the GraphPAD Prism package (San Diego, CA).

Functional Assay: Intracellular calcium mobilization

5 The intracellular free calcium concentration was measured by microspectrofluorimetry using the fluorescent indicator dye Fura-2/AM. Stably transfected cells were seeded onto a 35 mm culture dish containing a glass coverslip insert. 10 Cells were washed with HBS and loaded with 100 μ l of Fura-2/AM (10 μ M) for 20 to 40 min. After washing with HBS to remove the Fura-2/AM solution, cells were equilibrated in HBS for 10 to 20 min. Cells were then visualized under the 40X objective of a Leitz Fluovert FS microscope and 15 fluorescence emission was determined at 510 nm with excitation wave lengths alternating between 340 nm and 380 nm. Raw fluorescence data were converted to calcium concentrations using standard calcium concentration curves and software analysis techniques.

Materials

20 Cell culture media and supplements were from Specialty Media (Lavallette, NJ). Cell culture plates (150 mm and 96-well microtiter) were from Corning (Corning, NY). Sf9, 25 Sf21, and High Five insect cells, as well as the baculovirus transfer plasmid, pBlueBacIIITM, were purchased from Invitrogen (San Diego, CA). TMN-FH insect medium complemented with 10% fetal calf serum, and the baculovirus DNA, BaculoGoldTM, was obtained from Pharmingen 30 (San Diego, CA.). Ex-Cell 400TM medium with L-Glutamine was purchased from JRH Scientific. Polypropylene 96-well microtiter plates were from Co-star (Cambridge, MA). All radioligands were from New England Nuclear (Boston, MA). Commercially available NPY and related peptide analogs 35 were either from Bachem California (Torrance, CA) or Peninsula (Belmont, CA); [D-Trp³²]NPY and PP C-terminal

201

fragments were synthesized by custom order from Chiron
Mimotopes Peptide Systems (San Diego, CA). Bio-Rad Reagent
was from Bio-Rad (Hercules, CA). Bovine serum albumin
(kdfjas;jflka;ljf;dalkjf;lkajf;lka(ultra-fat free, A-7511)
5 was from Sigma (St. Louis. MO).1 other materials were
reagent grade.

Radioligand Binding Assay Results

10 The compounds described above were assayed using cloned
human NPY receptors. The preferred compounds were found
to be selective NPY (Y5) antagonists. The binding
affinities of several compounds for NPY (Y5) are
illustrated in Tables 1-3.

15

20

25

30

Table 1








EXAMPLE No.	STRUCTURE	K _i , nM hNPY-5	K _i , nM hNPY-1,2,4
1		2.1	>10000
2		1.6	
3		4.5	>10000
4		19.5	>10000
5		19.8	>10000
6		3.1	
7		23	>10000

Table 1 continued








EXAMPLE No.	STRUCTURE	K _i , nM hNPY-5	K _i , nM hNPY-1, 2, 4
8		94	
9		15.4	>10000
10		2.4	>10000
11		6.4	>10000
12		6.1	>1000
13		132	
14		2.1	>10000

Table 1 continued

EXAMPLE No.	STRUCTURE	K _i , nM hNPY-5	K _i , nM hNPY-1,2,4
15		3.2	
16		4.7	>10000
17		11	>10000
18		7.1	>10000
19		5	>10000
20		5.8	>10000
21		2.5	>10000

Table 1 continued








EXAMPLE No.	STRUCTURE	K_i , nM hNPY-5	K_i , nM hNPY-1, 2, 4
22		2.9	>10000
23		15.5	>10000
24		3.5	>10000
25		2.9	>10000
26		13.1	>10000
27		9.2	>10000
28		4	>10000

Table 1 continued








EXAMPLE No.	STRUCTURE	K _i , nM hNPY-5	K _i , nM hNPY-1,2,4
29		10.3	>10000
30		4.2	>10000
31		3.9	>10000
32		2.9	>10000
33		6.8	>10000
34		34	
35		6.5	>10000

Table 1 continued








EXAMPLE No.	STRUCTURE	K_i , nM hNPY-5	K_i , nM hNPY-1,2,4
36		60	
37		7.4	>10000
38		15.8	>10000
39		4.8	>10000
40		19.5	>10000
41		30	
42		3.9	>10000

Table 1 continued








EXAMPLE No.	STRUCTURE	K _i , nM hNPY-5	K _i , nM hNPY-1,2,4
43		4.4	>10000
44		2.3	>10000
45		65	>10000
46		34	>10000
47		22	>10000
48		15.1	>10000
49		2.9	

Table 1 continued








EXAMPLE No.	STRUCTURE	K _i , nM hNPY-5	K _i , nM hNPY-1,2,4
50		15.3	
51		8.3	
52		1.7	
53		3.2	
54		11.9	>10000
55		3.4	
56		26	

Table 1 continued








EXAMPLE No.	STRUCTURE	K_i , nM hNPY-5	K_i , nM hNPY-1, 2, 4
57		66	
58		3	
59		2.3	
60		31	
61		162	
62		1.8	
63		31	

Table 1 continued








EXAMPLE No.	STRUCTURE	K _i , nM hNPY-5	K _i , nM hNPY-1, 2, 4
64		12.2	
65		11.6	
66		2	
67		5.5	
68		4	
69		4.1	
70		21	

Table 1 continued








EXAMPLE No.	STRUCTURE	K _i , nM hNPY-5	K _i , nM hNPY-1,2,4
71		2.5	
72		22	
73		8.7	
74		37	
75		36	
76		4.1	
77		14.6	

Table 1 continued








EXAMPLE No.	STRUCTURE	K_i , nM hNPY-5	K_i , nM hNPY-1, 2, 4
78		6	
79		20	
80		33	
81		62	
82		11.9	
83		12	
84		40	

Table 1 continued







EXAMPLE No.	STRUCTURE	K_i , nM hNPY-5	K_i , nM hNPY-1,2,4
85		150	
86		68	
87		86	
88		36	
89		9.2	
90		5.7	

Table 2








EXAMPLE No.	STRUCTURE	K_i , nM hNPY-5	K_i , nM hNPY-1, 2, 4
91		12.3	
92		912	
93		150	
94		240	
95		106	
96		56	
97		335	

Table 2 continued











EXAMPLE No.	STRUCTURE	K_i , nM hNPY-5	K_i , nM hNPY-1, 2, 4
98		125	
99		45	

Table 3


EXAMPLE	STRUCTURE	K _i , nM hNPY-5	K _i , nM hNPY-1,2,4
100		24	
101		5.9	
102		5.2	>10000
103		3.7	>10000
104		3.2	>10000
105		91	>10000
106		2.6	

Scheme 1A. Synthesis of Thioureas




- a. benzoylisothiocyanate
b. K_2CO_3 , MeOH


A =




Scheme 1B. Synthesis of Thioureas




- a. Benzoylisothiocyanate
b. K_2CO_3 , MeOH
c. alkyl halide or acyl halide followed by borane reduction
d. formylating agent such as 1H-benzotriazole-1-carboxaldehyde
e. HCl or TFA




Scheme 1C. Synthesis of Thioureas




- a. Benzoylisothiocyanate
 b. K_2CO_3 , MeOH
 c. alkyl halide or acyl halide followed by borane reduction
 d. R_{12}COCl
 e. HCl or TFA




Scheme 1D. Synthesis of Thioureas



- a. benzoylisothiocyanate
 b. K_2CO_3 , MeOH
 c. alkyl halide or acyl halide followed by borane reduction
 d. R_{19}COCl
 e. HCl or TFA






Scheme 1E. Synthesis of Thioureas



- a. benzoylisothiocyanate
b. K_2CO_3 , MeOH
f. HNR_3R_4
g. reduction


Scheme 2A. Synthesis of Bromoketones




L = leaving group such as Br

[illegible]

Scheme 2B. Synthesis of Bromoketones




Scheme 2C. Synthesis of Bromoketones





224

Scheme 3A. Synthesis of the Tricycles




Scheme 3B. Synthesis of the Tricycles






b. TFA or HCl




A =






Scheme 3C. Synthesis of the Tricycles





Scheme 3D. Synthesis of the Tricycles




Scheme 3E. Synthesis of the Tricycles




Scheme 3F. Synthesis of the Tricycles




Scheme 3G. Synthesis of the Tricycles





Scheme 4A. Synthesis of Tricycles





Scheme 4B. Synthesis of Tricycles




Scheme 5. Transamination of Sulfonylureas




A =





Scheme 6: Synthesis of Side Chains




a. Diphenylphosphoryl azide, triethylamine, toluene;
b. heat; c. HOCH_2Ph

Scheme 7A. Synthesis of Thioureas

- a. benzoylisothiocyanate
b. K_2CO_3 , MeOH


**Scheme 7B. Synthesis of Thioureas**

- a. Benzoylisothiocyanate
b. K_2CO_3 , MeOH
c. alkyl halide or acyl halide followed by borane reduction
d. formylating agent such as 1H-benzotriazole-1-carboxaldehyde
e. HCl or TFA



233

Scheme 7C. Synthesis of Thioureas




a. Benzoylisothiocyanate



b. K_2CO_3 , MeOH

c. alkyl halide or acyl halide followed by borane reduction



d. R_{12}COCl

e. HCl or TFA





Scheme 8A. Synthesis of Bromoketones**Scheme 8B. Synthesis of Chloroketones**



Scheme 9A. Synthesis of the Bicycles




b. TFA or HCl






Scheme 9B. Synthesis of the Bicycles






b. TFA or HCl





Scheme 9C. Synthesis of the Bicycles



Scheme 9D. Synthesis of the Bicycles



Scheme 9E. Synthesis of the Bicycles




b. TFA or HCl

c. $RCOCl$



d. reduction

e. formylating agent such as
1H-benzotriazole-1-carboxaldehyde



239


Scheme 9F. Synthesis of the Bicycles




b. TFA or HCl

c. $R_{12}I$


d. formylating agent such as
1H-benzotriazole-1-carboxaldehyde




Scheme 10A. Synthesis of Bicycles




Scheme 10B. Synthesis of Bicycles





Scheme 11: Synthesis of Side Chains

a. Diphenylphosphoryl azide, triethylamine, toluene;
b. heat; c. HOCH2Ph

Scheme 12A. Synthesis of Thioureas


- a. benzoylisothiocyanate
b. K_2CO_3 , MeOH

A =


**Scheme 12B. Synthesis of Thioureas**

- a. Benzoylisothiocyanate
b. K_2CO_3 , MeOH
c. alkyl halide or acyl halide followed by borane reduction.
d. formylating agent such as 1H-benzotriazole-1-carboxaldehyde
e. HCl or TFA

A =



Scheme 12C. Synthesis of Thioureas²⁴³




a. Benzoylisothiocyanate

b. K_2CO_3 , MeOH


c. alkyl halide or acyl halide followed by borane reduction

d. R_{17}COCl

e. HCl or TFA



Scheme 12D. Synthesis of Thioureas




a. Benzoylisothiocyanate

b. K_2CO_3 , MeOH


c. alkyl halide or acyl halide followed by borane reduction

d. R_{12}COCl

e. HCl or TFA




Scheme 12E. Synthesis of Thioureas




- a. benzoylisothiocyanate
b. K_2CO_3 , MeOH
f. HNR_3R_4
g. reduction

Scheme 13. Synthesis of Bromoketones




L = leaving group such as Br
 X = S, SO, SO₂
 DMD = dimethyldioxirane
 mCPBA = m-chloroperbenzoic acid







Scheme 14A. Synthesis of the Tricycles





Scheme 14B. Synthesis of the Tricycles








247



Scheme 14C. Synthesis of the Tricycles**Scheme 14D. Synthesis of the Tricycles**

Scheme 14E. Synthesis of the Tricycles





Scheme 14F. Synthesis of the Tricycles





Scheme 15A. Synthesis of Tricycles**Scheme 15B. Synthesis of Tricycles**


Scheme 16. Transamination of Sulfonylureas



A =



Scheme 17: Synthesis of Side Chains



a. Diphenylphosphoryl azide, triethylamine, toluene;
 b. heat; c. HOCH₂Ph

Functional Assay Results

The functional in vitro activity of several compounds were characterized using a radioimmunoassay of cAMP, the results of which are summarized in Table 4.

Table 4. Functional Antagonism Data

Example #	K _i (h NPY-5), nM	pK _b
1	2.1	7.9
3	4.5	8.5
4	19.5	8.6
11	6.4	8.5
15	3.2	8.6
20	0.3	8.3
34	33.5	7.2
43	4.4	7.8
44	2.3	8.8
50	15.3	8.1
79	20.1	7.9
80	32.9	7.4

References

5 Balasubramaniam, A., Sheriff, S., Johnson, M. E.,
Prabhakaran, M., Huang, Y., Fischer, J. E., and Chance, W.
T. (1994). [D-Trp³²]Neuropeptide Y: A competitive
antagonist of NPY in rat hypothalamus. J. Med. Chem. 37: 3

10 Bradford, M. M. (1976). A rapid and sensitive method for
the quantitation of microgram quantities of protein
utilizing the principle of protein-dye binding. Anal.
Biochem. 72: 248-254.

15 Chabaka, L. M., et al., "Facile Synthesis of 2-Furyl-, 2-
Pyrrolyl-, 2-Imidazolyl- and Pyrrolo- Azoles from 2-
Substituted Methylazoles." Pol. J. Chem. (1994) 68(7):
1317-1325.

20 Clark, J. T., Kalra, P. S., Crowley, W. R., and Kalra, S.
P. (1984). Neuropeptide Y and human pancreatic
polypeptide stimulate feeding behavior in rats.
Endocrinology 115: 427-429.

25 Crangk, G. and Foulis, M.J., " Oxazoles from ureas" J.
Med. Chem. (1971) 14: 1075.

30 Criscione, L., Rigollier, P., Batzl-Hartmann, C., Rueger,
H., Stricker-Krongrad, A., Wyss, P., Brunner, L.,
Whitebread, S., Yamaguchi, Y., Gerald, C., Heurich, R. O.,
Walker, M. W., Chiesi, M., Schilling, W., Hofbauer, K. G.,
Levens, N. (1998) Food intake in free-feeding and energy-
deprived lean rats is mediated by the neuropeptide Y5
receptor. J. Clin. Invest. 102(12): 2136-45.

Critcher, D.J. and Pattenden, G., " Synthetic Studies Towards Pateamine, a Novel Thiazole-Based 19- Membered Bis-lactone from Mycale sp." Tetrahedron. Lett. (1996) 37(50): 9107-9110.

Cullen, B. (1987). Use of eukaryotic expression technology in the functional analysis of cloned genes. Methods Enzymol. 152: 685-704.

De Kimpe, N., et al., "Synthesis of 2-Imino-4-thiazolines, 2-Imino-4-alkoxythiazolidines, Thiazoles, and 4-Imidazolin-2-ones from alpha-Halomethyl Ketimines.", J. Heterocycl. Chem. (1996) 33(4): 1179-1183.

Demchenko, A. M., et al., " Preparation and Application of alpha -Bromomono- and -bisdifluoromethoxyacetophenones in the Course of Synthesis of Polymethyleneimidazoles Containing a Bridge Nitrogen Atom", Khim. Geterotsikl. Soedin. (1997) 10: 1371-1376.

Di Fabio, R. and Pentassuglia, G., "Novel Synthesis of Ethyl 3-(Bromoacetyl)-4,6-dichloro-1H-indole-2-carboxylate as Useful Intermediate in the Preparation of Potential Glycine Site Antagonists", Synth. Commun. (1998) 28(1): 51-60.

Dryden, S., Frankish, H., Wang, Q., and Williams, G. (1994). Neuropeptide Y and energy balance: one way ahead for the treatment of obesity? Eur. J. Clin. Invest. 24: 293-308.

Dumont, Y., Martel, J.-C., Fournier, A., St-Pierre, S., and Quirion, R. (1992). Neuropeptide Y and neuropeptide Y

receptor subtypes in brain and peripheral tissues.
Progress in Neurobiology 38: 125-167.

5 Eva, C., Oberto, A., Sprengel, R. and Genazzani, E.
(1992). The murine NPY-1 receptor gene: structure and
delineation of tissue specific expression. FEBS lett.
314: 285-288.

10 Eva, C., Keinänen, K., Monyer, H., Seeburg, P., and
Sprengel, R. (1990). Molecular cloning of a novel G
protein-coupled receptor that may belong to the
neuropeptide receptor family. FEBS Lett. 271: 80-84.

15 Friedman, B. S., et al., "Thiazoles from thioamides", J.
Am. Chem. Soc. (1937) 59: 2262.

Hammar, W.J. and Rustad, M.A., "Oxazoles from alpha-
bromoketones" J. Heterocycl. Chem. (1981) 18: 885.

20 Herzog, H., Hort, Y. J., Ball, H. J., Hayes, G., Shine,
J., and Selbie, L. (1992). Cloned human neuropeptide Y
receptor couples to two different second messenger
systems. Proc. Natl. Acad. Sci. USA 89: 5794-5798.

25 Kalra, S. P., Dube, M. G., Fournier, A., and Kalra, P. S.
(1991). Structure-function analysis of stimulation of
food intake by neuropeptide Y: Effects of receptor
agonists. Physiology & Behavior 50: 5-9.

30 Kearney, P. C., et al., "Solid-Phase Synthesis of 2-
Aminothiazoles", J. Org. Chem. (1998) 63(1): 196-200.

Larhammar, D., Blomqvist, A. G., Yee, F., Jazin, E., Yoo,
H., and Wahlestedt, C. (1992). Cloning and functional

expression of a human neuropeptide Y/peptide YY receptor of the Y1 type. J. Biol. Chem. 267: 10935-10938.

Levine, A. S., and Morley, J. E. (1984). Neuropeptide Y: A potent inducer of consummatory behavior in rats. Peptides 5: 1025-1029.

Little, T. L. and Webber, S. E., "A Simple and Practical Synthesis of 2-Aminoimidazoles" J. Org. Chem. (1994) 59(24): 7299-7305.

Marchetti, E., et al., "Oxazoles from ureas" J. Med. Chem. (1968) 11: 1092.

Michel, M.C. (1991). Receptors for neuropeptide Y: multiple subtypes and multiple second messengers. Trends Pharmacol. 12: 389-394.

Nagao, Y., et al., "Novel Nonprostanoid Prostacyclin (PGI₂) Mimetics with Heterocyclic Moiety", Heterocycles (1996) 42(2): 517-523.

Novikova, A. P., et al., "Synthesis and Properties of 1,3,4-Thiadiazine Derivatives. Part 1. Condensation of Substituted Phenacyl Bromides and Bromoacetylpyridines with Thiosemicarbazide", Khim. Geterotsikl. Soedin. (1991) (6): 843-846.

Pathak, V.N., et al., "Synthesis of Some New Fluorine Containing Oxazoles, Oxadiazoles, Thiadiazoles and Triazines"; J. Indian Chem. Soc. (1993) 70(6): 539-542.

Plazzi, P.V., et al., " Heteroaryl aminoethyl and Heteroarylthioethylimidazoles. Synthesis and H₃-Receptor Affinity", Eur. J. Med. Chem. (1995) 30(11): 881-889.

Sahu, A., and Kalra, S.P. (1993). Neuropeptidergic regulation of feeding behavior (neuropeptide Y). Trends Endocrinol. Metab. 4: 217-224.

5

Stanley, B. G., and Leibowitz, S. F. (1984). Neuropeptide Y: Stimulation of feeding and drinking by injection into the paraventricular nucleus. Life Sci. 35: 2635-2642.

10

Stanley, B. G., Magdalin, W., Seirafi, A., Nguyen, M. M., and Leibowitz, S. F. (1992). Evidence for neuropeptide Y mediation of eating produced by food deprivation and for a variant of the Y₁ receptor mediating this peptide's effect. Peptides 13: 581-587.

15

Wahlestedt, C., Edvinsson, L., Ekblad, E., and Hakanson, R. Effects of neuropeptide Y at sympathetic neuroeffector junctions: Existence of Y₁ and Y₂ receptors. In: *Neuronal messengers in vascular function*, Fernstrom Symp. No 10., pp. 231-242. Eds A. Nobin and C.H. Owman. Elsevier: Amsterdam (1987).

20

Wahlestedt, C., and Reis, D. J. (1993). Neuropeptide Y-Related Peptides and Their Receptors--Are the Receptors Potential Therapeutic Targets? Ann. Rev. Pharmacol. Tox. 32: 309-352.

25

Zhao, Z., et al., "Synthesis of trans-4-Alkenyloxazoles" Tetrahedron. Lett. (1991) 32(13): 1609-1612.